

# Does the Hippocampal Atrophy Correlate With the Cortical Theta Power in Elderly Subjects With a Range of Cognitive Impairment?

Martin Grunwald,\* Anke Hensel,† Henrike Wolf,‡ Thomas Weiss,‡ and Hermann-Josef Gertz†

**Summary:** A previous study with a small sample ( $N = 39$ ) showed a significant correlation between the cortical theta activity and the hippocampal volume in different stages of cognitive impairment in aged subjects. The recent study was aimed to replicate these results in a much bigger sample. The authors examined a sample of 121 right-handed subjects. The sample consisted of 37 healthy controls, 40 patients with questionable dementia, and 44 patients with mild dementia assessed by Clinical Dementia Rating. All subjects underwent EEG and brain MRI. Mean spectral power was calculated, and volume of hippocampal segments was measured. EEG theta power of the left and right hemisphere correlated significantly with the hippocampal volume on the left and right side in different stages of cognitive impairment. An increase of theta power was associated with decreased hippocampal volume. No other significant correlations were found for alpha or beta band power. No correlation was found between cortical theta and global brain volume. There seems to be a direct relationship between neuronal loss of the hippocampus and changed cortical theta activity for different stages of cognitive impairment in aged subjects.

**Key Words:** Cortical theta, EEG, Hippocampus, Mild cognitive impairment, Mild dementia.

(*J Clin Neurophysiol* 2007;24: 22–26)

Increased brain electrical theta activity and a decrease of hippocampal volume described two relevant parameters that discriminate patients with Alzheimer's disease (AD), mild cognitive impairment (MCI), and healthy controls (CO) (Förstl et al., 1996; Prichep et al., 1994). Functional relations seem to exist between cortical theta activity and the hippocampus (Miller 1991; Vinogradova 1995). A high correlation

between theta activity and an excessive neuronal loss in the hippocampi was found at autopsy in AD patients (Rae-Grant et al., 1987). Therefore, it can be presumed that a neuronal loss in the hippocampus may lead to changes in the slow-wave theta activity of the neocortex.

It was demonstrated in 39 subjects that the cortical spectral power density of the theta frequency band (4–8 Hz) within the EEG (theta power) under rest condition (eyes closed) is negatively correlated with the volume of the hippocampus in subjects with different degrees of cognitive impairment (MCI, CO, mild dementia). Such a negative correlation was found only for the theta power and not for other bands of the EEG spectra (Grunwald et al., 2001).

The aim of the present study was to replicate this examination in a bigger sample size of patients. According to the first study, we tested the hypothesis that a decrease of hippocampal volumes correlates negatively with an increase of cortical theta power over frontal regions under rest conditions (eyes closed). No correlations were expected for all other EEG bands. We also expect that the theta activity correlates specifically with the hippocampal volume and is not relative to the global brain volume.

## METHODS

### Participants

A total of 121 participants within aged 52 to 90 years were enrolled. Patients without dementia symptoms (53%) were recruited for this study from the Leipzig Longitudinal Study of the Aged (LEILA 75+) (Riedel-Heller et al. 2001). The remaining participants were recruited from referrals to the local memory clinic. To ensure that participants represented a cognitive continuum, they were recruited according to Mini-Mental State Examination strata (for more details see Wolf et al., 2004). Exclusion criteria were physical and/or neurologic disabilities (such as blindness, deafness, severe movement disorders or paralysis) that would have interfered with the subject's ability to complete neuropsychological tests or paraclinical examinations (such as computed tomography, EEG or MRI procedures).

All subjects were clinically assessed as described previously (Wolf et al., 2001). The degree of cognitive impairment was rated using the Clinical Dementia Rating (CDR) (Hughes et al., 1982) following a clinical interview with patient and a relative as described previously (Wolf et al.,

Department of Psychiatry, \*EEG Research Laboratory and †Memory Clinic, Interdisciplinary Center for Clinical Research, Leipzig, Germany; ‡Department of Biological Psychology, Friedrich-Schiller University Jena, Jena, Germany.

This paper was supported by the Bundesministerium für Bildung und Technologie (BMBF), by the Interdisciplinary Center for Clinical Research (IZKF) of the University of Leipzig (Projekt C8).

Address correspondence and reprint requests to Dr. Martin Grunwald, University of Leipzig, Department of Psychiatry, EEG-Research Lab, Johannisallee 34, 04103 Leipzig, Germany; e-mail: mgrun@medizin.uni-leipzig.de.

Copyright © 2007 by the American Clinical Neurophysiology Society  
ISSN: 0736-0258/07/2401-0022

**TABLE 1.** Characteristics of the Patient Groups

	<b>Controls</b> CDR 0.0 Mean (SD)	<b>QD</b> CDR 0.5 Mean (SD)	<b>MD</b> CDR 1.0 Mean (SD)
No. of subjects	37	40	44
Female/male	23/14	26/14	28/16
Age (y)	79.50 (2.83)	80.09 (3.68)	76.69 (6.09)
MMSE	28.2 (1.3)	25.6 (1.0)	22.4 (2.0)

CDR, Clinical Dementia Rating; SD, standard deviation; MMSE, Mini-Mental State Examination (Folstein 1975); QD, questionable dementia; MD, mild dementia.

2001). A CDR global score was assigned according to published rules. A CDR global score of 0 indicates no dementia, and CDR global scores of 0.5 to 1 indicate questionable and mild dementia. According to the CDR, the samples were divided into three groups: 37 controls (C, CDR = 0), 40 patients with questionable dementia (QD, CDR = 0.5), and 44 patients with mild dementia (MD, CDR = 1). All patients with a CDR global score of 1 were diagnosed with AD according to the research criteria of the ICD-10. This sample included 39 subjects from our preliminary sample (Grunwald et al., 2001). Demographic and neuropsychiatric characteristics of the groups are shown in Table 1. The study was approved by the local ethics committee. All subjects gave written informed consent to participate in this study.

### EEG Recording and Parameters

EEG recordings were performed  $102 \pm 169$  days before the MRI study. The EEG data recording procedure was similar to that described in a previous report (Grunwald et al., 2001). Segmentation of EEG data and subsequent calculations of the mean spectral power density were performed with the EEG analytical software package Brain Vision (Brain Products, Munich, Germany). Artifact-free segments of 1.53 seconds (256 samples/channel) of the rest period were chosen after automatic artifact rejection. For the artifact rejection, an amplitude criterion of  $\pm 70 \mu\text{V}$  was chosen. 150 to 250 artifact-free EEG-segments per subject under rest conditions were used for spectral analysis. The segments were submitted to a fast Fourier transform analysis. Mean spectral absolute power ( $\mu\text{V}^2$ ) was calculated as the mean amplitude of the spectral lines of the EEG bands (theta: 4.0–8.0 Hz, alpha: 8.0–13 Hz, beta: 13–24 Hz). The mean spectral power parameters per channel and per subject were  $\ln 10$ -transformed before statistical analysis.

### Magnetic Resonance Imaging

#### MRI Acquisition

Within 4 months after the clinical assessment, all participants were investigated by a 1.5 T tomograph (Siemens Vision) with a volumetric T1-weighted-MPRAGE sequence (TR 11.4 milliseconds, TE 4.4 milliseconds, 128 slices, orientation transverse, matrix  $256 \times 256$ , voxel size  $0.9 \times 0.9 \times 1.5$  mm).

### Analysis of MRI Data Sets

Datasets were analyzed using in-house software of the Max-Planck-Institute of Human Cognitive and Brain Sciences in Leipzig, Germany (Mon-A) without knowing the cognitive state or other clinical data of the participants.

The T1-weighted data sets were aligned with the stereotactical coordinate system and interpolated to an isotropical voxel size of 1 mm using fourth-order b-spline interpolation. Six cross sections of the hippocampus were manually outlined in the coronal plane on both sides (by A.H.). Hippocampal measures started behind the amygdala at the slice in which the area of the hippocampal head appeared maximal and were continued posteriorly at 3-mm intervals. The first two measured hippocampal segments are located in the anterior part of the hippocampus. The manual outlining technique of the hippocampus has previously been shown to have a high interrater reliability with an intraclass correlation coefficient of 0.996 (Wolf et al., 2001, 2004). To yield an estimate of the hippocampal volumes (HcV), areas one to six were multiplied by slice thickness (3 mm) and summed on each side (HcVL, resp. HcVR). The HcVR and HcVL per subject were  $\ln 10$ -transformed before statistical analysis.

In a large subsample of subjects ( $n = 104$ ), the brain volume was determined using a previously described automated region growing segmentation method (Wolf et al., 2003).

### Statistical Evaluation


All statistical analyses were conducted using the SPSS for Windows (version 10.0.7). To evaluate correlations between  $\ln 10$ -transformed EEG-data and  $\ln 10$ -transformed hippocampal volumes on each side throughout the entire sample ( $N = 121$ ), partial correlation coefficients of these parameters were computed and controlled for gender, age (Juarez and Corsi-Cabrera 1995, Coffey et al., 1998), and interval between MRI and EEG.

Eight single correlations per side (left, right) were calculated. Therefore, the critical alpha was adapted to  $\alpha' = 0.006$  ( $\alpha/8$ ). Because we used directed hypotheses, one-tailed tests were performed for the partial correlation.

To evaluate correlations between  $\ln 10$ -transformed theta power and  $\ln 10$ -transformed global brain volumes (BV), we used the data set of a subsample of 104 subjects. The partial correlations of 19 single tests were calculated ( $BV \times 19$  channel). The critical alpha was adapted to  $\alpha' = 0.002$  ( $\alpha/19$ ).

### RESULTS

Within the entire sample ( $N = 121$ ), partial correlation analysis (corrected for gender, age, time interval between MRI and EEG) between right hippocampal volume (HcVR) and spectral power of the theta band (right-side electrode positions revealed significant negative correlations [critical  $\alpha' < 0.006$ ] for frontal electrodes Fp2, F4; Fig 1). Partial correlations between left hippocampal volume (HcVL) and spectral power of the theta band (left side electrode positions) were significantly negative (critical  $\alpha' < 0.006$ ) for all left side electrodes (Fig 1). Partial correlation coefficients of all



**FIGURE 1.** Regression plot for significant linear correlations between right hippocampal volumes and right-frontal theta power (Fp2) and between left hippocampal volumes and left-frontal theta power (Fp1).

comparisons are shown in Table 2. For spectral power of the alpha band and beta band, no significant correlation coefficients were found.

No significant correlation was found between the theta power and the global brain volume (BV) in a subsample of  $n = 104$ .

## DISCUSSION

The main results of the present study are significant negative correlations between the theta power (left/right side) and the hippocampal volume (left/right side). This is in accordance with our previous results (Grunwald et al., 2001) that the increase of theta power at different stages of memory impairment is primarily associated with an atrophy of the hippocampus. Impressively, this study not only replicates the results of the previous study in a much bigger sample ( $N =$

**TABLE 2.** Partial Correlation Coefficients (Corrected for Age, Gender, and Time Interval Between MRI and EEG) Between Spectral Power of Theta Band ( $\mu V^2$ ) and Hippocampal Volume ( $mm^3$ )

Channel	Left Side		Channel	Right Side	
	r Value	P Value		r Value	P Value
Fp1	-0.270	0.002	Fp2	-0.270	0.002
F3	-0.260	0.002	F4	-0.236	0.005
F7	-0.276	0.001	F8	-0.135	0.072
C3	-0.263	0.002	C4	-0.188	0.020
P3	-0.265	0.002	P4	-0.184	0.023
T3	-0.272	0.001	T4	-0.150	0.052
T5	-0.257	0.002	T6	-0.159	0.042
O1	-0.282	0.001	O2	-0.192	0.018

*P* value is the one-tailed significance of partial correlation coefficients.

121), but also extends those results by significant negative correlations between size of the hippocampus and theta power for all left side electrodes. Two frontal electrodes on the right side reached the critical alpha level ( $\alpha' < 0.006$ ). However, the *r* values of the partial correlations are relatively low. The explained variance is on average 7%.

No significant correlation was found between theta power and the global brain volume in a subsample of 104 subjects. These results confirm the hypothesis that the relationship between cortical theta activity and the hippocampal volume is not mediated through the global brain volume.

We conclude for both hemispheres that theta activity is associated with hippocampal volume. A small hippocampal volume is correlated with a higher cortical theta power and *vice versa*. This result is specific for the theta activity because we have not found any significant correlations between hippocampal volume and alpha or beta power. This is in line with previous results (Grunwald et al., 2001).

It is well known that the theta activity in neocortical regions is induced by synchronized bursts of a small set of hippocampal pyramidal cells in the CA1 field, controlled by the medial and lateral septum (Bland et al., 1999; Vinogradova 1995) and regulated by hippocampo-cortical feedback loops (Grunwald et al., 2002; Klimesch et al. 1996; Lopes da Silva 1992; Miller 1991; Stewart and Fox, 1990). Although neuropathological mechanisms leading to an increase of cortical theta activity in humans at different stages of cognitive impairment are not completely understood, it seems comprehensible that the striking increase of synchronized hippocampal theta activity in these patients might reflect a process of degeneration, particularly of inhibitory interneurons in the CA1 field. Rae-Grant et al. (1987) provided some evidence for a possible connection between cortical theta activity and hippocampal atrophy. In their study, a high correlation between theta activity and an excessive neuronal loss in hippocampi was found at autopsy in AD patients. It might be assumed that the atrophy of the hippocampal formation is mainly due to a loss of neurons primarily in the CA1 region (Fukutani et al., 1995; Jack et al.,



2002; Price et al., 2001; Velez-Pardo et al., 2004; Wang et al., 2003), even in different stages of AD (Brady and Mufson 1991; Bobinski et al., 1995; Jack et al., 2000; Jobst et al., 1992, 1994).

A direct link between loss of CA1 neurons and changes of slow wave brain electrical activity was shown only in epileptic research. Diehl et al. (2002) showed relationships between brain electrical seizure spikes and cell loss in the hippocampal CA1 area in patients with hippocampal sclerosis. These investigators found a significant correlation between the amount of cell loss in CA1 and fronto-temporal spikes.

Whether the neuronal loss in the hippocampus is an epiphenomenon for the increase on theta activity or whether it is the causal reason cannot be distinguished on the basis of our data. The epiphenomenologic interpretation is that cortical theta activity can be generated independently of the hippocampus, e.g., in the cingulate cortex (Pizzagalli et al., 2003; Talk et al., 2004). This is in line with Kirk and Mackay's (2003) animal and human study findings that cortical theta activity is generated independently by two different networks, the hippocampo-anterior thalamic system and the perirhinal-mediodorsal thalamic system. Bastiaansen and Hagoort (2003) suggested that the cortical theta activity reflects an interplay between hippocampal and cortical generators whereby the theta generator(s) in the neocortex can produce theta activity by itself independently of or in relation to the hippocampal generators. We cannot exclude that the increased theta activity found for different forms of cognitive impairment in older subjects or in AD patients is caused by changes in other cortical regions and not by atrophy in the hippocampus.

A limitation of our study is the small spatial resolution of the EEG and the individual variation of electrode placements in relation to the brain anatomy. We used only the standard clinical resolution of the 10 to 20 system (19 electrodes). By this means, a strong spatial attribution between EEG signals and brain anatomy cannot be guaranteed. However, the cortical theta activity is not strongly focalized but rather involves broader cortical networks. Therefore, it seems reasonable to conclude that we were able to detect the related changes in the brain regions.

## ACKNOWLEDGMENTS

The authors thank I. Thomas for data collection, and U. Müller for his constructive comments on an earlier draft.

## REFERENCES

- American Psychiatric Association diagnostic and statistical manual of mental disorders, 3rd ed., rev. Washington, DC: American Psychiatric Press, 1987.
- Bannerman DM, Grubb M, Deacon RM, et al. Ventral hippocampal lesions affect anxiety but not spatial learning. *Behav Brain Res* 2003;139:197–213.
- Bannerman DM, Yee BK, Good MA, et al. Double dissociation of function within the hippocampus: a comparison of dorsal, ventral, and complete hippocampal cytotoxic lesions. *Behav Neurosci* 1999;113:1170–1188.
- Bast T, Feldon J. Hippocampal modulation of sensorimotor processes. *Prog Neurobiol* 2003;70:319–345.
- Bastiaansen M, Hagoort P. Event-induced theta responses as a window on the dynamics of memory. *Cortex* 2003;39:967–992.
- Berg L. Clinical dementia rating. *Br J Psychiatry* 1984;145:339.
- Bland BH, Oddie SD, Colom LV. Mechanisms of neural synchrony in the septohippocampal pathways underlying hippocampal theta generation. *J Neurosci* 1999;19:3223–3237.
- Bliss TV, Collingridge GL. A synaptic model of memory: long-term potentiation in the hippocampus. *Nature* 1993;361:31–39.
- Bobinski M, Wegiel J, Wisniewski HM, et al. Atrophy of hippocampal formation subdivisions correlates with stage and duration of Alzheimer disease. *Dementia* 1995;6:205–210.
- Brady DR, Mufson EJ. Alz-50 immunoreactive neuropil differentiates hippocampal complex subfields in Alzheimer's disease. *J Comp Neurol* 1991;305:489–507.
- Coffey CE, Lucke JF, Saxton JA, et al. Sex differences in brain aging. *Arch Neurol* 1998;55:169–179.
- Davis MH, Johnsruide IS. Hierarchical processing in spoken language comprehension. *J Neurosci* 2003;23:3423–3431.
- Diehl B, Najm I, Mohamed A, et al. Interictal EEG, hippocampal atrophy, and cell densities in hippocampal sclerosis and hippocampal sclerosis associated with microscopic cortical dysplasia. *J Clin Neurophysiol* 2002;19(2):157–62.
- Dolan RJ, Fletcher PF. Encoding and retrieval in human medial temporal lobes: an empirical investigation using functional magnetic resonance imaging (fMRI). *Hippocampus* 1999;9:25–34.
- Duva CA, Floresco SB, Wunderlich GR, et al. Disruption of spatial but not object-recognition memory by neurotoxic lesions of the dorsal hippocampus in rats. *Behav Neurosci* 1997;111:1184–1196.
- Fernandez A, Arrazola J, Maestu F, et al. Correlations of hippocampal atrophy and focal low-frequency magnetic activity in Alzheimer disease: volumetric MR imaging-magnetoencephalographic study. *Am J Neuroradiol* 2003;24:481–487.
- Floresco SB, Seamans JK, Phillips AG. Selective roles for hippocampal, prefrontal cortical, and ventral striatal circuits in radial-arm maze tasks with or without a delay. *J Neurosci* 1997;17:1880–1890.
- Folstein MF, Folstein SE, McHugh PR. "Mini-mental state." A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 1975;12:189–198.
- Förstl H, Besthorn C, Sattel H. Volumetrische Hirnveränderungen und quantitativen EEG bei normalem Altern und Alzheimer-Demenz. *Nervenarzt* 1996;67:53–61.
- Fukutani Y, Kobayashi K, Nakamura I, et al. Neurons, intracellular and extracellular neurofibrillary tangles in subdivisions of the hippocampal cortex in normal ageing and Alzheimer's disease. *Neurosci Lett* 1995;200:57–60.
- Giovanello KS, Schnyer DM, Verfaellie M. A critical role for the anterior hippocampus in relational memory: evidence from an fMRI study comparing associative and item recognition. *Hippocampus* 2004;14:5–8.
- Grunwald M, Busse F, Hensel A, et al. Correlation between cortical theta activity and hippocampal volumes in health, mild cognitive impairment, and mild dementia. *J Clin Neurophysiol* 2001;18:178–184.
- Grunwald M, Busse F, Hensel A, et al. Theta-power differences in patients with mild cognitive impairment under rest condition and during haptic tasks. *Alzheimer Dis Assoc Disord* 2002;16:40–48.
- Grunwald T, Lehnertz K, Heinze HJ, et al. Verbal novelty detection within the human hippocampus proper. *Proc Natl Acad Sci U S A* 1998;95:3193–3197.
- Hock BJ Jr, Bunsey MD. Differential effects of dorsal and ventral hippocampal lesions. *J Neurosci* 1998;18:7027–7032.
- Hojjatolleslami SA, Kruggel F, von Cramon DY. Segmentation of white matter lesions from volumetric MR images. In: Taylor C, Colchester A, eds. *Medical image computing and computer-aided intervention*. Heidelberg: Springer, 1999:52–61.
- Hughes CP, Berg L, Danziger WL, et al. A new clinical scale for the staging of dementia. *Br J Psychiatry* 1982;140:566–572.
- Izaki Y, Takita M, Nomura M, Akema T. Effects of ventral hippocampal long-term potentiation and depression on the gamma-band local field potential in anesthetized rats. *Exp Brain Res* 2004;157:147–151.
- Jack CR Jr, Dickson DW, Parisi JE, et al. Antemortem MRI findings correlate with hippocampal neuropathology in typical aging and dementia. *Neurology* 2002;58:750–757.
- Jack CR Jr, Petersen RC, Xu Y, et al. Rates of hippocampal atrophy correlate

- with change in clinical status in aging and AD. *Neurology* 2000;55:484–489.
- Jack CR, Petersen RC, Xu YC, et al. Medial temporal atrophy on MRI in normal aging and very mild Alzheimer's disease. *Neurology* 1997;49:786–794.
- Juarez J, Corsi-Cabrera M. Sex differences in interhemispheric correlation and spectral power of EEG activity. *Brain Res Bull* 1995;38:149–151.
- Jung MW, Wiener SI, McNaughton BL. Comparison of spatial firing characteristics of units in dorsal and ventral hippocampus of the rat. *J Neurosci* 1994;14:7347–7356.
- Jobst KA, Smith AD, Szatmari M. Detection in life of confirmed Alzheimer's disease using a simple measurement of medial temporal lobe atrophy by computed tomography. *Lancet* 1992;340:1179–1183.
- Jobst KA, Smith AD, Szatmari M. Rapidly progressing atrophy of medial temporal lobe in Alzheimer's disease. *Lancet* 1994;343:829–830.
- Kahana MJ, Sekuler R, Caplan JB, et al. Human theta oscillations exhibit task dependence during virtual maze navigation. *Nature* 1999;399:781–784.
- Kirk IJ, Mackay JC. The role of theta-range oscillations in synchronising and integrating activity in distributed mnemonic networks. *Cortex* 2003;39:993–1008.
- Klimesch W, Doppelmayr M, Russegger H, Pachinger T. Theta band power in the human scalp EEG and the encoding of new information. *Neuroreport* 1996;7:1235–1240.
- Lopes da Silva FH. The rhythmic slow activity (theta) of the limbic cortex: an oscillation in search of a function. In: Basar E, Bullock TH, eds. *Induced rhythms in the brain*. Boston, Birkhäuser, 1992:83–102.
- Maguire EA, Frackowiak RS, Frith CD. Learning to find your way: a role for the human hippocampal formation. *Proc R Soc Lond B Biol Sci* 1996;263:1745–1750.
- McKhann G, Drachman D, Folstein M, et al. Clinical diagnosis of Alzheimer's disease: report of the NINCDS-ADRDA Work Group under the auspices of Department of Health and Human Services Task Force on Alzheimer's Disease. *Neurology* 1984;34:939–944.
- Miller R. *Cortico-hippocampal interplay and the representation of contexts in the brain*. Berlin, Heidelberg, New York: Springer, 1991.
- Milner B, Squire LR, Kandel ER. Cognitive neuroscience and the study of memory. *Neuron* 1998;20:445–468.
- Mizuno K, Wakai M, Takeda A, Sobue G. Medial temporal atrophy and memory impairment in early stage of Alzheimer's disease: an MRI volumetric and memory assessment study. *J Neurol Sci* 2000;173:18–24.
- Montgomery SA, Asberg M. A new depression scale designed to be sensitive to change. *Br J Psychiatry* 1979;134:382–389.
- Moser MB, Moser EI. Functional differentiation in the hippocampus. *Hippocampus* 1998;8:608–619.
- Moser MB, Moser EI, Forrest E, et al. Spatial learning with a minislab in the dorsal hippocampus. *Proc Natl Acad Sci U S A* 1995;92:9697–9701.
- Narr KL, Thompson PM, Szeszko P, et al. Regional specificity of hippocampal volume reductions in first-episode schizophrenia. *Neuroimage* 2004;21(4):1563–575.
- Pegues MP, Rogers LJ, Amend D, et al. Anterior hippocampal volume reduction in male patients with schizophrenia. *Schizophr Res* 2003;60:105–115.
- Pihlajamäki M, Tanila H, Kononen M, et al. Visual presentation of novel objects and new spatial arrangements of objects differentially activates the medial temporal lobe subareas in humans. *Eur J Neurosci* 2004;19:1939–1949.
- Pizzagalli DA, Oakes TR, Davidson RJ. Coupling of theta activity and glucose metabolism in the human rostral anterior cingulate cortex: an EEG/PET study of normal and depressed subjects. *Psychophysiology* 2003;40:939–949.
- Price JL, Ko AI, Wade MJ, et al. Neuron number in the entorhinal cortex and CA1 in preclinical Alzheimer disease. *Arch Neurol* 2001;58:1395–1402.
- Pritchard LS, John ER, Ferris SH. Quantitative EEG correlates of cognitive deterioration in the elderly. *Neurobiol Aging* 1994;15:85–90.
- Rae-Grant A, Blume W, Lau C. The electroencephalogram in Alzheimer-type dementia: a sequential study correlating the electroencephalogram with psychometric and quantitative pathologic data. *Arch Neurol* 1987;44:50–54.
- Richmond MA, Yee BK, Pouzet B, et al. Dissociating context and space within the hippocampus: effects of complete, dorsal, and ventral excitotoxic hippocampal lesions on conditioned freezing and spatial learning. *Behav Neurosci* 1999;113:1189–1203.
- Riedel-Heller SG, Busse A, Aurich C, et al. Prevalence of dementia according to DSM-III-R and ICD-10: results of the Leipzig Longitudinal Study of the Aged (LEILA75+) Part 1. *Br J Psychiatry* 2001;179:250–254.
- Sawrie SM, Martin RC, Gilliam FG, et al. Visual confrontation naming and hippocampal function: a neural network study using quantitative (1)H magnetic resonance spectroscopy. *Brain* 2000;123:770–780.
- Stewart M, Fox SE. Do septal neurons pace the hippocampal theta rhythm? *Trends Neurosci* 1990;13:163–168.
- Strange BA, Fletcher PC, Henson RN. Segregating the functions of human hippocampus. *Proc Natl Acad Sci U S A* 1999;96:4034–4039.
- Szeszko PR, Goldberg E, Gunduz-Bruce H, et al. Smaller anterior hippocampal formation volume in antipsychotic-naïve patients with first-episode schizophrenia. *Am J Psychiatry* 2003;160:2190–2197.
- Talairach J. *Co-planar stereotaxic atlas of the human brain*. New York: Thieme, 1988.
- Talk A, Kang E, Gabriel M. Independent generation of theta rhythm in the hippocampus and posterior cingulate cortex. *Brain Res* 2004;1015:15–24.
- Tesche CD, Karhu J. Theta oscillations index human hippocampal activation during a working memory task. *Proc Natl Acad Sci U S A* 2000;97:919–924.
- Trivedi MA, Coover GD. Lesions of the ventral hippocampus, but not the dorsal hippocampus, impair conditioned fear expression and inhibitory avoidance on the elevated T-maze. *Neurobiol Learn Mem* 2004;81:172–184.
- Velez-Pardo C, Arellano JJ, Cardona-Gomez P, et al. CA1 hippocampal neuronal loss in familial Alzheimer's disease: presenilin-1 E280A mutation is related to epilepsy. *Epilepsia* 2004;45:751–756.
- Vinogradova OS. Expression, control, and probable functional significance of the neuronal theta-rhythm. *Prog Neurobiol* 1995;45:523–583.
- Wang L, Swank JS, Glick IE, et al. Changes in hippocampal volume and shape across time distinguish dementia of the Alzheimer type from healthy aging. *Neuroimage* 2003;20:667–682.
- Wolf H, Grunwald M, Kruggel F, et al. Hippocampal volume discriminates between normal cognition; questionable and mild dementia in the elderly. *Neurobiol Aging* 2001;22:177–186.
- Wolf H, Kruggel F, Hensel A, et al. The relationship between head size and intracranial volume in elderly subjects. *Brain Res* 2003;973:74–80.
- Wolf H, Hensel A, Arendt T, et al. Serum lipids and hippocampal volume: the link to Alzheimer's disease? *Ann Neurol* 2004;56:745–748.