

Normality results – 2 cluster

```
> normality_results
$pvt_reaction_time

      Shapiro-wilk normality test

data:  residuals(lm(clean_data[[variable]] ~ as.factor(cluster), data = cle
an_data))
W = 0.82109, p-value = 9.095e-08

$nback_miss_1

      Shapiro-wilk normality test

data:  residuals(lm(clean_data[[variable]] ~ as.factor(cluster), data = cle
an_data))
W = 0.95694, p-value = 0.01702

$nback_miss_2

      Shapiro-wilk normality test

data:  residuals(lm(clean_data[[variable]] ~ as.factor(cluster), data = cle
an_data))
W = 0.97487, p-value = 0.1715

$tmt_a_time

      Shapiro-wilk normality test

data:  residuals(lm(clean_data[[variable]] ~ as.factor(cluster), data = cle
an_data))
W = 0.96891, p-value = 0.07907

$tmt_b_time

      Shapiro-wilk normality test

data:  residuals(lm(clean_data[[variable]] ~ as.factor(cluster), data = cle
an_data))
W = 0.83599, p-value = 2.481e-07

$tmt_diff

      Shapiro-wilk normality test

data:  residuals(lm(clean_data[[variable]] ~ as.factor(cluster), data = cle
an_data))
W = 0.8319, p-value = 1.873e-07
```

```
> homogeneity_results
$pvt_reaction_time
Levene's Test for Homogeneity of Variance (center = median)
      Df F value    Pr(>F)
group 1  7.7683 0.006887 **
      68
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$nback_miss_1
Levene's Test for Homogeneity of Variance (center = median)
      Df F value    Pr(>F)
group 1  3.7301 0.05761 .
      68
```

```

---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$nbck_miss_2
Levene's Test for Homogeneity of Variance (center = median)
  Df F value  Pr(>F)
group 1  3.2772 0.07467 .
  68

---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$tmt_a_time
Levene's Test for Homogeneity of Variance (center = median)
  Df F value  Pr(>F)
group 1 10.176 0.002153 **
  68

---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$tmt_b_time
Levene's Test for Homogeneity of Variance (center = median)
  Df F value  Pr(>F)
group 1  5.3301 0.02401 *
  68

---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

$tmt_diff
Levene's Test for Homogeneity of Variance (center = median)
  Df F value  Pr(>F)
group 1  4.0407 0.04838 *
  68

---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

```

```

shapiro_per_cluster
$pvt_reaction_time
$pvt_reaction_time$cluster_1
$pvt_reaction_time$cluster_1$W
  W
0.8264

$pvt_reaction_time$cluster_1$p_value
[1] 0

$pvt_reaction_time$cluster_2
$pvt_reaction_time$cluster_2$W
  W
0.9222

$pvt_reaction_time$cluster_2$p_value
[1] 0.0347

$nbck_miss_1
$nbck_miss_1$cluster_1
$nbck_miss_1$cluster_1$W
  W
0.9412

$nbck_miss_1$cluster_1$p_value
[1] 0.0345

$nbck_miss_1$cluster_2
$nbck_miss_1$cluster_2$W
  W
0.9063

```

```
$nback_miss_1$cluster_2$p_value  
[1] 0.0139
```

```
$nback_miss_2  
$nback_miss_2$cluster_1  
$nback_miss_2$cluster_1$w  
W  
0.9521
```

```
$nback_miss_2$cluster_1$p_value  
[1] 0.0828
```

```
$nback_miss_2$cluster_2  
$nback_miss_2$cluster_2$w  
W  
0.9626
```

```
$nback_miss_2$cluster_2$p_value  
[1] 0.3806
```

```
$tmt_a_time  
$tmt_a_time$cluster_1  
$tmt_a_time$cluster_1$w  
W  
0.9604
```

```
$tmt_a_time$cluster_1$p_value  
[1] 0.163
```

```
$tmt_a_time$cluster_2  
$tmt_a_time$cluster_2$w  
W  
0.9697
```

```
$tmt_a_time$cluster_2$p_value  
[1] 0.5529
```

```
$tmt_b_time  
$tmt_b_time$cluster_1  
$tmt_b_time$cluster_1$w  
W  
0.8246
```

```
$tmt_b_time$cluster_1$p_value  
[1] 0
```

```
$tmt_b_time$cluster_2  
$tmt_b_time$cluster_2$w  
W  
0.9771
```

```
$tmt_b_time$cluster_2$p_value  
[1] 0.7608
```

```
$tmt_diff  
$tmt_diff$cluster_1  
$tmt_diff$cluster_1$w  
W  
0.8089
```

```
$tmt_diff$cluster_1$p_value
[1] 0

$tmt_diff$cluster_2
$tmt_diff$cluster_2$w
0.9609

$tmt_diff$cluster_2$p_value
[1] 0.3467
```

4-Cluster solution

```
> # Overall Kruskal
> kruskal_results
$pvt_reaction_time

    Kruskal-wallis rank sum test

data:  clean_data[[variable]] by as.factor(cluster)
Kruskal-wallis chi-squared = 47.029, df = 3, p-value = 3.426e-10

$nbback_miss_1

    Kruskal-wallis rank sum test

data:  clean_data[[variable]] by as.factor(cluster)
Kruskal-wallis chi-squared = 22.672, df = 3, p-value = 4.726e-05

$nbback_miss_2

    Kruskal-wallis rank sum test

data:  clean_data[[variable]] by as.factor(cluster)
Kruskal-wallis chi-squared = 27.642, df = 3, p-value = 4.318e-06

$tmt_a_time

    Kruskal-wallis rank sum test

data:  clean_data[[variable]] by as.factor(cluster)
Kruskal-wallis chi-squared = 38.848, df = 3, p-value = 1.869e-08

$tmt_b_time

    Kruskal-wallis rank sum test

data:  clean_data[[variable]] by as.factor(cluster)
Kruskal-wallis chi-squared = 34.861, df = 3, p-value = 1.303e-07

$tmt_diff

    Kruskal-wallis rank sum test

data:  clean_data[[variable]] by as.factor(cluster)
Kruskal-wallis chi-squared = 17.649, df = 3, p-value = 0.0005195
```

```
> effect_sizes
$pvt_reaction_time
[1] 15.6764
```

```
$nback_miss_1  
[1] 7.5575
```

```
$nback_miss_2  
[1] 9.214
```

```
$tmt_a_time  
[1] 12.9494
```

```
$tmt_b_time  
[1] 11.6204
```

```
$tmt_diff  
[1] 5.8831
```

```
> # By group  
> kruskal_results_withPCS  
$pvt_reaction_time
```

```
    Kruskal-wallis rank sum test
```

```
data: withPCS_data[[variable]] by as.factor(withPCS_data$cluster)  
Kruskal-wallis chi-squared = 30.167, df = 3, p-value = 1.273e-06
```

```
$nback_miss_1
```

```
    Kruskal-wallis rank sum test
```

```
data: withPCS_data[[variable]] by as.factor(withPCS_data$cluster)  
Kruskal-wallis chi-squared = 7.044, df = 3, p-value = 0.07051
```

```
$nback_miss_2
```

```
    Kruskal-wallis rank sum test
```

```
data: withPCS_data[[variable]] by as.factor(withPCS_data$cluster)  
Kruskal-wallis chi-squared = 15.919, df = 3, p-value = 0.001178
```

```
$tmt_a_time
```

```
    Kruskal-wallis rank sum test
```

```
data: withPCS_data[[variable]] by as.factor(withPCS_data$cluster)  
Kruskal-wallis chi-squared = 22.547, df = 3, p-value = 5.019e-05
```

```
$tmt_b_time
```

```
    Kruskal-wallis rank sum test
```

```
data: withPCS_data[[variable]] by as.factor(withPCS_data$cluster)  
Kruskal-wallis chi-squared = 24.533, df = 3, p-value = 1.933e-05
```

```
$tmt_diff
```

```
    Kruskal-wallis rank sum test
```

```
data: withPCS_data[[variable]] by as.factor(withPCS_data$cluster)  
Kruskal-wallis chi-squared = 14.291, df = 3, p-value = 0.002534
```

```
> effect_sizes_withPCS  
$pvt_reaction_time  
[1] 10.0558
```

```
$nback_miss_1  
[1] 2.348  
  
$nback_miss_2  
[1] 5.3062  
  
$tmt_a_time  
[1] 7.5156  
  
$tmt_b_time  
[1] 8.1778  
  
$tmt_diff  
[1] 4.7638
```

```
> kruskal_results_withoutPCS  
$pvt_reaction_time  
  
Kruskal-wallis rank sum test  
data: withoutPCS_data[[variable]] by as.factor(withoutPCS_data$cluster)  
Kruskal-wallis chi-squared = 11.296, df = 3, p-value = 0.01023  
  
$nback_miss_1  
  
Kruskal-wallis rank sum test  
data: withoutPCS_data[[variable]] by as.factor(withoutPCS_data$cluster)  
Kruskal-wallis chi-squared = 9.7076, df = 3, p-value = 0.02122  
  
$nback_miss_2  
  
Kruskal-wallis rank sum test  
data: withoutPCS_data[[variable]] by as.factor(withoutPCS_data$cluster)  
Kruskal-wallis chi-squared = 9.9672, df = 3, p-value = 0.01885  
  
$tmt_a_time  
  
Kruskal-wallis rank sum test  
data: withoutPCS_data[[variable]] by as.factor(withoutPCS_data$cluster)  
Kruskal-wallis chi-squared = 7.876, df = 3, p-value = 0.04864  
  
$tmt_b_time  
  
Kruskal-wallis rank sum test  
data: withoutPCS_data[[variable]] by as.factor(withoutPCS_data$cluster)  
Kruskal-wallis chi-squared = 6.3102, df = 3, p-value = 0.09746  
  
$tmt_diff  
  
Kruskal-wallis rank sum test  
data: withoutPCS_data[[variable]] by as.factor(withoutPCS_data$cluster)  
Kruskal-wallis chi-squared = 3.4946, df = 3, p-value = 0.3215
```

```
> effect_sizes_withoutPCS  
$pvt_reaction_time  
[1] 3.7653  
  
$nback_miss_1  
[1] 3.2359
```

```
$nback_miss_2  
[1] 3.3224
```

```
$tmt_a_time  
[1] 2.6253
```

```
$tmt_b_time  
[1] 2.1034
```

```
$tmt_diff  
[1] 1.1649
```

```
# Dunn post-hoc  
> dunn_results  
$pvt_reaction_time  
Dunn (1964) Kruskal-Wallis multiple comparison  
p-values adjusted with the Bonferroni method.
```

	Comparison	Z	P.unadj	P.adj
1	1 - 2	5.502634	3.741581e-08	2.244949e-07
2	1 - 3	3.315185	9.158248e-04	5.494949e-03
3	2 - 3	-1.689386	9.114557e-02	5.468734e-01
4	1 - 4	-1.045791	2.956577e-01	1.000000e+00
5	2 - 4	-5.331670	9.731369e-08	5.838821e-07
6	3 - 4	-3.702221	2.137205e-04	1.282323e-03

```
$nback_miss_1  
Dunn (1964) Kruskal-Wallis multiple comparison  
p-values adjusted with the Bonferroni method.
```

	Comparison	Z	P.unadj	P.adj
1	1 - 2	3.582942	3.397466e-04	0.002038480
2	1 - 3	1.019628	3.079049e-01	1.000000000
3	2 - 3	-2.373948	1.759904e-02	0.105594268
4	1 - 4	-1.147434	2.512025e-01	1.000000000
5	2 - 4	-3.972442	7.113967e-05	0.000426838
6	3 - 4	-1.956317	5.042783e-02	0.302566953

```
$nback_miss_2  
Dunn (1964) Kruskal-Wallis multiple comparison  
p-values adjusted with the Bonferroni method.
```

	Comparison	Z	P.unadj	P.adj
1	1 - 2	2.8226171	4.763342e-03	0.0285800516
2	1 - 3	-1.3210439	1.864867e-01	1.0000000000
3	2 - 3	-4.2461507	2.174744e-05	0.0001304846
4	1 - 4	-1.7702442	7.668647e-02	0.4601188136
5	2 - 4	-4.0595290	4.917181e-05	0.0002950308
6	3 - 4	-0.6901557	4.900963e-01	1.0000000000

```
$tmt_a_time  
Dunn (1964) Kruskal-Wallis multiple comparison  
p-values adjusted with the Bonferroni method.
```

	Comparison	Z	P.unadj	P.adj
1	1 - 2	3.186161	1.441745e-03	8.650473e-03
2	1 - 3	-1.501766	1.331576e-01	7.989458e-01
3	2 - 3	-4.804869	1.548527e-06	9.291160e-06
4	1 - 4	-2.455671	1.406220e-02	8.437319e-02
5	2 - 4	-5.073475	3.906152e-07	2.343691e-06
6	3 - 4	-1.223456	2.211574e-01	1.000000e+00

```
$tmt_b_time  
Dunn (1964) Kruskal-Wallis multiple comparison  
p-values adjusted with the Bonferroni method.
```

	Comparison	Z	P.unadj	P.adj
1	1 - 2	0.7671256	4.430068e-01	1.000000e+00
2	1 - 3	-3.2177077	1.292194e-03	7.753165e-03

```
3      2 - 3 -4.3513620 1.352945e-05 8.117669e-05
4      1 - 4 -3.8317902 1.272142e-04 7.632852e-04
5      2 - 4 -4.7005873 2.594143e-06 1.556486e-05
6      3 - 4 -1.2057146 2.279275e-01 1.000000e+00
```

\$tmt_diff

Dunn (1964) Kruskal-wallis multiple comparison
p-values adjusted with the Bonferroni method.

	Comparison	Z	P.unadj	P.adj
1	1 - 2	-0.9296137	0.3525711459	1.000000000
2	1 - 3	-2.6902166	0.0071405647	0.042843388
3	2 - 3	-2.0971096	0.0359838737	0.215903242
4	1 - 4	-3.6396211	0.0002730395	0.001638237
5	2 - 4	-3.1964563	0.0013912692	0.008347615
6	3 - 4	-1.4397525	0.1499374281	0.899624568

dunn_results_withPCS

\$pvt_reaction_time

Dunn (1964) Kruskal-wallis multiple comparison
p-values adjusted with the Bonferroni method.

	Comparison	Z	P.unadj	P.adj
1	1 - 2	4.073680	4.627602e-05	2.776561e-04
2	1 - 3	3.108111	1.882876e-03	1.129726e-02
3	2 - 3	-1.175352	2.398539e-01	1.000000e+00
4	1 - 4	-1.060334	2.889927e-01	1.000000e+00
5	2 - 4	-4.485691	7.267792e-06	4.360675e-05
6	3 - 4	-3.660766	2.514624e-04	1.508775e-03

\$nback_miss_2

Dunn (1964) Kruskal-wallis multiple comparison
p-values adjusted with the Bonferroni method.

	Comparison	Z	P.unadj	P.adj
1	1 - 2	1.423425	0.1546129166	0.92767750
2	1 - 3	-1.484052	0.1377951157	0.82677069
3	2 - 3	-2.807311	0.0049956999	0.02997420
4	1 - 4	-2.585591	0.0097212071	0.05832724
5	2 - 4	-3.674611	0.0002382116	0.00142927
6	3 - 4	-1.343944	0.1789663454	1.00000000

\$tmt_a_time

Dunn (1964) Kruskal-wallis multiple comparison
p-values adjusted with the Bonferroni method.

	Comparison	Z	P.unadj	P.adj
1	1 - 2	2.301185	0.0213811718	0.1282870308
2	1 - 3	-1.561865	0.1183198883	0.7099193300
3	2 - 3	-3.757631	0.0001715293	0.0010291761
4	1 - 4	-2.449222	0.0143165075	0.0858990448
5	2 - 4	-4.294074	0.0000175424	0.0001052544
6	3 - 4	-1.142473	0.2532576377	1.0000000000

\$tmt_b_time

Dunn (1964) Kruskal-wallis multiple comparison
p-values adjusted with the Bonferroni method.

	Comparison	Z	P.unadj	P.adj
1	1 - 2	0.4405916	6.595087e-01	1.0000000000
2	1 - 3	-2.8817490	3.954746e-03	0.0237284781
3	2 - 3	-3.1278364	1.760982e-03	0.0105658897
4	1 - 4	-3.8335883	1.262875e-04	0.0007577247
5	2 - 4	-4.0229604	5.747116e-05	0.0003448269
6	3 - 4	-1.4225441	1.548684e-01	0.9292101972

\$tmt_diff

Dunn (1964) Kruskal-wallis multiple comparison
p-values adjusted with the Bonferroni method.

	Comparison	Z	P.unadj	P.adj
1	1 - 2	-0.4398468	0.6600480919	1.000000000
2	1 - 3	-2.0460426	0.0407521764	0.244513058
3	2 - 3	-1.4680978	0.1420776526	0.852465916
4	1 - 4	-3.4815545	0.0004985123	0.002991074
5	2 - 4	-2.9365925	0.0033183982	0.019910389
6	3 - 4	-1.7697124	0.0767750588	0.460650353

dunn_results_withoutPCS

\$pvt_reaction_time

Dunn (1964) Kruskal-Wallis multiple comparison
p-values adjusted with the Bonferroni method.

	Comparison	Z	P.unadj	P.adj
1	1 - 2	2.9169862	0.003534315	0.02120589
2	1 - 3	1.7045579	0.088276902	0.52966141
3	2 - 3	-0.4874180	0.625962171	1.000000000
4	1 - 4	-0.3380617	0.735316691	1.000000000
5	2 - 4	-1.9324538	0.053303522	0.31982113
6	3 - 4	-1.4547859	0.145728533	0.87437120

\$nback_miss_1

Dunn (1964) Kruskal-Wallis multiple comparison
p-values adjusted with the Bonferroni method.

	Comparison	Z	P.unadj	P.adj
1	1 - 2	2.77915418	0.005450065	0.03270039
2	1 - 3	1.02229063	0.306643368	1.000000000
3	2 - 3	-1.20413183	0.228538629	1.000000000
4	1 - 4	0.01425219	0.988628780	1.000000000
5	2 - 4	-1.47462445	0.140313538	0.84188123
6	3 - 4	-0.66238206	0.507726395	1.000000000

\$nback_miss_2

Dunn (1964) Kruskal-Wallis multiple comparison
p-values adjusted with the Bonferroni method.

	Comparison	Z	P.unadj	P.adj
1	1 - 2	2.2538313	0.02420678	0.14524071
2	1 - 3	-0.4350091	0.66355583	1.000000000
3	2 - 3	-2.5305324	0.01138896	0.06833374
4	1 - 4	0.8348999	0.40377405	1.000000000
5	2 - 4	-0.2986713	0.76519088	1.000000000
6	3 - 4	1.0961199	0.27302633	1.000000000

\$tmt_a_time

Dunn (1964) Kruskal-Wallis multiple comparison
p-values adjusted with the Bonferroni method.

	Comparison	Z	P.unadj	P.adj
1	1 - 2	1.3865944	0.16556547	0.9933928
2	1 - 3	-0.6050279	0.54516049	1.00000000
3	2 - 3	-1.9716205	0.04865294	0.2919177
4	1 - 4	-1.0708557	0.28423434	1.00000000
5	2 - 4	-1.9104163	0.05607963	0.3364778
6	3 - 4	-0.6366632	0.52434424	1.00000000