TITLE GOES HERE

A. J. Martin

Supervisor: Dr Trent Mattner

October 20??

Thesis submitted for the degree of Honours in ??????



Declaration

| Except where stated this thesis is, to the best of my knowledge, my own work and my supervisor has approved its submission. |
|---|
| |
| Signed by student: |
| Date: |
| Signed by supervisor: |
| Date: |

Acknowledgements

Abstract

| | 1 | 1 | | - 1 |
|---|------------------------|----|---|--------------|
| А | $\mathbf{b}\mathbf{s}$ | τr | a | \mathbf{T} |

Abstract

Contents

| 1 | Introduction 1.1 Theory | 2 2 |
|---------------------------|-----------------------------------|---------------|
| 2 | The Squire-Long model | 3 |
| 3 | | 4 |
| | 3.1 Rotating Flow | 4 |
| | 3.2 Rankine Vortex | 4 |
| | 3.3 Lamb-Oseen Vortex | 4 |
| 4 | Solver | 5 |
| | 4.1 Computational Efficiency | 5 |
| $\mathbf{A}_{\mathtt{l}}$ | ppendices | 6 |
| \mathbf{A} | Derivation of Squire-Long | 7 |
| В | Code | 8 |
| | B.1. Code1 | 8 |

Introduction

1.1 Theory

Definition 1.1.1. Stream Function

Definition 1.1.2. Vorticity

Definition 1.1.3. Circulation

Theorem 1.1.4.

Lemma 1.1.5. 1.1.1 Finite Differences

First order:

Backward difference:

$$f'(x) = \frac{f(x) - f(x - h)}{h} + \mathcal{O}(h)$$

Forward Difference

$$f'(x) = \frac{f(x+h) - f(x)}{h} + \mathcal{O}(h)$$

Central Differences

$$f'(x) = \frac{f(x+h) - f(x-h)}{2h} + \mathcal{O}(h^2)$$

The Squire-Long model

2.1 Vorticity form

Azimuthal vorticity, η .

Flow Examples

- 3.1 Rotating Flow
- 3.2 Rankine Vortex
- 3.3 Lamb-Oseen Vortex (Possibly Bachelor Vortex?)

Solver

4.1 Computational Efficiency?

Appendices

Appendix A Derivation of Squire-Long

Appendix B

Conclusion

Appendix C

Code

C.1 Code1

Bibliography