

# School of Mathematical Sciences

## Assignment Cover Sheet



Student Name	
Student ID	
Assessment Title	<b>Assignment 3</b>
Due Date	Thursday, 12 September, 2019 @ 12:00 noon
Course / Program	APP MTH 3022–Optimal Functions & Nanomechanics
Date Submitted	
<b>OFFICE USE ONLY</b> Date Received	

### KEEP A COPY

It is a good idea to keep a copy of your work for your own records. If you have submitted assessment work electronically make sure you have a backup copy.

### PLAGIARISM AND COLLUSION

**Plagiarism:** using another person's ideas, designs, words or works without appropriate acknowledgement.

**Collusion:** another person assisting in the production of an assessment submission without the express requirement, or consent or knowledge of the assessor.

**In this course you are encouraged to work with other students but the work you submit must be your own. This means you must understand it and be able to explain it if required.**

### CONSEQUENCES OF PLAGIARISM AND COLLUSION

The penalties associated with plagiarism and collusion are designed to impose sanctions on offenders that reflect the seriousness of the University's commitment to academic integrity. Penalties may include: the requirement to revise and resubmit assessment work, receiving a result of zero for the assessment work, failing the course, expulsion and/or receiving a financial penalty.

I declare that all material in this assessment is my own work except where there is clear acknowledgement and reference to the work of others. I have read the University Policy Statement on Plagiarism, Collusion and Related Forms of Cheating (<http://www.adelaide.edu.au/policies/230/>).

I give permission for my assessment work to be reproduced and submitted to other academic staff for the purposes of assessment and to be copied, submitted and retained in a form suitable for electronic checking of plagiarism.

Signed ..... Date .....

# OFN Assignment 3

Andrew Martin

September 11, 2019

1. Find the form of extremals to the following

(a)

$$F\{y(x), z(x)\} = \int_{x_0}^{x_1} (8yz - 5y^2 + y'^2 - 4z'^2) dx$$

Let  $f = 8yz - 5y^2 + y'^2 - 4z'^2$

The Euler-Lagrange equations are

$$\begin{aligned}\frac{d}{dx} \left( \frac{\partial f}{\partial y'} \right) - \frac{\partial f}{\partial y} &= 0 \\ \frac{d}{dx} \left( \frac{\partial f}{\partial z'} \right) - \frac{\partial f}{\partial z} &= 0\end{aligned}$$

Where

$$\begin{aligned}\frac{\partial f}{\partial y'} &= 2y', & \frac{\partial f}{\partial y} &= 8z - 10y \\ \frac{\partial f}{\partial z'} &= -8z', & \frac{\partial f}{\partial z} &= 8y\end{aligned}$$

Plugging into EL

$$\begin{aligned}2y'' - 8z + 10y &= 0 \\ -8z'' - 8y &= 0\end{aligned}$$

We get

$$y = -z''$$

Use this in the first equation to get a fourth order ODE

$$-2z^{(4)} - 8z - 10z'' = 0 \implies z^{(4)} + 4z + 5z'' = 0$$

Using the characteristic equation:

$$\begin{aligned}\lambda^4 + 5\lambda^2 + 4 &= 0 \\ \mu^2 + 5\mu + 4 &= 0 \\ (\mu + 1)(\mu + 4) &= 0 \\ \implies \mu = -1, & \quad \mu = -4 \\ \implies \lambda = \pm i, & \quad \lambda = \pm 2i\end{aligned}$$

Hence the  $z$  solution is

$$z(x) = c_1 \sin(x) + c_2 \cos(x) + c_3 \sin(2x) + c_4 \cos(2x)$$

And hence

$$y(x) = -z'' = c_1 \sin(x) + c_2 \cos(x) + 4c_3 \sin(2x) + 4c_4 \cos(2x)$$

Giving the extremal

$$F = \int_{x_0}^{x_1} (8yz - 5y^2 + y'^2 - 4z'^2) dx$$

(b)

$$F\{\mathbf{q}\} = \int_{t_0}^{t_1} (\dot{q}_1 q_2 + \dot{q}_2 q_3 + q_1 \dot{q}_3 - \dot{q}_1^2) dt$$

Where  $\dot{q}_i := \frac{dq_i}{dt}$

EL is the set of equations

$$\frac{d}{dt} \left( \frac{\partial f}{\partial \dot{q}_i} \right) - \frac{\partial f}{\partial q_i} = 0$$

$$\begin{aligned} \frac{\partial f}{\partial \dot{q}_1} &= q_2 - 2\dot{q}_1, & \frac{\partial f}{\partial q_1} &= \dot{q}_3 \\ \frac{\partial f}{\partial \dot{q}_2} &= q_3, & \frac{\partial f}{\partial q_2} &= \dot{q}_1 \\ \frac{\partial f}{\partial \dot{q}_3} &= q_1, & \frac{\partial f}{\partial q_3} &= \dot{q}_2 \end{aligned}$$

So the EL equations are:

$$\begin{aligned} \dot{q}_2 - 2\ddot{q}_1 - \dot{q}_3 &= 0 \\ \dot{q}_3 - \dot{q}_1 &= 0 \\ \dot{q}_1 - \dot{q}_2 &= 0 \end{aligned}$$

$$\dot{q}_2 - 2\ddot{q}_1 - \dot{q}_3 = 0 \tag{1}$$

$$\dot{q}_1 = \dot{q}_3 \tag{2}$$

$$\dot{q}_1 = \dot{q}_2 \tag{3}$$

Using (2), (3) shows that

$$\dot{q}_1 = \dot{q}_2 = \dot{q}_3$$

And hence (1) gives

$$-2\ddot{q}_1 = 0 \implies q_1 = at + c_1$$

And hence

$$q_1 = at + c_1$$

$$q_2 = at + c_2$$

$$q_3 = at + c_3$$

Hence the extremal has form

$$F = \int_{t_0}^{t_1} (\dot{q}_1 q_2 + \dot{q}_2 q_3 + q_1 \dot{q}_3 - \dot{q}_1^2) dt$$

2. Find the extremal to

$$F\{y\} = \int_0^1 (y''^2 - 360x^2y) dx$$

Subject to  $y(0) = 0, y'(0) = 1, y(1) = 1$ , and  $y'(1) = 5/2$

$$f = y''^2 - 360x^2y$$

$$\frac{\partial f}{\partial y''} = 2y'', \quad \frac{\partial f}{\partial y'} = 0, \quad \frac{\partial f}{\partial y} = -360x^2$$

EL gives

$$\begin{aligned} 0 &= \frac{\partial f}{\partial y} - \frac{d}{dx} \left( \frac{\partial f}{\partial y'} \right) + \frac{d^2}{dx^2} \left( \frac{\partial f}{\partial y''} \right) \\ &= -360x^2 + 2y^{(4)} \\ y^{(4)} &= 180x^2 \\ y &= a + bx + cx^2 + dx^3 + \frac{x^6}{2} \end{aligned}$$

BCs

$$\begin{aligned} y(0) = 0 &\implies a = 0 \\ y'(0) = 1 &\implies b = 1 \\ y(1) = 1 &\implies 1 + c + d + \frac{1}{2} = 1 \\ y'(1) = 5/2 &\implies 1 + 2c + 3d + 5/2 = 5/2 \end{aligned}$$

$$\begin{aligned} c + d &= -\frac{1}{2} \\ c + \frac{3}{2}d &= -\frac{1}{2} \end{aligned}$$

Hence  $d = 0$  and  $c = -\frac{1}{2}$  Hence

$$\boxed{y(x) = x - \frac{1}{2}x^2 + \frac{x^6}{2}}$$

3. (a) Consider the integral definition of the beta function

$$B(x, y) = \int_0^1 t^{x-1} (1-t)^{y-1} dt \quad (4)$$

i. Write an integral expression for  $B(z - \frac{1}{2}, z + \frac{1}{2})$

$$\begin{aligned} B(z - \frac{1}{2}, z + \frac{1}{2}) &= \int_0^1 t^{z-\frac{1}{2}-1} (1-t)^{z+\frac{1}{2}-1} dt \\ &= \int_0^1 t^{z-\frac{3}{2}} (1-t)^{z-\frac{1}{2}} dt \end{aligned}$$

- ii. Substitute  $2t = 1 + s$  into it (take care on the limits)  
 $t = 0$  gives  $s = -1$  and  $t = 1$  gives  $s = 1$ , and  $dt = 1/2ds$

$$\begin{aligned}
 B\left(z - \frac{1}{2}, z + \frac{1}{2}\right) &= \int_0^1 t^{z-\frac{3}{2}}(1-t)^{z-\frac{1}{2}} dt \\
 &= \frac{1}{2} \int_{-1}^1 \left(\frac{1+s}{2}\right)^{z-\frac{3}{2}} \left(1 - \left(\frac{1+s}{2}\right)\right)^{z-\frac{1}{2}} ds \\
 &= \frac{1}{2} \int_{-1}^1 \left(\frac{1+s}{2}\right)^{z-\frac{3}{2}} \left(\frac{1-s}{2}\right)^{z-\frac{1}{2}} ds \\
 &= 2^{-1} \int_{-1}^1 (1+s)^{z-\frac{3}{2}} 2^{\frac{3}{2}-z} (1-s)^{z-\frac{1}{2}} 2^{\frac{1}{2}-z} ds \\
 &= 2^{1-2z} \int_{-1}^1 (1+s)^{z-\frac{3}{2}} (1-s)^{z-\frac{1}{2}} ds \\
 &= 2^{1-2z} \int_{-1}^1 (1-s^2)^z (1+s)^{-\frac{3}{2}} (1-s)^{-\frac{1}{2}} ds
 \end{aligned}$$

- iii. Decompose into even and odd parts in  $s$

To decompose a function  $f(x)$  into even and odd parts:

$$f(x) = f_e(x) + f_o(x) = \frac{f(x) + f(-x)}{2} + \frac{f(x) - f(-x)}{2}$$

So

$$f(s) = (1-s^2)^z (1+s)^{-3/2} (1-s)^{-1/2}$$

$$\begin{aligned}
 f_e(s) &= \frac{(1-s^2)^z (1+s)^{-3/2} (1-s)^{-1/2}}{2} + \frac{(1-s^2)^z (1-s)^{-3/2} (1+s)^{-1/2}}{2} \\
 f_o(s) &= \frac{(1-s^2)^z (1+s)^{-3/2} (1-s)^{-1/2}}{2} - \frac{(1-s^2)^z (1-s)^{-3/2} (1+s)^{-1/2}}{2}
 \end{aligned}$$

So we have

$$\begin{aligned}
 &= 2^{1-2z} \int_{-1}^1 (1-s^2)^z (1+s)^{-\frac{3}{2}} (1-s)^{-\frac{1}{2}} ds \\
 &= 2^{1-2z} \left( \int_{-1}^1 f_e(s) ds + \int_{-1}^1 f_o(s) ds \right) \\
 &= 2^{1-2z} \left( \int_{-1}^1 \frac{(1-s^2)^z (1+s)^{-3/2} (1-s)^{-1/2}}{2} + \frac{(1-s^2)^z (1-s)^{-3/2} (1+s)^{-1/2}}{2} ds \right. \\
 &\quad \left. + \int_{-1}^1 \frac{(1-s^2)^z (1+s)^{-3/2} (1-s)^{-1/2}}{2} - \frac{(1-s^2)^z (1-s)^{-3/2} (1+s)^{-1/2}}{2} ds \right)
 \end{aligned}$$

- iv. Use parity to half the integration ranges in each integral. *Hint* bisect the interval and sub  $\sigma = -s$  into the negative interval

Is this just to show that integrals over symmetric endpoints are

- Double one side for even;
- equal to zero for odd?

Even interval,  $F_e$

$$\begin{aligned}
 F_e(z) &= \int_{-1}^1 \frac{(1-s^2)^z(1+s)^{-3/2}(1-s)^{-1/2}}{2} + \frac{(1-s^2)^z(1-s)^{-3/2}(1+s)^{-1/2}}{2} ds \\
 &= \int_0^1 \frac{(1-s^2)^z(1+s)^{-3/2}(1-s)^{-1/2}}{2} + \frac{(1-s^2)^z(1-s)^{-3/2}(1+s)^{-1/2}}{2} ds \\
 &\quad - \int_1^0 \frac{(1-\sigma^2)^z(1-\sigma)^{-3/2}(1+\sigma)^{-1/2}}{2} + \frac{(1-\sigma^2)^z(1+\sigma)^{-3/2}(1-\sigma)^{-1/2}}{2} d\sigma \\
 &= \int_0^1 \frac{(1-s^2)^z(1+s)^{-3/2}(1-s)^{-1/2}}{2} + \frac{(1-s^2)^z(1-s)^{-3/2}(1+s)^{-1/2}}{2} ds \\
 &\quad + \int_0^1 \frac{(1-\sigma^2)^z(1-\sigma)^{-3/2}(1+\sigma)^{-1/2}}{2} + \frac{(1-\sigma^2)^z(1+\sigma)^{-3/2}(1-\sigma)^{-1/2}}{2} d\sigma \\
 &= \int_0^1 (1-s^2)^z(1+s)^{-3/2}(1-s)^{-1/2} + (1-s^2)^z(1-s)^{-3/2}(1+s)^{-1/2} ds \\
 \\
 F_o(z) &= \int_{-1}^1 \frac{(1-s^2)^z(1+s)^{-3/2}(1-s)^{-1/2}}{2} - \frac{(1-s^2)^z(1-s)^{-3/2}(1+s)^{-1/2}}{2} ds \\
 &= \int_0^1 \frac{(1-s^2)^z(1+s)^{-3/2}(1-s)^{-1/2}}{2} + \frac{(1-s^2)^z(1-s)^{-3/2}(1+s)^{-1/2}}{2} ds \\
 &\quad + \int_1^0 \frac{(1-\sigma^2)^z(1-\sigma)^{-3/2}(1+\sigma)^{-1/2}}{2} + \frac{(1-\sigma^2)^z(1+\sigma)^{-3/2}(1-\sigma)^{-1/2}}{2} d\sigma \\
 &= \int_0^1 \frac{(1-s^2)^z(1+s)^{-3/2}(1-s)^{-1/2}}{2} + \frac{(1-s^2)^z(1-s)^{-3/2}(1+s)^{-1/2}}{2} ds \\
 &\quad - \int_0^1 \frac{(1-\sigma^2)^z(1-\sigma)^{-3/2}(1+\sigma)^{-1/2}}{2} + \frac{(1-\sigma^2)^z(1+\sigma)^{-3/2}(1-\sigma)^{-1/2}}{2} d\sigma \\
 &= 0
 \end{aligned}$$

Hence the full integral is

$$2^{1-2z}(F_e+F_o) = 2^{1-2z} \int_0^1 (1-s^2)^z(1+s)^{-3/2}(1-s)^{-1/2} + (1-s^2)^z(1-s)^{-3/2}(1+s)^{-1/2} ds$$

v. Use  $\tau = s^2$  to get the form from eqn 4

$\tau = s^2$  gives  $ds = \frac{1}{2\sqrt{\tau}}d\tau = 2^{-1}\tau^{-1/2}d\tau$  and bounds stay the same

$$\begin{aligned}
 &= 2^{1-2z} \int_0^1 (1-s^2)^z (1+s)^{-3/2} (1-s)^{-1/2} + (1-s^2)^z (1-s)^{-3/2} (1+s)^{-1/2} ds \\
 &= 2^{1-2z} \int_0^1 ((1-s) + (1+s)) (1-s^2)^z (1-s)^{-3/2} (1+s)^{-3/2} ds \quad *** \\
 &= 2^{1-2z} \int_0^1 2(1-s^2)^z (1-s)^{-3/2} (1+s)^{-3/2} ds \\
 &= 2^{1-2z} \int_0^1 2(1-s^2)^z ((1-s)(1+s))^{-3/2} ds \\
 &= 2^{1-2z} \int_0^1 2(1-s^2)^z (1-s^2)^{-3/2} ds \\
 &= 2^{1-2z} \int_0^1 (1-\tau)^{z-3/2} \tau^{-1/2} d\tau \\
 &= 2^{1-2z} \int_0^1 \tau^{-1/2} (1-\tau)^{z-3/2} d\tau \\
 &= 2^{1-2z} B(1/2, z-1/2)
 \end{aligned}$$

\*\*\* putting the two parts even, I have multiplied the left part by  $(1-s)^{2/2}(1-s)^{-2/2}$  and the right part by  $(1+s)^{2/2}(1+s)^{-2/2}$ , i.e. multiplying by 1.

- vi. Using all the previous parts, and the relationship between the beta and gamma functions, derive

$$\Gamma(2z) = 2^{2z-1} \pi^{-1/2} \Gamma(z) \Gamma(z+1/2)$$

We have the beta/gamma relationship:

$$B(x, y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)}$$

$$\begin{aligned}
 B(z - \frac{1}{2}, z + \frac{1}{2}) &= 2^{1-2z} B(\frac{1}{2}, z - \frac{1}{2}) \\
 \frac{\Gamma(z - \frac{1}{2})\Gamma(z + \frac{1}{2})}{\Gamma(z - \frac{1}{2} + z + \frac{1}{2})} &= 2^{1-2z} \frac{\Gamma(\frac{1}{2})\Gamma(z - \frac{1}{2})}{\Gamma(z)} \\
 \frac{\Gamma(z + \frac{1}{2})}{\Gamma(2z)} &= 2^{1-2z} \frac{\pi^{-1/2}}{\Gamma(z)} \\
 \implies \Gamma(2z) &= 2^{2z-1} \pi^{-1/2} \Gamma(z) \Gamma(z+1/2)
 \end{aligned}$$

- (b) From the integral definitions given in class, show

$$K(k) = \frac{\pi}{2} F(1/2, 1/2; 1; k^2)$$

Note that, using Euler's integral formula

$$\begin{aligned}
 F(a, b; c; z) &= \frac{\Gamma(c)}{\Gamma(b)\Gamma(c-b)} \int_0^1 t^{b-1} (1-t)^{c-b-1} (1-tz)^{-a} dt \\
 F(1/2, 1/2; 1; k^2) &= \frac{\Gamma(1)}{\Gamma(1/2)\Gamma(1/2)} \int_0^1 t^{-1/2} (1-t)^{-1/2} (1-tk^2)^{-1/2} dt \\
 &= \frac{1}{\pi} \int_0^1 t^{-1/2} (1-t)^{-1/2} (1-tk^2)^{-1/2} dt
 \end{aligned}$$

Sub  $t = \sin^2 \vartheta$ ,  $dt = 2 \sin \vartheta \cos \vartheta d\vartheta$ , and  $t = 1$  gives  $\vartheta = \pi/2$

$$\begin{aligned} F(1/2, 1/2; 1; k^2) &= \frac{1}{\pi} \int_0^1 t^{-1/2} (1-t)^{-1/2} (1-tk^2)^{-1/2} dt \\ &= \frac{1}{\pi} \int_0^{\pi/2} \frac{1}{\sin \vartheta} \frac{1}{\sqrt{1-\sin^2 \vartheta}} \frac{1}{\sqrt{1-k^2 \sin^2 \vartheta}} 2 \sin \vartheta \cos \vartheta d\vartheta \\ &= \frac{2}{\pi} \int_0^{\pi/2} \frac{1}{\cos \vartheta} \frac{1}{\sqrt{1-k^2 \sin^2 \vartheta}} \cos \vartheta d\vartheta \\ &= \frac{2}{\pi} \int_0^{\pi/2} \frac{1}{\sqrt{1-k^2 \sin^2 \vartheta}} d\vartheta \end{aligned}$$

Where  $F(\varphi, k)$  denotes the elliptic integral of the first kind, we get:

$$\begin{aligned} K(k) &= F(\pi/2, k) \\ &= \int_0^{\pi/2} \frac{d\vartheta}{\sqrt{1-k^2 \sin^2 \vartheta}} \\ &= \frac{\pi}{2} \frac{2}{\pi} \int_0^{\pi/2} \frac{1}{\sqrt{1-k^2 \sin^2 \vartheta}} d\vartheta \\ &= \frac{\pi}{2} F(1/2, 1/2; 1; k^2) \end{aligned}$$

As required