

# Topic C Assignment 5

Andrew Martin

June 4, 2019

# Practical Asymptotics (APP MTH 4051/7087)

## Assignment 5 (5%)

Due 14 June 2019

1. Consider the integral

$$I(x) = \int_{-\infty}^{\infty} \frac{e^{x2it(1-t^2/6+3it/4)}}{t^2 + 9} dt$$

- (a) Identify any saddle points, then find the paths of steepest descent and ascent through each of these saddle(s).
- (b) Sketch (or plot) the saddles and paths of steepest descent/ascent.
- (c) Sketch a deformed contour that passes through a saddle point in a direction that will permit  $I(x)$  to be evaluated by the method of steepest descent.
- (d) Use the deformed contour from part (c) to approximate  $I(x)$  to leading-order as  $x \rightarrow \infty$ .

2. Use the method of steepest descents to show that

$$I(x) = \frac{1}{2} \int_{-1}^1 e^{-4xt^2+5ixt-ixt^3} dt \sim \frac{1}{2} e^{-2x} \sqrt{\pi/x}, \quad \text{as } x \rightarrow \infty.$$

A complete solution should go through similar steps to Question 1, but will require a few extra details. [Hints:

- The deformed steepest descent path should have the same endpoints as the original contour (this might look a bit weird).
- The contributions from the end points are negligible compared to that from a saddle point (you need to show this).]

**The following is an extension question which you may do as an alternative the short project.**

3. Continue the analysis of the Airy function  $\text{Ai}(x)$ . Recall that the integral representation was

$$\text{Ai}(x) = \frac{1}{2\pi} \int_{C'} e^{i(t^3/3+xt)} dt,$$

where the contour  $C'$  starts at infinity with  $2\pi/3 < \arg(t) < \pi$  and ends at infinity with  $0 < \arg(t) < \pi/3$ .

- (a) Use the method of steepest descents to find  $\text{Ai}(x)$  to leading-order as  $x \rightarrow \infty$ .
- (b) Now investigate **Stokes phenomenon**, the idea that the behaviour of these integrals depends on the direction  $x$  approaches infinity. We have already seen this is the case along the real axis, and will now extend this to the complex plane. Consider the integral:

$$\text{Ai}(z) = \frac{1}{2\pi} \int_{C'} e^{i(t^3/3+zt)} dt, \quad \text{as } |z| \rightarrow \infty.$$

where  $C'$  is as above and  $z = e^{i\theta}x$ , that is  $\arg(z) = \theta$ .

- i. Determine the location of the two saddle points, which will now vary with  $\theta$ .
  - ii. Find expressions for the steepest descent paths through each saddle (these will now also depend on  $\theta$ ).
  - iii. Write a MATLAB code to plot the saddles and steepest descent paths for any value of  $\theta$ .
  - iv. With reference to the original contour (thinking about how it can be deformed) and the above analysis, discuss why and for what value of  $\theta$  there is a qualitative change in leading-order behaviour.
- (c) The following paper (available on MyUni) extends the above analysis:  
 Berry, M.V., **Asymptotics, superasymptotics, hyperasymptotics**, in *Asymptotics Beyond All Orders*, 1991.
- Briefly summarise the contents of this paper, and discuss how it relates to part (b). You may be particularly interested in Figure 6, which will (hopefully) look familiar.