



COMP30026

Models of Computation

Cezary Kaliszyk and William Umboh

Lecture 3

Consequence and Satisfaction



Recap: Models

\models is short for “is a model of” or “satisfies”.

$v \models F$ iff F is true under v .



Recap: Models

\models is short for “is a model of” or “satisfies”.

$v \models F$ iff F is true under v .

Examples:

$$\{P \mapsto 1, Q \mapsto 0\} \models P \vee Q$$

$$\{P \mapsto 0\} \models \neg P$$



Recap: Models

\models is short for “is a model of” or “satisfies”.

$v \models F$ iff F is true under v .

Examples:

$$\{P \mapsto 1, Q \mapsto 0\} \models P \vee Q$$

$$\{P \mapsto 0\} \models \neg P$$

Non-examples:

$$\{P \mapsto 1, Q \mapsto 0\} \not\models P \rightarrow Q$$

$$\{P \mapsto 1\} \not\models \neg P$$



Recap: Equivalence

“ \equiv ” is short for “is logically equivalent to”.

$A \equiv B$ iff A and B always have equal truth values.



Recap: Equivalence

“ \equiv ” is short for “is logically equivalent to”.

$A \equiv B$ iff A and B always have equal truth values.

Examples:

$$\begin{aligned}P \rightarrow Q &\equiv \neg P \vee Q \\ \neg(P \wedge Q) &\equiv \neg P \vee \neg Q\end{aligned}$$



Recap: Equivalence

“ \equiv ” is short for “is logically equivalent to”.

$A \equiv B$ iff A and B always have equal truth values.

Examples:

$$\begin{aligned}P \rightarrow Q &\equiv \neg P \vee Q \\ \neg(P \wedge Q) &\equiv \neg P \vee \neg Q\end{aligned}$$

Non-examples:

$$\begin{aligned}P \rightarrow Q &\not\equiv R \rightarrow S \\ P \wedge Q &\not\equiv P \vee Q\end{aligned}$$

Semantic Consequence

Definition

G is a *semantic consequence* of F **if and only if** every model of F is a model of G .

For short, we write “ $F \models G$ ”.

“ \models ” is pronounced “(semantically) entails”.



Semantic Consequence

Definition

G is a *semantic consequence* of F **if and only if** every model of F is a model of G .

For short, we write “ $F \models G$ ”.

“ \models ” is pronounced “(semantically) entails”.

Note: $F \equiv G$ iff $F \models G$ and $G \models F$.



Consequence and Implication

models used before defined later?

Let F and G be formulas.

Theorem

$F \models G$ if and only if $\models F \rightarrow G$.

Consequence and Implication

models used before defined later?

Let F and G be formulas.

Theorem

$F \models G$ if and only if $\models F \rightarrow G$.

As an immediate corollary:

Corollary

$F \equiv G$ if and only if $\models F \leftrightarrow G$.



Poll

Of the following formulas, which allow us to conclude $P \rightarrow Q$?

1. P
2. $\neg P$
3. Q
4. $P \rightarrow (Q \wedge R)$
5. $(P \vee R) \rightarrow Q$
6. $\neg P \vee Q$
7. $\neg Q \rightarrow \neg P$
8. $P \rightarrow (Q \vee R)$
9. $(P \rightarrow Q) \vee R$



Poll

Of the following formulas, which allow us to conclude $P \rightarrow Q$?

- | | |
|---------------------------------|-------|
| 1. P | ■ No |
| 2. $\neg P$ | ■ Yes |
| 3. Q | ■ Yes |
| 4. $P \rightarrow (Q \wedge R)$ | ■ Yes |
| 5. $(P \vee R) \rightarrow Q$ | ■ Yes |
| 6. $\neg P \vee Q$ | ■ Yes |
| 7. $\neg Q \rightarrow \neg P$ | ■ Yes |
| 8. $P \rightarrow (Q \vee R)$ | ■ No |
| 9. $(P \rightarrow Q) \vee R$ | ■ No |



Poll

Of the following formulas, which allow us to conclude $P \rightarrow Q$?

1. P
2. $\neg P$
3. Q
4. $P \rightarrow (Q \wedge R)$
5. $(P \vee R) \rightarrow Q$
6. $\neg P \vee Q$
7. $\neg Q \rightarrow \neg P$
8. $P \rightarrow (Q \vee R)$
9. $(P \rightarrow Q) \vee R$

Which of the formulas are consequences of $P \rightarrow Q$?



Tautology

Tautology: a logical formula which is always true.

Tautology

Tautology: a logical formula which is always true.

$(\neg P \wedge Q) \rightarrow (P \rightarrow R)$ is a tautology:

P	Q	R	$(\neg P \wedge Q) \rightarrow (P \rightarrow R)$						
1	1	1	0	1	0	1	1	1	1
1	1	0	0	1	0	1	1	0	0
1	0	1	0	1	0	0	1	1	1
1	0	0	0	1	0	0	1	0	0
0	1	1	1	0	1	1	1	0	1
0	1	0	1	0	1	1	1	0	0
0	0	1	1	0	0	0	1	0	1
0	0	0	1	0	0	0	1	0	0



Contradiction

Contradiction: A logical formula which is always false.

$Q \wedge \neg Q$ is a contradiction.



Contradiction

Contradiction: A logical formula which is always false.

$Q \wedge \neg Q$ is a contradiction.

$P \wedge Q \wedge (\neg Q \leftrightarrow (\neg P \vee Q))$ is a contradiction.

No need for whole truth table.



Contradiction

Contradiction: A logical formula which is always false.

$Q \wedge \neg Q$ is a contradiction.

$P \wedge Q \wedge (\neg Q \leftrightarrow (\neg P \vee Q))$ is a contradiction.

No need for whole truth table.

If either P or Q are false, \wedge makes whole formula false.



Contradiction

Contradiction: A logical formula which is always false.

$Q \wedge \neg Q$ is a contradiction.

$P \wedge Q \wedge (\neg Q \leftrightarrow (\neg P \vee Q))$ is a contradiction.

No need for whole truth table.

If either P or Q are false, \wedge makes whole formula false.

When P and Q are both true:

P	Q	$(P \wedge Q) \wedge (\neg Q \leftrightarrow (\neg P \vee Q))$									
1	1	1	1	1	0	0	1	0	0	1	1



Contradiction

Contradiction: A logical formula which is always false.

$Q \wedge \neg Q$ is a contradiction.

$P \wedge Q \wedge (\neg Q \leftrightarrow (\neg P \vee Q))$ is a contradiction.

No need for whole truth table.

If either P or Q are false, \wedge makes whole formula false.

When P and Q are both true:

P	Q	$(P \wedge Q) \wedge (\neg Q \leftrightarrow (\neg P \vee Q))$									
1	1	1	1	1	0	0	1	0	0	1	1

Negating a contradiction gives a tautology and vice versa.



Tautologies Are Valid

Consider: “If the program works, then the program works.”

It is **true** *regardless* of what “program” or “works” mean.



Tautologies Are Valid

Consider: “If the program works, then the program works.”

It is **true** *regardless* of what “program” or “works” mean.

Valid: Always true.

Non-valid: Sometimes false.

$\models F$ is short for “ F is valid”.



Contradictions Are Unsatisfiable

Consider:

“the application is good and the application is not good.”

It is **false** *regardless* of what “the application” or “good” mean.



Contradictions Are Unsatisfiable

Consider:

“the application is good and the application is not good.”

It is **false** *regardless* of what “the application” or “good” mean.

Unsatisfiable: Never true.

Satisfiable: Sometimes true.



Most Statements Are Contingent

Consider: “It is currently raining.”

It is true **if and only if** it is currently raining.



Most Statements Are Contingent

Consider: “It is currently raining.”

It is true **if and only if** it is currently raining.

Contingent: Sometimes true, sometimes false.



Poll

Classify the following formulas as valid, contingent, or unsatisfiable:

1. P
2. $P \leftrightarrow \neg P$
3. $P \rightarrow (\neg Q \vee P)$
4. $\neg Q \vee \neg(P \wedge \neg Q)$

Classify the following formulas as valid, contingent, or unsatisfiable:

- | | |
|--|-----------------|
| 1. P | ■ contingent |
| 2. $P \leftrightarrow \neg P$ | ■ unsatisfiable |
| 3. $P \rightarrow (\neg Q \vee P)$ | ■ valid |
| 4. $\neg Q \vee \neg(P \wedge \neg Q)$ | ■ valid |



Substitution

Replacing **all** uses of a propositional variable with a given formula.



Substitution

Replacing **all** uses of a propositional variable with a given formula.

Example

Consider " $P \rightarrow P$ ".



Substitution

Replacing **all** uses of a propositional variable with a given formula.

Example

Consider " $P \rightarrow P$ ".

Substitute P with " $(Q \wedge R)$ ".

Substitution

Replacing **all** uses of a propositional variable with a given formula.

Example

Consider " $P \rightarrow P$ ".

Substitute P with " $(Q \wedge R)$ ".

Result: " $(Q \wedge R) \rightarrow (Q \wedge R)$ ".



Substitution

Replacing **all** uses of a propositional variable with a given formula.

Example

Consider " $P \rightarrow P$ ".

Substitute P with " $(Q \wedge R)$ ".

Result: " $(Q \wedge R) \rightarrow (Q \wedge R)$ ".

Substitution preserves validity! Students do not know what preserves MEANS?



Poll

Does substitution preserve unsatisfiability?



Poll

Does substitution preserve unsatisfiability?

Yes!



Poll

Does substitution preserve unsatisfiability?

Yes!

Negation of contradiction is a tautology.



Poll

Does substitution preserve satisfiability?



Poll

Does substitution preserve satisfiability?

No — a counterexample is easy:



Poll

Does substitution preserve satisfiability?

No — a counterexample is easy:

Take P (which is clearly satisfiable).

Then substitute P by $Q \wedge \neg Q$.



Substitution Preserves Logical Equivalence

Denote by $F[A := B]$ the result of substituting A with B in F .

Example: $(P \rightarrow P)[P := Q]$ is $(Q \rightarrow Q)$.

Substitution Preserves Logical Equivalence

Denote by $F[A := B]$ the result of substituting A with B in F .

Example: $(P \rightarrow P)[P := Q]$ is $(Q \rightarrow Q)$.

Theorem

Let F, G, H be formulas and P be any propositional variable.

If $F \equiv G$, then $F[P := H] \equiv G[P := H]$.



Interchange of Equivalents

If $F \equiv G$, then F can be freely replaced with G .

Interchange of Equivalents

If $F \equiv G$, then F can be freely replaced with G .

Theorem

Let H' be the result of replacing an instance of F with G in H .

If $F \equiv G$, then $H \equiv H'$.

Interchange of Equivalents

If $F \equiv G$, then F can be freely replaced with G .

Theorem

Let H' be the result of replacing an instance of F with G in H .

If $F \equiv G$, then $H \equiv H'$.

Result is equivalent: all semantic properties preserved.

Interchange of Equivalents

If $F \equiv G$, then F can be freely replaced with G .

Theorem

Let H' be the result of replacing an instance of F with G in H .

If $F \equiv G$, then $H \equiv H'$.

Result is equivalent: all semantic properties preserved.

Rewrite formulas algebraically!

Some Equivalences

Absorption: $P \wedge P \equiv P$
 $P \vee P \equiv P$

Commutativity: $P \wedge Q \equiv Q \wedge P$
 $P \vee Q \equiv Q \vee P$

Associativity: $P \wedge (Q \wedge R) \equiv (P \wedge Q) \wedge R$
 $P \vee (Q \vee R) \equiv (P \vee Q) \vee R$

Distributivity: $P \wedge (Q \vee R) \equiv (P \wedge Q) \vee (P \wedge R)$
 $P \vee (Q \wedge R) \equiv (P \vee Q) \wedge (P \vee R)$



More Equivalences

Double negation: $P \equiv \neg\neg P$

De Morgan: $\neg(P \wedge Q) \equiv \neg P \vee \neg Q$
 $\neg(P \vee Q) \equiv \neg P \wedge \neg Q$

Implication: $P \rightarrow Q \equiv \neg P \vee Q$

Contraposition: $\neg P \rightarrow \neg Q \equiv Q \rightarrow P$
 $P \rightarrow \neg Q \equiv Q \rightarrow \neg P$
 $\neg P \rightarrow Q \equiv \neg Q \rightarrow P$

Biimplication: $P \leftrightarrow Q \equiv (P \wedge Q) \vee (\neg P \wedge \neg Q)$



Last Equivalences

\perp is equivalent to any unsatisfiable formula

\top is equivalent to any valid formula

Duality: $\neg \top \equiv \perp$
 $\neg \perp \equiv \top$

Negation from absurdity: $P \rightarrow \perp \equiv \neg P$

Identity: $P \vee \perp \equiv P$
 $P \wedge \top \equiv P$

Dominance: $P \wedge \perp \equiv \perp$
 $P \vee \top \equiv \top$

Contradiction: $P \wedge \neg P \equiv \perp$

Excluded middle: $P \vee \neg P \equiv \top$



Poll

Which of these claims hold?

1. $P \rightarrow Q \equiv (Q \leftrightarrow (P \vee Q))$
2. $(P \rightarrow Q) \wedge (P \rightarrow R) \equiv P \rightarrow (Q \wedge R)$
3. $(P \rightarrow R) \wedge (Q \rightarrow R) \models (P \wedge Q) \rightarrow R$