## GRAVITATIONAL POTENTIAL OF A SPHERE



Fig. E.1 shows a point mass m with Cartesian coordinates (x, y, z) as well a system of N point masses  $m_1$ ,  $m_2, m_3, \dots, m_N$ . The ith one of these particles has mass  $m_i$  and coordinates  $(x_i, y_i, z_i)$ . The total mass of the N particles is M,

$$M = \sum_{i=1}^{N} m_i \tag{E.1}$$

The position vector drawn from  $m_i$  to m is  $\mathbf{r}_i$  and the unit vector in the direction of  $\mathbf{r}_i$  is

$$\hat{\mathbf{u}}_i = \frac{\mathbf{r}_i}{r_i}$$




FIG. E.1

A system of point masses and a neighboring test mass m.

The gravitational force exerted on m by  $m_i$  is opposite in direction to  $\mathbf{r}_i$ , and is given by

$$\mathbf{F}_i = -\frac{Gmm_i}{r_i^2}\hat{\mathbf{u}}_i = -\frac{Gmm_i}{r_i^3}\mathbf{r}_i$$

The potential energy of this force is

$$V_i = -G\frac{mm_i}{r_i} \tag{E.2}$$

The total gravitational potential energy of the system due to the gravitational attraction of all the Nparticles is

$$V = \sum_{i=1}^{N} V_i \tag{E.3}$$

Therefore, the total force of gravity  $\mathbf{F}$  on the mass m is

$$\mathbf{F} = -\nabla V = -\left(\frac{\partial V}{\partial x}\hat{\mathbf{i}} + \frac{\partial V}{\partial y}\hat{\mathbf{j}} + \frac{\partial V}{\partial z}\hat{\mathbf{k}}\right)$$
(E.4)

where  $\nabla$  is the gradient operator.

Consider the solid sphere of mass M and radius  $R_0$  illustrated in Fig. E.2. Instead of a discrete system as above, we have a continuum with mass density  $\rho$ . Each "particle" is a differential element  $dM = \rho dv$ of the total mass M. Eq. (E.1) becomes

$$M = \iiint_{v} \rho \, dv \tag{E.5}$$




FIG. E.2

Sphere with a spherically symmetric mass distribution.

where dv is the volume element, and v is the total volume of the sphere. In this case, Eq. (E.2) becomes

$$dV = -G\frac{m \ dM}{r} = -Gm\frac{\rho \ dv}{r}$$

where r is the distance from the differential mass dM to the finite point mass m. Eq. (E.3) is replaced by

$$V = -Gm \iiint \frac{\rho \, dv}{r} \tag{E.6}$$

Let the mass of the sphere have a spherically symmetric distribution, which means that the mass density  $\rho$  depends only on r', the distance from the center C of the sphere. An element of mass dM has spherical coordinates  $(r', \theta, \phi)$ , where the angle  $\theta$  is measured in the xy plane of a Cartesian coordinate system with origin at C, as shown in Fig. E.2. In spherical coordinates the volume element is

$$dv = r'^2 \sin \phi \ d\phi \ dr' \ d\theta \tag{E.7}$$

Therefore Eq. (E.5) becomes

$$M = \int_{\theta=0}^{2\pi} \int_{r'=0}^{R_0} \int_{\phi=0}^{\pi} \rho r'^2 \sin\phi \, d\phi \, dr' \, d\theta = \left(\int_{0}^{2\pi} d\theta\right) \left(\int_{0}^{\pi} \sin\phi \, d\phi\right) \left(\int_{0}^{R_0} \rho r'^2 \, dr'\right) = (2\pi)(2) \left(\int_{0}^{R_0} \rho r'^2 \, dr'\right)$$

so that the mass of the sphere is given by

$$M = 4\pi \int_{r'=0}^{R_0} \rho r'^2 dr'$$
 (E.8)

Substituting Eq. (E.7) into Eq. (E.6) yields

$$V = -Gm \int_{\theta=0}^{2\pi} \int_{r'=0}^{R_0} \int_{\phi=0}^{\pi} \frac{\rho r'^2 \sin \phi \, d\phi \, dr' \, d\theta}{r} = -2\pi Gm \left[ \int_{0}^{R_0} \left( \int_{0}^{\pi} \frac{\sin \phi \, d\phi}{r} \right) \rho r'^2 \, dr' \right]$$
 (E.9)

The distance r is found by using the law of cosines,

$$r = \left(R^2 + {r'}^2 - 2r'R\cos\phi\right)^{1/2}$$

where R is the distance from the center of the sphere to the mass m. Differentiating this equation with respect to  $\phi$ , holding r' constant, yields

$$\frac{\mathrm{d}r}{\mathrm{d}\phi} = \frac{1}{2} \left( R^2 + r'^2 - 2r'R\cos\phi \right)^{-1/2} (2r'R\sin\phi \,\mathrm{d}\phi) = \frac{r'R\sin\phi}{r}$$

so that

$$\sin\phi \, d\phi = \frac{r \, dr}{r'R}$$

It follows that

$$\int_{\phi=0}^{\pi} \frac{\sin \phi \, d\phi}{r} = \frac{1}{r'R} \int_{R-r'}^{R+r'} dr = \frac{2}{R}$$

Substituting this result along with Eq. (E.8) into Eq. (E.9) yields

$$V = -\frac{GMm}{R} \tag{E.10}$$

We conclude that the gravitational potential energy, and hence (from Eq. E.4) the gravitational force, of a sphere with a spherically symmetric mass distribution M is the same as that of a point mass M located at the center of the sphere.