GYRE Stellar Model (GSM) Format

GSM-format files store data describing a stellar model in an HDF5 file. There are a number of variants of this format, which can be distinguished by the presence and/or value of the version attribute.

Version 0.00

The root group atributes and datasets of version-0.00 GSM files are as follows:

Variable	Object name	(A)ttribute /	Object datatype	Definition
		(D)ataset		
R_*	R_star	A	H5T_IEEE_F64LE	Stellar radius (cm)
M_*	M_star	A	H5T_IEEE_F64LE	Stellar mass (g)
L_*	$L_{ extsf{star}}$	A	H5T_IEEE_F64LE	Stellar luminosity (erg s $^{-1}$)
n	n	A	H5T_STD_I64LE	Number of grid points
r	r	D	H5T_IEEE_F64LE	Radius (cm)
w	w	D	H5T_IEEE_F64LE	$M_r/(M_*-M_r)$
L_r	L_r	D	H5T_IEEE_F64LE	Luminosity $(erg s^{-1})$
p	р	D	H5T_IEEE_F64LE	Total pressure $(dyn cm^{-2})$
T	T	D	H5T_IEEE_F64LE	Temperature (K)
ρ	rho	D	H5T_IEEE_F64LE	Density $(g cm^{-3})$
N^2	N2	D	H5T_IEEE_F64LE	Brunt-Väisälä frequency squared (s^{-2})
Γ_1	${\tt Gamma_1}$	D	H5T_IEEE_F64LE	$(\partial \ln p/\partial \ln \rho)_{\rm ad}$
$\nabla_{ m ad}$	nabla_ad	D	H5T_IEEE_F64LE	$(\mathrm{d}\ln T/\mathrm{d}\ln p)_{\mathrm{ad}}$
δ	delta	D	H5T_IEEE_F64LE	$-(\partial \ln \rho/\partial \ln T)_p$
∇	nabla	D	H5T_IEEE_F64LE	$d \ln T / d \ln p$
κ	kappa	D	H5T_IEEE_F64LE	Opacity $(cm^2 g^{-1})$
κ_T	$kappa_{-}T$	D	H5T_IEEE_F64LE	$(\partial \ln \kappa / \partial \ln T)_{\rho}$
$\kappa_{ ho}$	kappa_rho	D	H5T_IEEE_F64LE	$(\partial \ln \kappa / \partial \ln \rho)_T$
ϵ	epsilon	D	H5T_IEEE_F64LE	Energy generation rate $(\operatorname{erg} s^{-1} \operatorname{g}^{-1})$
ϵ_T	$epsilon_T$	D	H5T_IEEE_F64LE	$(\partial \epsilon/\partial \ln T)_{\rho} (\operatorname{erg} s^{-1} \operatorname{g}^{-1})$
$\epsilon_{ ho}$	epsilon_rho	D	H5T_IEEE_F64LE	$(\partial \epsilon/\partial \ln \rho)_T \ (\text{erg } s^{-1} \text{g}^{-1})$
$\Omega_{ m rot}$	Omega_rot	D	H5T_IEEE_F64LE	Rotation angular velocity $(rad s^{-1})$

Version 1.00

The root group atributes and datasets of version-1.00 GSM files are as follows:

Note that the definitions of κ_T and κ_ρ are slightly different than in previous versions.

Variable	Object name	(A)ttribute /	Object datatype	Definition
		(D)ataset		
version ×100	version	A	H5T_STD_I32LE	100
R_*	R_star	A	H5T_IEEE_F64LE	Stellar radius (cm)
M_*	M_star	A	H5T_IEEE_F64LE	Stellar mass (g)
L_*	L_star	A	H5T_IEEE_F64LE	Stellar luminosity (erg s ⁻¹)
n	n	A	H5T_STD_I64LE	Number of grid points
r	r	D	H5T_IEEE_F64LE	Radius (cm)
M_r	M_r	D	H5T_IEEE_F64LE	Interior mass (g)
L_r	L_r	D	H5T_IEEE_F64LE	Luminosity $(\operatorname{erg} \operatorname{s}^{-1})$
p	р	D	H5T_IEEE_F64LE	Total pressure $(dyn cm^{-2})$
T	T	D	H5T_IEEE_F64LE	Temperature (K)
ρ	rho	D	H5T_IEEE_F64LE	Density $(g cm^{-3})$
N^2	N2	D	H5T_IEEE_F64LE	Brunt-Väisälä frequency squared (s^{-2})
Γ_1	Gamma_1	D	H5T_IEEE_F64LE	$(\partial \ln p/\partial \ln \rho)_{\rm ad}$
$ abla_{ m ad}$	nabla_ad	D	H5T_IEEE_F64LE	$(\mathrm{d} \ln T/\mathrm{d} \ln p)_{\mathrm{ad}}$
δ	delta	D	H5T_IEEE_F64LE	$-(\partial \ln \rho/\partial \ln T)_p$
∇	nabla	D	H5T_IEEE_F64LE	$d \ln T / d \ln p$
κ	kappa	D	H5T_IEEE_F64LE	Opacity $(cm^2 g^{-1})$
κ_T	kappa_T	D	H5T_IEEE_F64LE	$(\partial \kappa/\partial \ln T)_{\rho} \ (\text{cm}^2 \text{g}^{-1})$
$\kappa_{ ho}$	kappa_rho	D	H5T_IEEE_F64LE	$(\partial \kappa/\partial \ln \rho)_T \ (\text{cm}^2 \text{g}^{-1})$
ϵ	epsilon	D	H5T_IEEE_F64LE	Energy generation rate (erg s^{-1} g ⁻¹)
ϵ_T	epsilon_T	D	H5T_IEEE_F64LE	$(\partial \epsilon / \partial \ln T)_{\rho} (\operatorname{erg} s^{-1} \operatorname{g}^{-1})$
$\epsilon_{ ho}$	epsilon_rho	D	H5T_IEEE_F64LE	$(\partial \epsilon/\partial \ln \rho)_T (\operatorname{erg} s^{-1} \operatorname{g}^{-1})$
$\Omega_{ m rot}^{'}$	Omega_rot	D	H5T_IEEE_F64LE	Rotation angular velocity $(rad s^{-1})$