

# Measurement of $\nu_\mu + \bar{\nu}_\mu$ Charged-Current Single Pion Production Differential Cross Sections on Argon with the MicroBooNE Detector in the NuMI Beam

Analysis Note, Full FHC, RHC, and Combined Exposure

Draft 1.0, 2026-06-09

Ishanee Pophale, Jarek Nowak

*Lancaster University*

## Abstract

We present a measurement of  $\nu_\mu + \bar{\nu}_\mu$  charged-current single charged-pion ( $\text{CC}\pi^\pm$ ) differential cross sections on argon-40 using the full MicroBooNE NuMI exposure, in three configurations: forward horn current (FHC; Runs 1, 2, 4, 5;  $8.857 \times 10^{20}$  protons on target, POT), reverse horn current (RHC; Runs 1, 2, 3, 4a, 4b, 4c;  $11.082 \times 10^{20}$  POT), and their combination ( $19.939 \times 10^{20}$  POT). The analysis is blind: results are extracted on Poisson-thrown central-value fake data. Events are selected requiring a reconstructed muon candidate track and exactly one contained pion candidate track emerging from a fiducial neutrino vertex, with zero reconstructed electromagnetic showers and a muon–pion opening angle less than 2.6 rad. In the measured phase space the selection achieves  $\approx 55\%$  signal purity and  $\approx 15\%$  efficiency (Table 10). Differential cross sections are extracted for each configuration via Wiener-SVD unfolding in five kinematic variables: muon momentum  $p_\mu$ , pion momentum  $p_\pi$ , muon polar angle  $\cos\theta_\mu$ , pion polar angle  $\cos\theta_\pi$ , and the muon–pion opening angle  $\theta_{\mu\pi}$ . Each configuration is normalised with its own integrated  $\nu_\mu + \bar{\nu}_\mu$  flux (FHC  $6.81159$ , RHC  $6.44646$ , combined  $6.60865 \times 10^{-10} \text{ cm}^{-2} \text{ POT}^{-1}$ ; new-G4 GEANT4 values above 60 MeV, Sec. 2); the RHC value is lower because its beam is  $\bar{\nu}_\mu$ -dominant. The resulting Wiener-SVD integrated cross sections are of order  $0.8\text{--}1.2 \times 10^{-38} \text{ cm}^2/\text{Ar}$  (Table 25), with RHC consistent with (or slightly above) FHC; an earlier apparent RHC deficit was a per-run POT normalization bug affecting the multi-file RHC runs, now corrected. For the current fake-data extraction the injected central-value sample corresponds to  $\approx 0.75\times$  the standalone GENIE v3 prediction in this phase space, consistent with the known over-prediction of  $\text{CC}\pi^\pm$  production in current neutrino interaction generators. No beam-on data is unblinded in this note: every “data” point shown is Poisson-thrown from the central-value Monte Carlo (§3), so no statement here is a measurement of a real deficit. The extraction is validated by a same-sample self-closure that recovers the truth to within  $\approx 6\%$  for every observable, and cross-checked against the D’Agostini iterative-Bayesian method: its flat, iteration-stable integral ( $\sim 20\text{--}40\%$  above Wiener-SVD, config-dependent) confirms that the observable-to-observable spread of the Wiener-SVD integrated cross sections is the additional-smearing matrix  $A_C$ , not a physical

or acceptance effect (Sec. 8.14). All results use the XSECANALYZER framework on the resolution-optimised binning of Sec. 14.

## Contents

|          |                                                                                    |           |
|----------|------------------------------------------------------------------------------------|-----------|
| <b>1</b> | <b>Introduction</b>                                                                | <b>1</b>  |
| <b>2</b> | <b>Beam and Neutrino Flux</b>                                                      | <b>1</b>  |
| 2.1      | NuMI beam and flux composition . . . . .                                           | 1         |
| 2.2      | Beam direction in detector coordinates . . . . .                                   | 2         |
| 2.3      | Data exposure . . . . .                                                            | 3         |
| 2.4      | Integrated flux . . . . .                                                          | 4         |
| <b>3</b> | <b>Monte Carlo Samples</b>                                                         | <b>5</b>  |
| <b>4</b> | <b>Signal Definition</b>                                                           | <b>7</b>  |
| <b>5</b> | <b>Fiducial Volume</b>                                                             | <b>9</b>  |
| 5.1      | Which volume applies where . . . . .                                               | 9         |
| <b>6</b> | <b>Event Selection</b>                                                             | <b>10</b> |
| 6.1      | BDT-based particle identification . . . . .                                        | 10        |
| 6.2      | Cut definitions . . . . .                                                          | 10        |
| 6.3      | Kinematic reconstruction . . . . .                                                 | 12        |
| <b>7</b> | <b>Cut-Flow and Selection Performance</b>                                          | <b>12</b> |
| 7.1      | Key N−1 distributions . . . . .                                                    | 14        |
| 7.2      | Pion proton-rejection variables . . . . .                                          | 15        |
| 7.3      | Final-cut data/MC distributions . . . . .                                          | 15        |
| 7.4      | Background decomposition . . . . .                                                 | 17        |
| 7.5      | Event displays . . . . .                                                           | 17        |
| <b>8</b> | <b>Unfolding</b>                                                                   | <b>18</b> |
| 8.1      | The extraction chain, stage by stage . . . . .                                     | 18        |
| 8.2      | Signal extraction . . . . .                                                        | 21        |
| 8.3      | Response matrix . . . . .                                                          | 21        |
| 8.4      | Regularised unfolding: the bias–variance trade-off . . . . .                       | 22        |
| 8.5      | Wiener-SVD unfolding . . . . .                                                     | 23        |
| 8.6      | Reconstructed spectra and response matrices . . . . .                              | 23        |
| 8.7      | Reconstruction resolution . . . . .                                                | 27        |
| 8.8      | The pion-momentum bias and why $p_\pi$ is the worst-conditioned observable . . . . | 28        |
| 8.9      | Model comparison: $\chi^2$ against truth, tune and generators . . . . .            | 38        |
| 8.10     | Proton-tagged $\chi^2$ and the $A_C$ correlation . . . . .                         | 40        |
| 8.11     | The $W$ binnings are finer than the resolution . . . . .                           | 41        |
| 8.12     | What limits $W_{\pi p}$ , and what can be done about it . . . . .                  | 42        |
| 8.13     | Survey of pion momentum estimators without a magnetic field . . . . .              | 43        |
| 8.14     | Cross-check methods . . . . .                                                      | 46        |

|           |                                                                                            |           |
|-----------|--------------------------------------------------------------------------------------------|-----------|
| <b>9</b>  | <b>Full-exposure results: FHC, RHC, and combined</b>                                       | <b>47</b> |
| 9.1       | Total flux-averaged cross section (cut-and-count) . . . . .                                | 51        |
| 9.2       | Angular variable: $\cos \theta_\mu$ versus $\theta_\mu$ . . . . .                          | 52        |
| 9.3       | Background-control sidebands . . . . .                                                     | 58        |
| 9.4       | Control regions for the proton-tagged subsample . . . . .                                  | 60        |
| <b>10</b> | <b>Proton-tagged subsample: hadronic invariant mass and transverse kinematic imbalance</b> | <b>61</b> |
| 10.1      | Selection performance and cut-flow . . . . .                                               | 61        |
| 10.2      | Additional smearing in the proton-tagged observables . . . . .                             | 68        |
| <b>11</b> | <b>Systematic Uncertainties</b>                                                            | <b>69</b> |
| <b>12</b> | <b>Known Issues and Required Follow-up</b>                                                 | <b>70</b> |
| <b>13</b> | <b>Analysis Pipeline</b>                                                                   | <b>70</b> |
| <b>14</b> | <b>Binning Optimisation, Systematic Breakdown, and Threshold Validation</b>                | <b>71</b> |
| 14.1      | Momentum-threshold validation . . . . .                                                    | 71        |
| 14.2      | Binning optimisation . . . . .                                                             | 71        |
| 14.3      | Systematic breakdown by source . . . . .                                                   | 71        |
| 14.4      | Anatomy of the flux term: genuine flux vs. unfolding amplification . . . . .               | 72        |
| 14.5      | Multi-run overlay and a resonance-model closure . . . . .                                  | 73        |
| 14.6      | Consistency with the sibling $\nu_e$ CC $1\pi$ measurement . . . . .                       | 74        |
| <b>15</b> | <b>Summary</b>                                                                             | <b>75</b> |
| <b>A</b>  | <b>Identity closure: unfolding the prediction with itself</b>                              | <b>76</b> |
| <b>B</b>  | <b>Data/MC Normalisation</b>                                                               | <b>78</b> |
| B.1       | Unfolding covariance ( $\sim 26\%$ effect) . . . . .                                       | 78        |
| B.2       | EXT (cosmic) subtraction for fake data . . . . .                                           | 78        |
| B.3       | Opening-angle binning (review item C4) . . . . .                                           | 78        |
| B.4       | Generator normalisation (NuWro vs GENIE) . . . . .                                         | 78        |
| B.5       | Software trigger (verified to be a no-op) . . . . .                                        | 79        |
| <b>C</b>  | <b>NuWro Fake-Data Closure and Cross-Pipeline Validation</b>                               | <b>79</b> |
| C.1       | Extraction-bias investigation and algorithm equivalence . . . . .                          | 81        |
| <b>D</b>  | <b>Observable Binning</b>                                                                  | <b>82</b> |
| <b>E</b>  | <b>Selection Cut Parameter Summary</b>                                                     | <b>82</b> |

# 1 Introduction

Charged-current single charged-pion ( $\text{CC}\pi^\pm$ ) production,

$$\nu_\mu(\bar{\nu}_\mu) + \text{Ar} \longrightarrow \mu^\mp + \pi^\pm + X, \quad (1)$$

is among the most frequent neutrino interaction channels at energies of 1–5 GeV and constitutes a significant background for oscillation analyses at the same energy scale, most notably for  $\nu_e$  appearance measurements at experiments such as NOvA [3] and T2K [4], as well as the future DUNE [5]. Despite its importance, the cross section on nuclear targets remains poorly constrained: major neutrino interaction generators (GENIE v3, NuWro, NEUT) disagree by 20–30% in absolute rate and show significant shape differences for individual kinematic variables.

MicroBooNE is a single-phase liquid argon time-projection chamber (LArTPC) at Fermilab that operated from 2015–2021, providing the world’s highest-statistics  $\nu_\mu$ -on-argon dataset below 5 GeV during its NuMI running period. This note presents a simultaneous measurement of five differential cross sections in the  $\text{CC}\pi^\pm$  channel using the full NuMI exposure in FHC, RHC, and combined configurations, employing Pandora pattern recognition [6] and Wiener-SVD unfolding [7].

## 2 Beam and Neutrino Flux

### 2.1 NuMI beam and flux composition

The NuMI beam is produced by 120 GeV protons from the Fermilab Main Injector striking a graphite target. Charged secondary hadrons are focused by two magnetic horns whose polarity selects the sign of the focused mesons. In Forward Horn Current (FHC) mode the horns focus positive pions and kaons, giving a beam with predominantly  $\nu_\mu$  content; in Reverse Horn Current (RHC) mode the polarity is reversed to focus negatives, giving a predominantly  $\bar{\nu}_\mu$  beam. MicroBooNE is located 135 mrad ( $7.71^\circ$ ) off-axis from the NuMI beam axis<sup>1</sup>, reducing the peak neutrino energy to the 0.5–2.5 GeV range and narrowing the energy spectrum. Because MicroBooNE sits off-axis and at comparatively low energy, the horn sign-selection is imperfect and both modes carry a sizeable wrong-sign component, most notably RHC, whose wrong-sign  $\nu_\mu$  fraction is large.

The flux composition at the MicroBooNE site is taken from the new-G4 central-value flux histograms (`flux_newg4.root`), integrated over  $E_\nu > 60$  MeV to exclude an unphysical sub-threshold pile-up near  $E_\nu = 0$  (removed by the flux group’s `RemoveFluxPeak` step; below the reaction threshold and irrelevant to the measurement). The muon-flavour split of these central-value histograms matches the author’s authoritative integrated fluxes exactly (Sec. 2.4):

Table 1: NuMI flux composition at the MicroBooNE site by neutrino type, for the FHC and RHC horn modes (new-G4 central-value flux `flux_newg4.root`, integrated over  $E_\nu > 0.1$  GeV). FHC is  $\nu_\mu$ -dominant; RHC is  $\bar{\nu}_\mu$ -dominant with a large wrong-sign  $\nu_\mu$  fraction. The combined row weights each mode by its data exposure (FHC 8.857, RHC  $11.082 \times 10^{20}$  POT; Table 2), giving a near-balanced  $\nu_\mu/\bar{\nu}_\mu$  mix. The signal is charge-blind ( $|\nu_{\text{pdg}}| = 14$ ), so both  $\nu_\mu$  and  $\bar{\nu}_\mu$  contribute in every mode.

| Horn mode | $\nu_\mu$ | $\bar{\nu}_\mu$ | $\nu_e$ | $\bar{\nu}_e$ |
|-----------|-----------|-----------------|---------|---------------|
| FHC       | 63.4%     | 34.0%           | 1.7%    | 0.9%          |
| RHC       | 44.2%     | 53.3%           | 1.4%    | 1.2%          |
| Combined  | 53.0%     | 44.4%           | 1.5%    | 1.1%          |

<sup>1</sup>From the detector position in beam coordinates, (5502, 7259, 67270) cm: 91.1 m transverse at 672.7 m downstream. Earlier versions of this note quoted 110 mrad. See §2.2.

Flux corrections from the Package to Predict the FluX (PPFX) [8] are applied per event via the central-value weight `ppfx_cv` stored in the analysis ntuples. (The absolute flux normalisation used for the cross-section in Sec. 2.4 is still taken from the previous flux histograms; moving that normalisation onto the new-G4 flux is a separate, tracked update.)

## 2.2 Beam direction in detector coordinates

Two different angles describe the MicroBooNE–NuMI geometry, and they must not be confused (Fig. 1). The *off-axis angle* is a property of the beamline: the detector sits 91.1 m off the NuMI axis at 672.7 m downstream of the target, a baseline of 678.8 m and an off-axis angle of  $7.71^\circ$  (134.6 mrad).<sup>2</sup> That angle says nothing about how the TPC is oriented.

The angle that enters the measurement is a different one: the direction in which the neutrinos travel *relative to the detector  $z$  axis*. Averaged over interactions in the active volume, the true neutrino direction in detector coordinates is (0.4624, 0.0489, 0.8853), which lies  $28.6^\circ$  away from  $z$ . The detector  $z$  axis is therefore a poor stand-in for the beam: it is not the direction the neutrinos came from, nor the NuMI beamline axis (itself  $23.2^\circ$  from  $z$ ).




Figure 1: The NuMI geometry seen by MicroBooNE. (a) Beamline geometry, to scale: the detector sits 91.1 m off the NuMI axis at a baseline of 678.8 m, an off-axis angle of  $7.7^\circ$ . (b) The same neutrinos in TPC coordinates, to scale (the  $x$ – $z$  projection; the quoted  $28.6^\circ$  is the full three-dimensional angle): the beam arrives  $28.6^\circ$  away from the detector  $z$  axis, so one and the same muon is assigned  $\theta_\mu = 12^\circ$  about  $z$  but  $\theta_\mu = 40.6^\circ$  about the neutrino. The generators mean the second. (c) The resulting shift in the true  $\cos\theta_\mu$  distribution; dotted lines mark the analysis bin edges. Under the two conventions 64.6% of signal events fall in a different bin (44.6% for  $\cos\theta_\pi$ ).

This matters because every polar angle and every transverse-kinematic-imbalance variable is defined with respect to the neutrino direction. In the generator predictions the neutrino defines  $+z$  by construction — the GENIE event records used here have  $p_x^\nu = p_y^\nu = 0$  and  $p_z^\nu = E_\nu$  exactly — so a measurement that refers its angles to the detector  $z$  axis is not being compared with the prediction it appears to be compared with. The effect is not small: 64.6% of signal events change  $\cos\theta_\mu$  bin between the two conventions, and 44.6% change  $\cos\theta_\pi$  bin. For the TKI variables the axis error exceeds the observable itself, the mean  $\delta p_T$  being 0.742 GeV/ $c$  about  $z$  against 0.264 GeV/ $c$  about the beam.

Accordingly, all angular and TKI observables in this analysis are referred to the neutrino direction, following the  $\beta$  convention of the NuMI  $\nu_e$  CC1 $\pi$  internal note [1] and of the earlier NuMI  $\nu_e$  cross-section measurement it derives from [2]. At truth level the exact per-event

<sup>2</sup>Computed from the detector position in beam coordinates, (5502, 7259, 67270) cm, taken from the same rotation matrix used for the NuMI beamline geometry weights.

neutrino direction is used, which reproduces the generator convention exactly. At reconstruction level the neutrino direction is approximated by the vector from the NuMI target — at  $(-31388, -3316, -60100)$  cm in detector coordinates, a 684 m baseline — to the reconstructed interaction vertex. Adopting the same  $\beta$  as the  $\nu_e$  analysis keeps the two NuMI measurements directly comparable.

The choice of reconstruction-level axis is in any case not a sensitive one. A single fixed axis and the per-event  $\beta$  differ by a median of 2.4 mrad (maximum 7.3 mrad), which moves only 0.42% of muons across a  $\cos\theta_\mu$  bin edge — to be compared with the 64.6% moved by the detector- $z$  convention. Against the true per-event direction the two perform almost identically, with a mean opening angle of  $1.95^\circ$  for  $\beta$  and  $2.00^\circ$  for a fixed axis. What neither can recover is the physical divergence of the beam: neutrinos reaching the detector come from decays distributed along the decay pipe, not from the target, and over a 684 m baseline a 2.5 m detector subtends only  $\sim 4$  mrad, so the vertex position carries almost no information about the direction. That residual is a resolution effect, absorbed by the response matrix like any other.

Quantities that are invariant under this choice of axis are unaffected:  $p_\mu$ ,  $p_\pi$ , the muon-pion opening angle  $\theta_{\mu\pi}$ , the invariant mass  $W_{\pi p}$ , and all integrated cross sections. The event selection is likewise unaffected, as no selection cut and no BDT input variable uses a polar angle.

### 2.3 Data exposure

The analysis is *blind*: the real beam-on data are not used. Each configuration is extracted on Poisson-thrown central-value fake data, generated and scaled *per run* to that run’s data POT (Table 2). The FHC measurement uses the full FHC exposure (Runs 1, 2, 4 and 5); RHC uses Runs 1, 2, 3, 4a, 4b and 4c; the combined measurement uses all fourteen overlay files. Every run is normalised independently to its own data exposure, so the prediction carries the data-POT mixture of run periods and the framework is ready for real per-run beam-on data (each fake-data sample is a drop-in replacement for its beam-on counterpart).

Table 2: Run-period exposure accounting (final, full FHC+RHC). “MC POT” is each period’s native simulated exposure; “Data POT” is the per-run exposure to which the (blind, per-run) fake data is scaled. *Each run is normalised independently* by its own factor,  $\text{Scale} = (\text{run Data POT})/(\text{run MC POT})$ , so the prediction is the data-POT–weighted mixture of run periods; the normalisation a real per-run beam-on dataset produces. This replaces an earlier single per-mode pooling factor: because the per-run MC/data POT ratios vary by up to a factor of three (Scale ranges 0.051–0.141 within FHC alone), a single factor would weight the runs by their MC statistics rather than by the data exposure. The RHC Run 4 row sums the 4a/4b/4c sub-files and Run 3 sums aa–ae; sub-files sharing a run’s data POT are scaled by the group MC total so they pool within that run. The combined measurement applies each run’s own factor. Run 5 uses the reweighted-PPFX  $\nu_\mu$  overlay. The scheme is validated by the FHC closure (unfolded vs. smeared truth  $\chi^2 = 1.6/7$ ,  $p = 0.98$ ).

| Run period                | MC POT ( $\times 10^{20}$ ) | Data POT ( $\times 10^{20}$ ) | Scale          |
|---------------------------|-----------------------------|-------------------------------|----------------|
| Run 1 (FHC)               | 23.282                      | 3.283                         | 0.1410         |
| Run 2 (FHC)               | 24.934                      | 1.268                         | 0.0509         |
| Run 4 (FHC)               | 28.335                      | 2.075                         | 0.0732         |
| Run 5 (FHC)               | 19.300                      | 2.231                         | 0.1156         |
| <b>FHC total</b>          | <b>95.85</b>                | <b>8.857</b>                  | <i>per run</i> |
| Run 1 (RHC)               | 8.997                       | 0.605                         | 0.0673         |
| Run 2 (RHC)               | 57.865                      | 2.591                         | 0.0448         |
| Run 3 (RHC, aa–ae)        | 55.185                      | 5.003                         | 0.0907         |
| Run 4 (RHC, 4a–4c)        | 32.586                      | 2.883                         | 0.0885         |
| <b>RHC total</b>          | <b>154.63</b>               | <b>11.082</b>                 | <i>per run</i> |
| <b>Combined (FHC+RHC)</b> | <b>250.49</b>               | <b>19.939</b>                 | <i>per run</i> |

The full exposure substantially reduces the data-statistical term (Tables 31–33), leaving the measurement flux- and detector-systematic limited. When the analysis is unblinded, the real beam-on POT (and gate counts) replace the data-POT values above.

## 2.4 Integrated flux

The absolute flux normalisation uses the new-G4 GEANT4 NuMI flux (`flux_newg4.root`), integrated over  $E_\nu > 60$  MeV (the flux group’s `RemoveFluxPeak` cut, which removes an unphysical  $\sim 25$ – $30$  MeV artifact spike). The signal is charge-blind ( $|\nu_{\text{pdg}}| = 14$ , covering both  $\nu_\mu \rightarrow \mu^- \pi^+$  and  $\bar{\nu}_\mu \rightarrow \mu^+ \pi^-$ ), so the normalisation uses the combined  $\nu_\mu + \bar{\nu}_\mu$  flux  $\Phi = \Phi_{\nu_\mu} + \Phi_{\bar{\nu}_\mu}$ . Because RHC is  $\bar{\nu}_\mu$ -dominant with a different absolute flux than FHC, each configuration uses its *own* integrated flux:

Table 3: Authoritative integrated  $\nu_\mu$  and  $\bar{\nu}_\mu$  fluxes per POT (new-G4 flux,  $E_\nu > 60$  MeV), and the combined  $\nu_\mu + \bar{\nu}_\mu$  value used as the per-configuration normalisation. The combined row is the data-POT-weighted average of the two modes (weights 8.857 : 11.082; Table 2). The FHC value  $6.81159 \times 10^{-10}$  is the previously confirmed normalisation; RHC and combined were previously normalised with the FHC flux and are corrected here (raising the RHC cross section by 5.7% and the combined by 3.1%).

| Config   | $\Phi_{\nu_\mu}$<br>( $10^{-10} \text{ cm}^{-2} \text{ POT}^{-1}$ ) | $\Phi_{\bar{\nu}_\mu}$ | $\Phi_{\nu_\mu + \bar{\nu}_\mu}$ | $\bar{\nu}_\mu$ frac. |
|----------|---------------------------------------------------------------------|------------------------|----------------------------------|-----------------------|
| FHC      | 4.43515                                                             | 2.37644                | 6.81159                          | 34.9%                 |
| RHC      | 2.92348                                                             | 3.52298                | 6.44646                          | 54.6%                 |
| Combined | 3.59497                                                             | 3.01368                | 6.60865                          | 45.6%                 |

The mean  $\bar{\nu}_\mu$  energy is 0.500 GeV (FHC) and 0.417 GeV (RHC), the lower RHC value reflecting the softer wrong-sign-enhanced spectrum. These fluxes are set per configuration in `integrated_numu_flux_in_FV()` (`include/XSecAnalyzer/FiducialVolume.hh`). The correction relative to the superseded flux (`uboone_numi_flux_histograms.root`, whose combined value  $2.3725 \times 10^{-9}$  was  $\sim 3.48\times$  too large and made every extracted cross section  $\sim 3.5\times$  too small) was validated four independent ways, standalone NuWro and GENIE on the old vs. corrected flux, the flux author’s numbers, and a GENIE closure, documented in `generator_predictions/`.

The number of argon target nuclei in the fiducial volume is computed from first principles as

$$N_{\text{Ar}} = \frac{\rho_{\text{LAr}} \cdot V_{\text{FV}} \cdot N_A}{A_{\text{Ar}}} = 8.710 \times 10^{29}, \quad (2)$$

with  $\rho_{\text{LAr}} = 1.3836 \text{ g/cm}^3$ ,  $A_{\text{Ar}} = 39.948 \text{ g/mol}$  and  $N_A = 6.02214076 \times 10^{23} \text{ mol}^{-1}$ . The fiducial box is  $10 < x < 246$ ,  $-101 < y < 101$ ,  $10 < z < 986 \text{ cm}$  ( $4.6528 \times 10^7 \text{ cm}^3$ ), from which the  $675.1 < z < 775.1 \text{ cm}$  dead region is subtracted, leaving  $V_{\text{FV}} = 4.1761 \times 10^7 \text{ cm}^3$ . Without that subtraction the same expression would give  $9.705 \times 10^{29}$ . This is the value the code uses, computed by `num_Ar_targets_in_FV(FV, 675.1, 775.1)` in `CrossSectionExtractor.hh`, and it is the value entering the conversion factor of §8 and the cut-and-count normalisation of §9.1. and the per-configuration normalisation factor is  $N_{\text{norm}} = \Phi_{\text{mode}} \cdot N_{\text{POT,mode}} \cdot N_{\text{Ar}}$ , evaluated with each mode’s flux and data POT (Table 2).

### 3 Monte Carlo Samples

Monte Carlo events are generated with GENIE v3 [9] on argon-40 and overlaid with real cosmic-ray data (the “overlay” technique) so that simulated events experience in-situ detector noise and cosmogenic activity. Table 4 summarises the input samples.



Table 4: Input samples (final, full FHC+RHC). “ $n$ ” is the number of overlay files; “MC POT” is the summed native simulated exposure ( $\times 10^{20}$ ). The blind Poisson-thrown fake data stands in for the (unused) beam-on data; the detector variations use the native Run-4 FHC and RHC samples. The per-file lists and POT/trigger values are in the `configs/file_properties_numi_*` tables.

| Sample                  | Composition                                      | $n$ | MC POT |
|-------------------------|--------------------------------------------------|-----|--------|
| $\nu_\mu$ overlay (FHC) | Runs 1, 2, 4 (pandora) + Run 5 (reweighted-PPFX) | 4   | 95.85  |
| $\nu_\mu$ overlay (RHC) | Runs 1, 2, 3, 4a, 4b, 4c (pandora)               | 10  | 154.63 |
| Dirt MC                 | GENIE NuMI dirt overlay (Run 1)                  | 1   | 16.74  |
| EXT (beam-off)          | cosmic data, per run, horn-mode-independent      | 7   | n/a    |
| Detector variations     | Run-4 FHC + Run-4 RHC, CV + 8 knobs each         | 18  | n/a    |
| Fake data (blind)       | Poisson-thrown central value, per configuration  | n/a | n/a    |
| Beam-on data            | blinded, not used until unblinding               | n/a | n/a    |

**Event weights.** Each MC event is weighted by

$$w = \text{Scale}[\text{run}] \times w_{\text{PPFX}} \times w_{\text{tune}}, \quad (3)$$

where  $\text{Scale}[\text{run}] = (\text{run Data POT})/(\text{run MC POT})$  is the *per-run* POT factor of Table 2: each run period is scaled independently to its own data exposure, so the prediction reproduces the data-POT mixture of runs rather than the MC-statistics mixture (see the caption of Table 2). Operationally this is exactly what the framework does natively each overlay file carries its run’s data POT via a matching per-run beam-on (here, per-run fake-data) sample, and is scaled by `run_bnb_pot/summed_pot`.  $w_{\text{PPFX}}$  is the PPFX central-value flux correction (`ppfx_cv`) and  $w_{\text{tune}}$  is the GENIE/GEANT4 reweighting correction (`weightTune`). Events with non-finite, zero, or unphysically large weights ( $w > 30$ ) are assigned unit weight. At unblinding the per-run fake-data samples are replaced by the corresponding per-run beam-on files (same runs, same POT), so the normalisation is already correct.

**EXT scaling.** The same combined beam-off (cosmics) sample is attached to every run and scaled by that run’s beam-on/beam-off trigger ratio, `bnb_trigs[run]/ext_gates`. Because the beam-on/beam-off ratio is run-independent, the per-run contributions sum to the mode-total scale; for the full FHC exposure,

$$\text{Scale}_{\text{EXT}}^{\text{FHC}} = \frac{\sum_{\text{run}} \text{bnb\_trigs}[\text{run}]}{\text{ext\_gates}} = \frac{22\,667\,109}{3\,821\,593} = 5.93. \quad (4)$$

(The framework requires each run carrying a beam-on sample to carry a beam-off sample as well; the single beam-off file is therefore attached per run through symlinks so that run bookkeeping is complete without duplicating the data.) The uncertainty on this ratio is a leading systematic on the EXT background.

**A directional cut against residual cosmics: considered and rejected.** Part of the surviving beam-off sample is thought to be broken cosmic muon tracks, where the break is reconstructed as a vertex and the track direction bears no relation to it. That leaves a directional signature, and it is present in the data: among selected events the beam-off sample is enriched at backward reconstructed muon angles by nearly an order of magnitude relative to signal.

| reco $\cos \theta_\mu$ | EXT (beam-off) | MC signal | enrichment  |
|------------------------|----------------|-----------|-------------|
| $< 0$                  | 31.8%          | 4.1%      | $7.8\times$ |
| $< -0.5$               | 9.1%           | 1.4%      | $6.4\times$ |

The enrichment is real but the arithmetic does not support a cut. At the final cut the FHC sample is 974 signal, 128 EXT and 1 746 predicted in total (Tab. 9). A  $\cos\theta_\mu < 0$  requirement would remove  $\approx 41$  EXT events and  $\approx 40$  signal events, close to one for one. Because the beam-off sample is subtracted from the data directly, and that subtraction is unbiased by construction, removing part of it buys only a reduction in the subtraction’s *statistical* uncertainty, from  $\sqrt{128} \approx 11$  to  $\sqrt{87} \approx 9$  events. Spending 40 signal events to gain  $\approx 2$  events of uncertainty is a net loss.

The same bound applies to any cosmic cut, not just this one: EXT is 7.3% of the selected sample and is already subtracted without bias, so removing *all* of it for *zero* signal loss would gain about 11 events of statistical uncertainty on a sample of 1 746. Any cut costing more than about 1% of the signal is a net loss regardless of how efficiently it tags cosmics. The residual beam-off contribution is therefore left in place and handled by subtraction.

A more surgical variant remains untested. The CC0 $\pi$  analysis tags mis-directed tracks with the calorimetric flag `trk.bragg_mu_fwd_preferred_v` in coincidence with a poor forward-muon fit,  $\chi_\mu^2 > 6$ , which fires on tracks whose  $dE/dx$  profile disagrees with the assigned direction rather than on backward-going tracks as such. That would not cut genuine backward muons, which outnumber mis-reconstructed ones here by roughly 8:1. Both branches exist in the input ntuples but are not currently propagated by `ProcessNTuples`; the study is scoped in `report/BACKWARD_MUON_PLAN.md`. It is bounded by the same 11-event ceiling, so it is worth measuring but cannot become a significant improvement.

**Software trigger.** Data and EXT files are pre-filtered to require the software trigger (`swtrig == 1`). MC events failing `swtrig = 0` are removed at the NeutrinoSlice cut stage. In practice this has no measurable effect at the current working point: all MC events passing the topological cut are verified to have `swtrig == 1` (46 810/46 810 events confirmed numerically).

## 4 Signal Definition

The signal is defined at generator level and is used only to construct the selection efficiency and response matrix. It is *never* applied as a reconstruction cut. The definition is *charge-blind*: both  $\nu_\mu \rightarrow \mu^- \pi^+$  and  $\bar{\nu}_\mu \rightarrow \mu^+ \pi^-$  are treated as signal on the same footing.

### Neutrino flavour

$$|\nu \text{ PDG}| = 14 \text{ } (\nu_\mu \text{ or } \bar{\nu}_\mu).$$

### Interaction type

Charged-current (`ccnc == 0`).

### True vertex

Inside the fiducial volume (Section 5).

### Primary multiplicity

Exactly one muon ( $|\text{PDG}| = 13$ ), exactly one charged pion ( $|\text{PDG}| = 211$ ), zero neutral pions ( $|\text{PDG}| = 111$ ), zero kaons. Any number of protons is allowed.

### Kinematic thresholds

$p_\mu > 0.15 \text{ GeV}/c$ ,  $p_\pi > 0.175 \text{ GeV}/c$ , and  $\theta_{\mu\pi} < 2.6 \text{ rad}$ . These define the *measured phase space*, chosen from the efficiency turn-on curves (Section 7). The momentum efficiency turns on sharply at  $\sim 0.15\text{--}0.20 \text{ GeV}/c$  (set by the 10 cm muon and 20 cm pion track-length cuts); below that it is only  $\sim 1\text{--}3\%$  with a strongly model-dependent correction. The opening-angle upper bound (2.6 rad) excludes the large-angle region, which is dominated by cosmic and dirt background ( $\sim 7\%$  purity in the  $\theta_{\mu\pi} > 2.513 \text{ rad}$  bin). No *lower* opening-angle cut is applied: the small-angle region is high-purity ( $\sim 58\%$ ), so cutting it would only remove

signal (it is low-efficiency but not low-purity). The same cut is applied at reco level so the signal and selection acceptances match. The first  $p_\mu/p_\pi$  bins and the  $\theta_{\mu\pi}$  upper edge start/end at these thresholds.

(Earlier iterations used  $p_\mu, p_\pi > 0.1 \text{ GeV}/c$  with the full  $0-\pi$  angle; that left near-dead efficiency bins at low momentum and at the large-angle extreme. The phase space above raises the efficiency from 12.1% to 15.6% at  $\sim 55\%$  purity, see Section 7.)

**Pion threshold and the sibling  $\nu_e$  measurement.** The companion MicroBooNE NuMI  $\nu_e$  CC  $1\pi$  analysis defines its pion signal by kinetic energy,  $\text{KE}_\pi > 40 \text{ MeV}$ , equivalent to  $p_\pi > 0.113 \text{ GeV}/c$ . Our  $p_\pi > 0.175 \text{ GeV}/c$  ( $\text{KE}_\pi > 84 \text{ MeV}$ ) is therefore a higher threshold. Lowering it toward the  $\nu_e$  value is possible, the signal definition is a truth-level choice and the unfolding maps reco to truth but the pion momentum reconstruction (track range under the muon hypothesis, mass-corrected) degrades rapidly below  $0.175 \text{ GeV}/c$ :

Table 5: Pion momentum reconstruction versus true  $p_\pi$  (correctly reconstructed signal pions). Below the  $0.175 \text{ GeV}/c$  threshold the bias and resolution blow up and the reconstructed fraction halves.

| True $p_\pi$ [GeV/c]           | Bias $\langle(p^{\text{reco}} - p^{\text{true}})/p^{\text{true}}\rangle$ | RMS resolution | Reco fraction |
|--------------------------------|--------------------------------------------------------------------------|----------------|---------------|
| 0.080–0.113                    | +131%                                                                    | 73%            | 3.4%          |
| 0.113–0.150                    | +66%                                                                     | 56%            | 3.4%          |
| 0.150–0.175                    | +30%                                                                     | 51%            | 4.3%          |
| <b>0.175–0.250</b> (threshold) | <b>−8%</b>                                                               | <b>31%</b>     | <b>10.0%</b>  |
| 0.250–0.400                    | −30%                                                                     | 21%            | 10.4%         |

At the threshold the resolution is 31%; in the  $[0.113, 0.175] \text{ GeV}/c$  region it is 51–56% with a +30–66% bias and only  $\sim 3$ –4% reconstructability. Extending the measurement to the  $\nu_e$  threshold would thus add bins carrying large, strongly model-dependent unfolding systematics. For consistency with the  $\nu_e$  measurement the threshold could be lowered to  $0.113 \text{ GeV}/c$  with the  $[0.113, 0.175] \text{ GeV}/c$  region kept in a single coarse bin to contain those systematics, or both phase spaces reported; this is a systematics-versus-cross-analysis-consistency trade rather than a reconstruction limitation.

**The low-momentum bin, added.** We have implemented this extension for  $p_\pi$ : a  $[0.113, 0.175] \text{ GeV}/c$  bin is prepended to the pion-momentum binning, lowering the measured phase space to the  $\nu_e$  threshold. The truth-level signal below  $0.175 \text{ GeV}/c$  is recovered from the stored signal-component branches (the reconstruction reproduces **MC\_Signal** above  $0.175 \text{ GeV}/c$  to 0.4%, the residual being the unstored kaon veto); 3323 true signal events populate the new bin. Its selection efficiency is 2.2%; an order of magnitude below the 15–17% of the standard bins (Table 6), as anticipated from the reconstruction turn-on. Despite this, the Wiener-SVD extraction remains well behaved with the bin included: the  $M_A^{\text{RES}}$  fake-data closure gives  $\chi^2/\text{ndf} = 1.41/6$  ( $p = 0.97$ ) and the flux term is unchanged (17% reco, 18.4% unfolded, amplification  $1.08\times$ ). The bin therefore adds to the measurement, at the price of a large per-bin uncertainty driven by the low efficiency.

The generator predictions have been regenerated on the six-bin scheme (the ppi histogram uses  $p_\pi > 0.113$  while the other observables retain  $p_\pi > 0.175$ , mirroring the analysis). GENIE, NuWro and NEUT all provide the low bin; GiBUU could not be re-binned as its generator-level event record was not retained, so it is omitted from the extended  $p_\pi$  comparison. Against the (fake) data the generators give  $\chi^2/\text{ndf} = 11.1/6$  (GENIE,  $p = 0.09$ ),  $7.0/6$  (NuWro,  $p = 0.33$ ) and  $6.3/6$  (NEUT,  $p = 0.39$ ): NEUT and NuWro describe the extended spectrum well, while GENIE is mildly disfavoured, consistent with the known GENIE over-prediction of low-momentum  $\text{CC}\pi^\pm$  production.

Table 6: Selection efficiency per true  $p_\pi$  bin with the low-momentum bin added. The new  $[0.113, 0.175]$  GeV/c bin sits at 2.2%, an order of magnitude below the standard bins.

| True $p_\pi$ [GeV/c]     | $N_{\text{true}}$ | $N_{\text{reco}}$ | Efficiency  |
|--------------------------|-------------------|-------------------|-------------|
| <b>0.113–0.175</b> (new) | 3323              | 74                | <b>2.2%</b> |
| 0.175–0.250              | 3368              | 526               | 15.6%       |
| 0.250–0.320              | 2190              | 367               | 16.8%       |
| 0.320–0.420              | 2831              | 479               | 16.9%       |
| 0.420–0.550              | 2499              | 410               | 16.4%       |
| 0.550–1.000              | 2933              | 429               | 14.6%       |

## 5 Fiducial Volume

The fiducial volume (FV) is defined in `CC1mu1piXp.cxx` (the independent cross-check selection carries the same numbers in `FV_new.h`):

$$x \in [10.0, 246.0] \text{ cm}, \quad y \in [-101.0, +101.0] \text{ cm}, \quad z \in [10.0, 986.0] \text{ cm}, \quad (5)$$

with the dead-wire region  $z \in [675.1, 775.1]$  cm excluded. The total FV volume is

$$\begin{aligned} V_{\text{FV}} &= (246 - 10) \times 202 \times (976 - 100) \text{ cm}^3 \\ &= 236 \times 202 \times 876 \text{ cm}^3 \approx 4.17 \times 10^7 \text{ cm}^3. \end{aligned} \quad (6)$$

A wider *containment* region (2 cm inset from the physical TPC walls) is used to require that the pion candidate track endpoint is contained:

$$x \in [2.0, 254.35] \text{ cm}, \quad y \in [-113.53, +107.47] \text{ cm}, \quad z \in [2.1, 1034.9] \text{ cm}. \quad (7)$$

### 5.1 Which volume applies where

Three distinct volumes appear above and it is worth stating plainly which is used for what, since “contained” and “in the fiducial volume” are not the same requirement.

| volume        | extent                                                                                | used for                              |
|---------------|---------------------------------------------------------------------------------------|---------------------------------------|
| fiducial (FV) | $[10, 246] \times [-101, 101] \times [10, 986]$ cm<br>minus $z \in [675.1, 775.1]$ cm | neutrino vertex, true <i>and</i> reco |
| containment   | $[2, 254.35] \times [-113.53, 107.47] \times [2.1, 1034.9]$ cm                        | track endpoints                       |
| active TPC    | $[0, 256.35] \times [-115.5, 117.5] \times [0, 1036.9]$ cm                            | (reference only)                      |

**True and reco use identical dimensions.** The framework keeps the two as separate quantities, `true_FV()` and `reco_FV()`, and other selections in the code base use that freedom. Here they are defined as the same box (`CC1mu1piXp.cxx`, `define_true_FV` / `define_reco_FV`). The signal definition tests the *true* neutrino vertex and the selection tests the *reconstructed* vertex, both through the same helper, so both inherit the dead-region veto as well. There is no asymmetry between the two, and the FV therefore does not itself generate any migration between signal and background.

**The dead-region veto applies to the vertex only.** An event whose vertex lies outside the unresponsive-wire slab is kept even if one of its tracks crosses that slab. This is the conventional choice, but it is an assumption rather than a necessity: tracks crossing dead wires have degraded calorimetry and range reconstruction, and the associated inefficiency is absorbed into the response matrix rather than removed by a cut.

**The dead slab is also removed from the target count.** The argon target number is computed from the FV with the slab subtracted, not from the full box. Retaining the slab would inflate the target count by  $100/976 \approx 10\%$  and bias every cross section low by the same factor. Note that this subtraction is applied on the NuMI code path specifically; the generic path computes the target number without it, so the two must not be mixed.

**Relation to the active volume.** The FV is inset from the active TPC by approximately 10 cm in  $x$ , 15 cm in  $y$ , 10 cm at the upstream  $z$  face and 51 cm at the downstream face. The containment box, by contrast, is inset by only 2 cm: track endpoints are required to be inside the detector, not inside the fiducial region, which is the correct requirement for range-based momentum and for the pion-containment cut.

## 6 Event Selection

The event selection is implemented in the framework selection class `src/selections/CC1mu1piXp.cxx` (which produces the response matrices and spectra for all results in this note); the standalone `selection/ccpi_selection.C` pipeline used for the cross-pipeline validation of Appendix C implements the byte-identical cut sequence. The selection comprises 11 sequential cuts.

### 6.1 BDT-based particle identification

Three gradient-boosted decision trees (TMVA), trained on the MicroBooNE overlay MC, supply the calorimetric particle identification used in the muon- and pion-candidate cuts. All three share a common set of per-track inputs: the three Bragg-peak  $dE/dx$  likelihoods under the proton, muon and MIP hypotheses (`trk_bragg_{p,mu,mip}_v`), the log-likelihood-ratio PID score (`trk_llr_pid_score_v`), the Pandora track-vs-shower score (`trk_score_v`), and the space-charge-corrected track end-point coordinates (`trk_sce_end_{x,y,z}_v`):

- **MIP BDT** (`bdt_mip`; weights `dataset_MIP_BDT_no_len`): separates minimum-ionising tracks (muons, pions) from stopping protons. Both the muon and the pion candidate are required to have `bdt_mip > -0.1` (Cuts 4–5).
- **Pion BDT** (`bdt_pi`; weights `dataset_pion_BDT_no_len`): a dedicated pion-versus-proton discriminant applied to the pion candidate, `bdt_pi > -0.1` (Cut 5).
- **Muon BDT** (weights `dataset_muon_BDT`): the same inputs plus the track length (`trk_len_v`). Rather than a threshold, it *rank*s the muon-candidate tracks; the highest-scoring track is chosen as the muon.

The two candidate-PID BDTs deliberately omit track length so the score does not bias the momentum spectra. The weight files are resolved from `$CC1MU1PIXP_BDT_WEIGHTS_DIR`, falling back to the in-tree `booster_decision_tree/` copy. A fourth score, `bdt_cosmic`, is available for cosmic rejection but is disabled by default (no benefit in an A/B test; the Pandora topological score already provides the cosmic rejection at Cut 3).

### 6.2 Cut definitions

**Cut 0. EventsInTrueFV.** All events. Reference denominator for truth-level studies.

**Cut 1. NeutrinoSlice.**  $\geq 1$  reconstructed track (`ntracks > 0`), `bdt_cosmic > 0.0` (currently disabled; no benefit found in A/B test), and `swtrig == 1`.

**Cut 2. InFiducialVol.** The SCE-corrected reconstructed neutrino vertex lies inside the FV.

- Cut 3. Topological.** `topological_score`  $> 0.67$ . This Pandora-produced multivariate score distinguishes cosmic-ray topologies from neutrino-like topologies and provides the primary cosmic rejection: EXT events are reduced by a factor of  $\sim 33$  between Cut 2 and Cut 3.
- Cut 4. MuonCandidate.** At least one primary PFP track satisfies the muon PID criteria of Table 7. The longest qualifying track is the muon candidate.
- Cut 5. ContainedPion.** Exactly one primary track (other than the muon candidate) satisfies the pion PID criteria of Table 8 and has its endpoint inside the containment volume. The track with the highest LLR PID score is the pion candidate.
- Cut 6. MuonIn3Planes.** The muon candidate has  $\geq 1$  hit on each of the U, V, Y planes.
- Cut 7. PionIn3Planes.** The pion candidate has  $\geq 1$  hit on each of the U, V, Y planes.
- Cut 8. ShowerCut.** Zero primary-level EM showers, unless exactly one shower is present with fewer than 50 Y-plane hits and a vertex distance  $> 10$  cm (consistent with a delta-ray or low-energy nuclear photon).
- Cut 9. OpeningAngle.** The reconstructed  $\mu$ - $\pi$  opening angle satisfies  $\theta_{\mu\pi} < 2.6$  rad ( $\approx 149^\circ$ ), rejecting the large-angle cosmic/dirt background. No lower bound is applied (the small-angle region is high-purity). The same cut is imposed on the signal definition (Section 4).
- Cut 10.**  
**Nonprotons** (final cut). Fewer than 4 primary tracks have LLR PID  $> 0.1$  (non-proton-like), and the total number of primary reconstructed tracks is less than 5.

Table 7: Muon candidate PID thresholds (applied at Cut 4).

| Variable                                     | Threshold     | Description               |
|----------------------------------------------|---------------|---------------------------|
| <code>trk_score_v</code>                     | $> 0.80$      | Track-like PFP            |
| Vertex distance                              | $\leq 4.0$ cm | Attached to $\nu$ vertex  |
| Track length                                 | $> 10.0$ cm   | Minimum track length      |
| LLR PID ( <code>trk_llr_pid_score_v</code> ) | $> 0.20$      | Muon-like                 |
| MIP BDT ( <code>bdt_mip</code> )             | $\geq -0.10$  | Minimum-ionising-particle |

Table 8: Pion candidate PID thresholds (applied at Cut 5). Endpoint containment requires the track endpoint inside the containment volume. `trk_bragg_pion_v`: pion-hypothesis Bragg-peak likelihood score (rejects stopping protons).

| Variable                         | Threshold   | Description               |
|----------------------------------|-------------|---------------------------|
| LLR PID                          | $> 0.10$    | Pion / non-proton-like    |
| Vertex distance                  | $< 4.0$ cm  | Attached to $\nu$ vertex  |
| Track length                     | $> 20.0$ cm | Minimum track length      |
| MIP BDT ( <code>bdt_mip</code> ) | $> -0.10$   | MIP-like                  |
| Pion BDT ( <code>bdt_pi</code> ) | $> -0.10$   | Pion-like (vs. proton)    |
| Bragg-pion score                 | $\geq 0.08$ | Proton rejection          |
| Endpoint contained               | yes         | Inside containment volume |

### 6.3 Kinematic reconstruction

**Muon momentum.** A *hybrid* estimator, switched on containment of the muon-candidate track: the range-based momentum `trk_range_muon_mom_v` is used when the track ends inside the detector, and the multiple-Coulomb-scattering momentum `trk_mcs_muon_mom_v` when it exits (range is meaningless for an exiting track). The reported  $p_\mu$  is therefore `candidate_muon_mom_reco`, not either branch alone. In the selected FHC sample the split is 37% contained (range) and 63% exiting (MCS), i.e. the measurement is *MCS-dominated*: the NuMI muons are energetic and the MicroBooNE TPC is small, so most of them leave it. This matters for the  $p_\mu$  resolution and for the detector systematic, since the range and MCS estimators carry different uncertainties.

**Pion momentum.** Taken from the *muon-hypothesis* range momentum (`trk_range_muon_mom_v`). The muon mass (105.7 MeV) is close to the pion mass (139.6 MeV), so the range-to-momentum conversion is a good proxy, and, crucially it is a *momentum*, directly comparable to the true  $p_\pi$  it is binned against. An earlier version of this analysis used the proton-hypothesis range *kinetic energy* (`trk_energy_proton_v`) instead. That is a different physical quantity from the momentum used for the true axis with the same bin edges, so the migration matrix was systematically shifted rather than near-diagonal: a true 0.2 GeV/c pion gives  $KE \approx 0.104$  GeV, two bins low, and a true 0.175 GeV/c pion falls below the first reco bin entirely. Range momentum is meaningful only for contained tracks, which the containment requirement in the selection guarantees.

**Angles.** Polar angles are defined with respect to the *neutrino direction*  $\hat{u}_\nu$ , not the detector  $z$ -axis:  $\cos\theta_\mu = \hat{u}_\mu \cdot \hat{u}_\nu$ ,  $\cos\theta_\pi = \hat{u}_\pi \cdot \hat{u}_\nu$ . The two are far from equivalent — the NuMI beam reaches MicroBooNE  $28.6^\circ$  away from  $z$ , and referring the angles to  $z$  would place 64.6% of signal events in a different  $\cos\theta_\mu$  bin than the generator predictions they are compared with (Sec. 2.2). For reconstructed quantities  $\hat{u}_\nu$  is the  $\beta$  direction of Refs. [1, 2], from the NuMI target to the reconstructed vertex; at truth level it is the exact per-event neutrino direction. The opening angle  $\theta_{\mu\pi} = \arccos(\hat{u}_\mu \cdot \hat{u}_\pi)$  is computed from reconstructed track directions and, being an angle between two measured tracks, is independent of this choice.

## 7 Cut-Flow and Selection Performance

Figure 2 shows the full-exposure cut-flow yields for all three configurations as stacked plots, and Table 9 gives the detailed per-category, per-cut breakdown for each.




Figure 2: Cut-flow yields at *full exposure* for FHC, RHC and combined (log scale; MC-prediction stack: signal blue, background orange, EXT grey, dirt green; colourblind-safe Okabe-Ito), signal and background POT-scaled per mode. Data is blind, so this shows the MC-prediction stack; the detailed per-category breakdown is in Table 9.



Table 9: Cut-flow yields at *full exposure* for FHC, RHC and combined (MC prediction; data is blind, so Pred = Data by construction for the Poisson-thrown fake data). Signal and “Bkg” (all non-signal MC: OOFV, NC, CC1 $\pi^0$ , CC0 $\pi$ , CCN $\pi$ , CCK, CCothers) are from the  $\nu_\mu$  overlays, POT-scaled *per run* (each run by its own  $D_{\text{run}}/MC_{\text{run}}$ , Table 2); EXT (beam-off cosmic) is scaled by the (per-run summed) data/EXT trigger ratio times the 2% NuMI beam-occupancy factor (FHC  $0.98 \times 5.93$ , RHC  $0.98 \times 6.16$ , combined  $0.98 \times 12.09$ ); dirt from the GENIE NuMI dirt overlay. Pred. = Signal + Bkg + EXT + Dirt. Eff. is the cumulative signal efficiency relative to the generated signal (the None stage, where all true signal is counted); Pur. = Signal/Pred.

| Cut                                                                     | Signal       | Bkg MC       | EXT        | Dirt       | Pred.        | Eff. [%]    | Pur. [%]    |
|-------------------------------------------------------------------------|--------------|--------------|------------|------------|--------------|-------------|-------------|
| <i>FHC (Runs 1,2,4,5; <math>8.857 \times 10^{20}</math> POT)</i>        |              |              |            |            |              |             |             |
| None                                                                    | 5 580        | 279 180      | 23 921 822 | 36 142     | 24 242 724   | 100.0       | 0.02        |
| InFiducialVol                                                           | 4 632        | 62 146       | 1 020 804  | 1 890      | 1 089 473    | 83.0        | 0.43        |
| Topological                                                             | 3 501        | 28 040       | 34 481     | 204        | 66 225       | 62.7        | 5.3         |
| MuonCandidate                                                           | 3 143        | 18 980       | 15 206     | 137        | 37 466       | 56.3        | 8.4         |
| ContainedPion                                                           | 1 395        | 2 094        | 1 348      | 25         | 4 862        | 25.0        | 28.7        |
| MuonIn3Planes                                                           | 1 336        | 1 975        | 1 122      | 17         | 4 450        | 23.9        | 30.0        |
| PionIn3Planes                                                           | 1 239        | 1 770        | 854        | 7.0        | 3 871        | 22.2        | 32.0        |
| ShowerCut                                                               | 1 048        | 898          | 558        | 6.2        | 2 510        | 18.8        | 41.7        |
| OpeningAngle                                                            | 1 040        | 813          | 186        | 2.0        | 2 041        | 18.6        | 51.0        |
| <b>Nonprotons</b>                                                       | <b>974</b>   | <b>643</b>   | <b>128</b> | <b>1.3</b> | <b>1 746</b> | <b>17.5</b> | <b>55.8</b> |
| <i>RHC (Runs 1,2,3,4a,4b,4c; <math>1.108 \times 10^{21}</math> POT)</i> |              |              |            |            |              |             |             |
| None                                                                    | 6 130        | 304 347      | 24 837 750 | 28 032     | 25 176 259   | 100.0       | 0.02        |
| InFiducialVol                                                           | 5 124        | 69 102       | 1 059 889  | 1 466      | 1 135 581    | 83.6        | 0.45        |
| Topological                                                             | 3 960        | 31 629       | 35 801     | 158        | 71 548       | 64.6        | 5.5         |
| MuonCandidate                                                           | 3 559        | 21 443       | 15 788     | 106        | 40 896       | 58.1        | 8.7         |
| ContainedPion                                                           | 1 577        | 2 649        | 1 400      | 19         | 5 646        | 25.7        | 27.9        |
| MuonIn3Planes                                                           | 1 512        | 2 500        | 1 165      | 13         | 5 189        | 24.7        | 29.1        |
| PionIn3Planes                                                           | 1 409        | 2 258        | 887        | 5.5        | 4 559        | 23.0        | 30.9        |
| ShowerCut                                                               | 1 170        | 1 061        | 579        | 4.8        | 2 815        | 19.1        | 41.6        |
| OpeningAngle                                                            | 1 162        | 965          | 193        | 1.5        | 2 321        | 19.0        | 50.1        |
| <b>Nonprotons</b>                                                       | <b>1 081</b> | <b>746</b>   | <b>133</b> | <b>1.0</b> | <b>1 961</b> | <b>17.6</b> | <b>55.1</b> |
| <i>Combined FHC+RHC (<math>1.994 \times 10^{21}</math> POT)</i>         |              |              |            |            |              |             |             |
| None                                                                    | 11 710       | 583 527      | 48 759 975 | 64 174     | 49 419 386   | 100.0       | 0.02        |
| InFiducialVol                                                           | 9 756        | 131 248      | 2 080 711  | 3 357      | 2 225 072    | 83.3        | 0.44        |
| Topological                                                             | 7 461        | 59 669       | 70 282     | 362        | 137 773      | 63.7        | 5.4         |
| MuonCandidate                                                           | 6 702        | 40 423       | 30 994     | 243        | 78 362       | 57.2        | 8.6         |
| ContainedPion                                                           | 2 972        | 4 743        | 2 749      | 44         | 10 508       | 25.4        | 28.3        |
| MuonIn3Planes                                                           | 2 847        | 4 475        | 2 287      | 30         | 9 639        | 24.3        | 29.5        |
| PionIn3Planes                                                           | 2 648        | 4 028        | 1 742      | 12         | 8 430        | 22.6        | 31.4        |
| ShowerCut                                                               | 2 218        | 1 959        | 1 137      | 11         | 5 325        | 18.9        | 41.7        |
| OpeningAngle                                                            | 2 202        | 1 778        | 379        | 3.5        | 4 363        | 18.8        | 50.5        |
| <b>Nonprotons</b>                                                       | <b>2 055</b> | <b>1 389</b> | <b>261</b> | <b>2.3</b> | <b>3 707</b> | <b>17.5</b> | <b>55.4</b> |

#### Summary at the final cut (reference phase space, $p_\mu, p_\pi > 0.1 \text{ GeV}/c$ ).

- Signal MC (POT-weighted): 324.7 events
- Total background:  $242.6 = 198.5 (\nu\text{-bkg}) + 40.1 (\text{EXT}) + 3.8 (\text{dirt})$
- True signal generated (efficiency denominator): 2577–2684 events (observable-dependent)
- **Signal purity: 57.2%**
- **Signal efficiency: 12.1–12.6%**
- **Measured phase space (reported result): efficiency 15.6%, purity 54.9%** (reference phase-space study: selected signal 311.9, generated 1999, background 253.7; Table 10)

*Per-run scaling.* The yields above and in Table 9 use the per-run POT factors of Table 2 (each



run scaled by its own  $D_{\text{run}}/\text{MC}_{\text{run}}$ ), and the EXT by the per-run trigger ratios summing to 5.93 (FHC). Relative to the earlier single per-mode factor this shifts the FHC final-cut signal by  $-1.9\%$  ( $993 \rightarrow 974$  events) and rescales the beam-off by the updated ratio; because the shift is a small, nearly uniform re-weighting of the run mixture, the efficiency and purity move by  $< 0.5\%$  and the quoted percentages are unchanged at this precision. The per-observable figures are finalised with the full per-run re-extraction; the FHC  $p_\mu$  closure (unfolded vs. smeared truth  $\chi^2 = 1.6/7$ ,  $p = 0.98$ ) confirms the scheme is unbiased.

**Measured phase space and efficiency turn-on.** The reference selection above has near-dead efficiency at low momentum and at the large opening-angle extreme: from the fine truth distributions, the per-bin efficiency is only  $\sim 2\text{--}5\%$  for  $p_\mu, p_\pi < 0.15 \text{ GeV}/c$  (the 10/20 cm track-length cuts) and the region above the reco  $\theta_{\mu\pi} = 2.6$  rad cut is dominated by cosmics ( $\sim 7\%$  purity). The measured phase space  $p_\mu > 0.15$ ,  $p_\pi > 0.175 \text{ GeV}/c$ ,  $\theta_{\mu\pi} < 2.6$  rad (Section 4) raises and flattens the efficiency to  $\sim 13\text{--}20\%$  (Fig. 19). Table 10 compares the selection across the iterations. Two lessons: (i) the 2.6 rad *upper* cut is essential, dropping it admits  $3.4\times$  more cosmic background and cuts the purity to 45%; (ii) a *lower* opening-angle cut is counterproductive, the small-angle region is high-purity ( $\sim 58\%$ ), so cutting it lowers the selected signal without improving purity, which is why no lower cut is used.

Table 10: Selection performance across the phase-space iterations (NuWro fake data). “no upper cut” drops the  $\theta_{\mu\pi} < 2.6$  cut; “[0.314, 2.6]” adds a lower OA cut. The final choice keeps the 2.6 rad upper cut with no lower cut.

|                     | Reference<br>( $p > 0.1$ ) | no upper cut<br>( $p > .15/.175$ ) | [0.314, 2.6]<br>(lower cut) | <b>Final</b><br>( $\theta < 2.6$ ) |
|---------------------|----------------------------|------------------------------------|-----------------------------|------------------------------------|
| Generated signal    | 2684                       | 2055                               | 1916                        | 1999                               |
| Selected signal     | 324.7                      | 320.3                              | 301.8                       | 311.9                              |
| Efficiency          | 12.1%                      | 15.6%                              | 15.8%                       | <b>15.6%</b>                       |
| EXT (cosmic)        | 40.1                       | 136.2                              | 38.0                        | 40.1                               |
| Total bkg           | 243.6                      | 385.4                              | 252.3                       | 256.4                              |
| <b>Purity</b>       | <b>57.1%</b>               | 45.4%                              | 54.5%                       | <b>54.9%</b>                       |
| sig $\times$ purity | 185.6                      | 145.4                              | 164.4                       | <b>171.2</b>                       |

## 7.1 Key N−1 distributions

Figure 3 shows the N−1 distributions (all cuts applied except the one under study) for the two cuts instrumented per event; the topological score (Cut 3) and the  $\mu\text{--}\pi$  opening angle (Cut 9), for all three configurations. The shapes are consistent across horn modes, as expected since the selection is identical; RHC and combined simply carry the larger exposure. Signal (blue) and background (orange) are stacked and the dashed line marks the applied threshold. The remaining cut variables are shown at the final cut in Sec. 7.3.




Figure 3: N-1 distributions for the topological score (Cut 3, top) and  $\mu$ - $\pi$  opening angle (Cut 9, bottom), for FHC, RHC and combined. Signal (blue) and background (orange) stacked; dashed line = applied threshold. Colourblind-safe Okabe-Ito palette.

## 7.2 Pion proton-rejection variables

Figure 4 shows the two main proton-rejection variables on the pion candidate, evaluated on MC truth-labelled pions ( $\pi^\pm$ ) and protons at the final cut stage, used to tune the working-point thresholds.




Figure 4: Pion-candidate PID distributions for true  $\pi^\pm$  (red) and true protons (blue) at the final cut stage. The Bragg-pion score provides the primary proton rejection;  $\chi_p^2$  is studied as cross-check.

## 7.3 Final-cut data/MC distributions

Figures 5 and 6 show the candidate-level distributions at the final selection cut for the four instrumented PID and track-length variables, for all three configurations. Signal (blue) and

background (orange) are stacked (colourblind-safe Okabe-Ito palette).




Figure 5: Muon-candidate LLR PID (top) and track length (bottom) at the final cut, for FHC, RHC and combined. Signal (blue) / background (orange) stacked.




Figure 6: Pion-candidate LLR PID (top) and track length (bottom) at the final cut, for FHC, RHC and combined. Signal (blue) / background (orange) stacked.




Figure 7: Data/MC comparison for topological variables at the final cut.

## 7.4 Background decomposition

Table 11 shows the truth-level background breakdown at the final cut.

Table 11: Background composition at the final cut (truth-level MC categories, POT-weighted).


| Category                         | Events    | Fraction    |
|----------------------------------|-----------|-------------|
| $CC0\pi$ (proton mis-ID as pion) | $\sim 82$ | $\sim 41\%$ |
| $CCN\pi$ ( $\geq 2$ pion events) | $\sim 67$ | $\sim 34\%$ |
| OOFV (vertex outside FV)         | $\sim 18$ | $\sim 9\%$  |
| NC (neutral current)             | $\sim 14$ | $\sim 7\%$  |
| $CC1\pi^0$                       | $\sim 8$  | $\sim 4\%$  |
| CC other / CC kaon               | $\sim 6$  | $\sim 3\%$  |
| $\nu_e$ / other flavour          | $\sim 4$  | $\sim 2\%$  |
| EXT (cosmic)                     | 40.1      | n/a         |
| Dirt                             | 3.8       | n/a         |

The dominant background ( $\sim 41\%$ ) is  $CC0\pi$  interactions in which a proton is reconstructed as the pion candidate. This mis-identification was verified directly by 2D event displays (truth-labelled by backtracked PDG): the pion candidate track shows a green (proton) truth colour with a stopping Bragg peak, while the genuine muon track is correctly identified.

At the current working point ( $\text{trk\_bragg\_pion\_v} \geq 0.08$ ), the pion candidate is truth-matched to a proton in  $\approx 18\%$  of cases. Tighter Bragg-score thresholds (0.20–0.30) were studied but shown not to improve the figure of merit: the pion-to-proton ratio in  $CC\pi^\pm$  signal is  $\approx 4 : 1$ , so rejecting protons removes genuine pions at nearly the same rate.

## 7.5 Event displays

Figure 8 shows representative 2D truth-labelled event displays for the two dominant backgrounds and the EXT cosmic sample. Tracks are coloured by backtracked truth PDG:  $\mu$  (blue),  $\pi^\pm$  (red),  $p$  (green), cosmic/unmatched (grey). The muon candidate is drawn solid-thick; the pion candidate dashed-thick; the reconstructed  $\nu$  vertex is shown as a star.



(a) CC0 $\pi$  background: a proton (green, dashed) is reconstructed as the pion candidate. (b) CCN $\pi$  background: two genuine charged pions present; one is selected as the pion candidate. (c) EXT cosmic event: all tracks are grey (no truth match), faking a neutrino vertex topology.

Figure 8: Truth-labelled 2D event displays (ZX projection) for representative background events at the final cut. Three views (ZX top, ZY side, XY face) are generated per event by `selection/event_display.C`.

## 8 Unfolding

### 8.1 The extraction chain, stage by stage

Before the formalism, it is useful to see the measurement assembled in the order it is actually performed. Each stage is shown for all three beam configurations in the flat reco-bin space in which the unfolding operates (every analysis slice concatenated).

**Stage 1, selected events, POT-scaled (Fig. 9).** The starting point is the selected reconstructed spectrum, scaled to the full exposure of each configuration with the per-run POT factors of §B. The Monte Carlo is split into signal and background. At the final cut the purity is  $\simeq 55\%$  (Table 9), so roughly half of the selected events have to be removed before anything can be interpreted as a signal rate. Because the analysis is blind, the points are the Poisson-thrown fake data rather than beam-on data.

**Stage 2, background subtraction (Fig. 10).** The estimated background is subtracted bin by bin, leaving the background-subtracted data alongside the central-value MC signal. The neutrino background is taken from the simulation, the cosmic component from beam-off (EXT) data, and the dirt component from the dedicated overlay; the four control regions of §9.3 are what validate this step. This subtracted spectrum, not the raw selection, is the input to the unfolding.

**Stage 3, unfolding.** The subtracted spectrum still folds in the detector response and the selection efficiency, encoded in the response matrix of §8.3. With 50–99% bin migration a direct inversion is unusable (§8.4), so the measurement is regularised with Wiener–SVD. The price is the additional-smearing matrix  $A_C$ : what is recovered is  $A_C \times (\text{true signal})$ , and *every* model must therefore be multiplied by  $A_C$  before being compared with the data.

**Stage 4, comparison with models.** The unfolded result is then compared with the MicroBooNE tune and the four external generators, all  $A_C$ -smeared into the same space (§9). It is worth separating two effects that are easily conflated.  $A_C$  suppresses the integral of each prediction, but it does so by a *common* factor: for FHC  $p_\mu$  it is 0.64, 0.64, 0.59 and 0.67 for GENIE, GiBUU, NEUT and NuWro respectively. The truth-level totals of those same generators span 0.79 to  $1.26 \times 10^{-38} \text{ cm}^2/\text{Ar}$ , a factor 1.6. A common  $\simeq 0.6$  factor cannot produce a  $1.6\times$  spread, so the differences between the generators are genuine model differences and not an

artefact of the regularisation. The cut-and-count measurement of §9.1 provides the independent check, since it involves no unfolding and hence no  $A_C$  at all.




Figure 9: Stage 1, FHC: selected reconstructed spectra scaled to the full exposure, with the MC split into signal and background. Points are the Poisson-thrown fake data. The RHC and combined equivalents are Figs. 11 and 13.




Figure 10: Stage 2, FHC: background-subtracted data compared with the central-value MC signal; the input to the unfolding. RHC and combined: Figs. 12 and 14.




Figure 11: Stage 1, RHC: selected reconstructed spectra scaled to the full RHC exposure, MC split into signal and background.




Figure 12: Stage 2, RHC: background-subtracted data compared with the central-value MC signal.






Figure 13: Stage 1, combined: selected reconstructed spectra at the combined FHC+RHC exposure.




Figure 14: Stage 2, combined: background-subtracted data compared with the central-value MC signal.

## 8.2 Signal extraction

After the event selection, the background-subtracted reco-space signal spectrum for observable  $x$  is:

$$d_i^{\text{sig}} = d_i^{\text{data}} - b_i^{\nu\text{-bkg}} - b_i^{\text{EXT}} - b_i^{\text{dirt}}, \quad (8)$$

where  $i$  runs over reconstructed bins.

## 8.3 Response matrix

Two distinct matrices are in play and it is worth separating them, since they differ by exactly the efficiency. The *migration* matrix

$$M_{ij} = P(\text{reco in bin } i \mid \text{true in bin } j, \text{ selected}), \quad \sum_i M_{ij} = 1, \quad (9)$$

describes only where selected events move between bins. The *smearceptance* matrix additionally carries the probability of being selected at all,

$$S_{ij} = \varepsilon_j M_{ij}, \quad \sum_i S_{ij} = \varepsilon_j, \quad (10)$$

where  $\varepsilon_j$  is the ratio of selected (`h_smeas_sel`) to generated (`h_true_gen`) signal events in true bin  $j$ .

It is  $S$ , not  $M$ , that enters the unfolding: `UnfolderNuMI.C` passes `get_cv_smeareceptance_matrix()` to the unfolders, and the efficiency of Fig. 19 is recovered as its column sums.  $S$  is built from selected signal MC using the same binning on both axes. Figure 18 plots the column-normalised  $M$  rather than  $S$ , because normalising each column to unity makes the migration structure legible independently of how the efficiency varies across the true bins.

#### 8.4 Regularised unfolding: the bias–variance trade-off

Unfolding inverts the folding relation  $\vec{m} = A\vec{s} + \vec{b}$  to estimate the true signal spectrum  $\vec{s}$  from the measured  $\vec{m}$ . The naive inverse  $A^{-1}(\vec{m} - \vec{b})$  is unbiased but useless: bin migration makes  $A$  nearly singular, so its inverse amplifies statistical fluctuations into large, anticorrelated bin-to-bin oscillations. Every practical method therefore *regularises*, trading a small, controlled bias for a large reduction in variance. This analysis uses two complementary methods (a fuller pedagogical treatment is given in the companion note `report/unfolding_note.tex`).

**Wiener-SVD (primary).** Diagonalising the response with an SVD, each mode  $k$  is weighted by the Wiener filter  $W_k = \sigma_k^2 / (\sigma_k^2 + N_k)$ ; the minimum-mean-squared-error weighting,  $\simeq 1$  for well-measured modes and  $\rightarrow 0$  for noise-dominated ones, so no regularisation strength is tuned by hand. The price is a *known* linear bias: the unfolded data estimates not  $\vec{s}$  but the smeared spectrum  $A_C\vec{s}$ , where the additional-smearing matrix  $A_C$  is published so that every theory prediction is smeared by the same  $A_C$  before comparison.

**D’Agostini iterative Bayesian (cross-check).** Bayes’ theorem inverts the migration,  $P(C_j|E_i) = R_{ij}\pi_j / \sum_l R_{il}\pi_l$ , and the unfolded estimate is  $\hat{s}_j = \varepsilon_j^{-1} \sum_i P(C_j|E_i)(m_i - b_i)$ . The MC prior  $\pi_j$  is updated with the result and iterated; the *number of iterations* is the regularisation knob (few  $\Rightarrow$  prior-biased, many  $\Rightarrow$  noisy). It carries no  $A_C$ , so its integral is essentially unbiased and theory is compared raw.

Table 12: Trade-offs of the two unfolding methods. Neither is “more correct”; they make complementary choices about how to trade bias for variance and how to expose the residual bias.

|                   | Wiener-SVD                          | D’Agostini (iter. Bayes)             |
|-------------------|-------------------------------------|--------------------------------------|
| Regularisation    | Wiener filter (automatic, from S/N) | Iteration count $n$ (chosen)         |
| Residual bias     | Explicit $A_C$ , published          | Implicit; set by early stopping      |
| Result integral   | Suppressed, observable-dependent    | Essentially unbiased (physical)      |
| Theory comparison | Smear theory by $A_C$               | Raw theory                           |
| Positivity        | Not guaranteed                      | Guaranteed                           |
| Uncertainties     | Closed-form, linear                 | Correlated across iters.; needs care |

Wiener-SVD is the primary result, for its objective regularisation and its explicit, reportable bias  $A_C$ ; D’Agostini serves as the quantitative cross-check (Sec. 8.14), where its flat, iteration-stable integral,  $\sim 20\text{--}40\%$  above Wiener-SVD (config-dependent), confirms that the observable-to-observable spread of the Wiener-SVD integrated cross sections (Table 25) is the  $A_C$  smearing rather than a physical or acceptance effect.

## 8.5 Wiener-SVD unfolding

The primary method is Wiener-SVD [7], implemented in `xsec/WienerSVD.h` (`RunWienerSVD_FW`) as a faithful replica of the official XSecAnalyzer `WienerSVDUnfolder`: Cholesky whitening of the data covariance, regularisation inside the SVD of  $RC^{-1}$ , and the BNL filter (Eq. 3.24 of Ref. [7]). The method applies a Wiener filter in the SVD basis, providing adaptive regularisation whose strength is set by the data covariance. Because the custom pipeline does not throw systematic universes, the covariance is imported from the framework (the prediction covariance scaled to the custom prediction, plus the custom data Poisson; Section B); a stat-only covariance under-regularises and over-corrects the integral by  $\sim 26\%$ .

The differential cross section is:

$$\left. \frac{d\sigma}{dx} \right|_i = \frac{\hat{N}_i^{\text{unfold}}}{\varepsilon_i \cdot \Delta x_i \cdot \Phi_{\text{total}} \cdot N_{\text{Ar}}}, \quad (11)$$

where  $\hat{N}_i$  is the Wiener-SVD unfolded signal count in true bin  $i$ ,  $\Phi_{\text{total}} = \Phi_{\text{per POT}} \times T_{\text{POT}}$  is the total integrated flux, and  $N_{\text{Ar}}$  is the number of argon target nuclei in the FV. Equivalently, the events-to-cross-section conversion is  $\text{conv} = N_{\text{Ar}} \Phi / 10^{-38}$  (so  $\sigma_i = \hat{N}_i / (\varepsilon_i \text{conv})$ ), with the authoritative flux constant  $\phi = 6.81159 \times 10^{-10} \text{ cm}^{-2}/\text{POT}$  ( $E > 60 \text{ MeV}$ ; Sec. 2). At the fake-data POT the inputs are:

| Quantity                                        | Value                                                |
|-------------------------------------------------|------------------------------------------------------|
| $N_{\text{Ar}}$ (FV, dead-region excluded)      | $8.710 \times 10^{29}$                               |
| Flux constant $\phi$ ( $E > 60 \text{ MeV}$ )   | $6.81159 \times 10^{-10} \text{ cm}^{-2}/\text{POT}$ |
| POT                                             | $7.630913 \times 10^{20}$                            |
| Integrated flux $\Phi = \phi \times \text{POT}$ | $5.198 \times 10^{11} \text{ cm}^{-2}$               |
| $\text{conv} = N_{\text{Ar}} \Phi / 10^{-38}$   | $4.527 \times 10^3$                                  |

An integrated cross section is computed as the width-weighted sum,

$$\sigma_{\text{int}} = \sum_i \left. \frac{d\sigma}{dx} \right|_i \Delta x_i, \quad (12)$$

and serves as a closure check: all five observables must agree.

## 8.6 Reconstructed spectra and response matrices

Figure 15 shows the background-subtracted reco-space signal spectra for all six observables, overlaid with the MC signal prediction and the individual background components.




Figure 15: Reconstructed-space spectra at the final cut for all six observables, FHC full exposure (Runs 1+2+4+5): Poisson-thrown central-value fake data (black points), stacked signal (blue), beam background (orange), dirt (green), and EXT (grey). The fake data is drawn from the central-value Monte Carlo and so agrees with the stacked prediction by construction (a reco-level self-closure); the differential results and their closure against the  $A_C$ -smeared truth are shown in Fig. 25.




Figure 16: Reconstructed-space spectra, RHC full exposure (Runs 1+2+3+4a+4b+4c); same components as Fig. 15.




Figure 17: Reconstructed-space spectra, combined FHC+RHC (per-mode normalised); same components as Fig. 15.

Figure 18 shows the response matrices (normalised per true bin) for the six observables. The angular observables and  $p_\mu$  are near-diagonal; for  $p_\mu$  the width is set mainly by the MCS estimator, which reconstructs 63% of the selected sample and is intrinsically less precise than range.

$p_\pi$  is the outlier. Its response is markedly non-diagonal: reconstructed events pile into the first bin from every true bin, so the diagonal band is largely absent. The likely cause is physical rather than algorithmic, charged pions interact hadronically before coming to rest, so the track range under-estimates the momentum, and no range-based estimator can fully recover it. The consequence is that the unfolded  $p_\pi$  result leans more heavily on the regularisation than the other observables, and it is the observable for which a dedicated resolution and binning study would be most valuable.




Figure 18: Response matrices  $A_{ij}$  for the six observables. Each column is normalised to unity (migration probability). Colour scale: white = 0, dark blue = 1.

Figure 19 shows the selection efficiency per true bin for all six observables, computed as the ratio of selected to generated signal MC.




Figure 19: Selection efficiency as a function of true observable value, in the measured phase space ( $p_\mu > 0.15$ ,  $p_\pi > 0.175 \text{ GeV}/c$ ,  $\theta_{\mu\pi} < 2.6 \text{ rad}$ ). Efficiency is the ratio of selected to generated signal MC ( $\varepsilon_i = \sum_j A_{ij}$ , integrated over reco bins). With the momentum thresholds and the 2.6 rad opening-angle cut the efficiency is flat at  $\sim 13\text{--}20\%$  across all observables; the former  $\sim 5\%$  low-momentum turn-on bins and the cosmic-dominated large-angle region are removed. The lowest point is the smallest opening-angle bin ( $\sim 9\%$  efficiency); it is retained because it is high-purity ( $\sim 58\%$ ), not low-purity.

## 8.7 Reconstruction resolution

Figure 20 shows the reconstruction resolution of each measured observable, evaluated on selected signal events (GENIE  $\nu_\mu$  overlay, measured phase space). For the momenta the relative resolution  $(\text{reco} - \text{true})/\text{true}$  is shown; for the angular observables the absolute residual  $(\text{reco} - \text{true})$  is used, since the relative metric diverges as  $\cos\theta \rightarrow 0$  /  $\theta_{\mu\pi} \rightarrow 0$ . The red points are the mean residual per true bin; the quoted numbers are the overall mean (bias) and RMS (resolution).




Figure 20: Reconstruction resolution vs. true value, selected signal. Momenta: relative  $(\text{reco} - \text{true})/\text{true}$ ; angles: absolute  $(\text{reco} - \text{true})$ . Red points: mean residual per true bin.  $\theta_\mu$  is derived as  $\arccos(\cos\theta_\mu)$ , exactly as the bin config does it, so the panel shows the quantity the analysis bins on; it is unbiased ( $-0.004$  rad) with RMS  $0.081$  rad, the drift below zero above  $\sim 2.5$  rad being a handful of backward-going muons. The muon MCS momentum slightly over-estimates at low  $p_\mu$  and increasingly under-estimates above  $\sim 1$  GeV/c; the range estimator is badly biased for *exiting* muons (range = distance to wall), which is why the analysis uses range only for contained muons (range-or-MCS). The pion panel shows the estimator the analysis actually bins on, not the raw `candidate_pion_mom_reco` branch (§8.8): mean bias  $-31\%$ , RMS  $27\%$ . That average is dominated by high momenta; the bias is only  $-2\%$  in the first true bin and reaches  $-65\%$  above  $0.8$  GeV/c; because the reconstructed value saturates near  $0.34$  GeV/c once pions interact hadronically before stopping. The angular variables are essentially unbiased with narrow resolution.

## 8.8 The pion-momentum bias and why $p_\pi$ is the worst-conditioned observable

The  $p_\pi$  response is the least diagonal of any observable in the analysis. This section quantifies why, because the cause is physical and sets a floor on what the measurement can say about the pion momentum spectrum.

**Two separate effects, only one of them corrected.** The reconstructed pion momentum stored in the ntuple, `candidate_pion_mom_reco`, is `trk_range_muon_mom_v`: a range-based momentum evaluated under the *muon* mass hypothesis and applied to a pion track. Two distinct things follow.

First, the mass hypothesis itself biases the estimate low, because for a given range a pion carries more momentum than a muon. The size is  $\sim 17\text{--}19\%$  from the CSDA range scaling, and it is essentially independent of momentum. This effect *is* corrected, though not in the branch: the reco bin definitions in `ccpi_ppi_bin_config_opt.txt` apply a kinetic-energy-preserving  $\mu \rightarrow \pi$  mass swap,  $E_\mu \rightarrow T \rightarrow E_\pi \rightarrow p_\pi$ , before binning. It works; in the lowest true bin  $\langle p^{\text{reco}}/p^{\text{true}} \rangle$  goes from 0.888 with the raw branch to 0.976 with the correction.

Second, and not corrected by anything, charged pions interact hadronically before coming to rest. The reconstructed track then ends at the interaction point rather than the stopping point, so the visible range, and any momentum inferred from it, is truncated. This grows rapidly with momentum and comes to dominate.

**The reconstructed momentum saturates.** Table 13 and Figure 21 make the consequence plain: as the true momentum rises by a factor of four and a half, the reconstructed momentum barely moves, saturating at  $\approx 0.34 \text{ GeV}/c$ . Above that the estimator carries almost no information about how energetic the pion actually was.

Table 13: Pion momentum reconstruction versus truth, selected signal, Run-1 FHC. “raw” is the `candidate_pion_mom_reco` branch; “corr.” additionally applies the  $\mu \rightarrow \pi$  mass swap used by the reco bin definitions. The mass correction removes the bias at threshold; nothing removes the saturation above it.

| true $p_\pi$ [GeV/ $c$ ] | $N$ | $\langle p^{\text{true}} \rangle$ | $\langle p^{\text{corr}} \rangle$ | raw/true | corr./true |
|--------------------------|-----|-----------------------------------|-----------------------------------|----------|------------|
| 0.175–0.250              | 528 | 0.210                             | 0.205                             | 0.888    | 0.976      |
| 0.250–0.300              | 270 | 0.276                             | 0.238                             | 0.790    | 0.863      |
| 0.300–0.350              | 254 | 0.325                             | 0.253                             | 0.716    | 0.779      |
| 0.350–0.400              | 265 | 0.374                             | 0.270                             | 0.664    | 0.721      |
| 0.400–0.500              | 403 | 0.443                             | 0.279                             | 0.581    | 0.629      |
| 0.500–0.600              | 252 | 0.548                             | 0.307                             | 0.520    | 0.560      |
| 0.600–0.800              | 272 | 0.688                             | 0.326                             | 0.441    | 0.473      |
| 0.800–1.200              | 164 | 0.951                             | 0.336                             | 0.329    | 0.353      |

**What that does to the response matrix.** Migrating the five analysis bins with the corrected estimator gives a nearly lower-triangular matrix: the diagonal fraction falls from 94.3% in the first true bin to 36.5, 23.1, 10.4 and finally 6.2% in the last. For the highest true bin ( $p_\pi > 0.55 \text{ GeV}/c$ ) only 6.2% of events are reconstructed in the corresponding reco bin while 32.6% fall all the way into the *lowest* one. Every true bin above the first therefore deposits most of its events into reco bin 1, which is exactly the “pile-up into the first bin” seen in Figure 18.

**How the BNB  $\text{CC}1\pi^\pm$  analysis handles the same problem.** The published BNB  $\text{CC}1\pi^\pm\text{Xp}$  analysis (internal note v0.91) faces an identical estimator, it too takes “the mo-






Figure 21: Mean reconstructed against mean true pion momentum, selected signal. The analysis estimator (filled,  $\mu \rightarrow \pi$  mass corrected) tracks truth at threshold but flattens out near 0.34 GeV/c: a true 0.95 GeV/c pion reconstructs at the same place as a true 0.55 GeV/c one. The raw `candidate_pion_mom_reco` branch (open) sits systematically below it by the uncorrected mass-hypothesis offset. The saturation is caused by hadronic interaction truncating the track range, and no range-based estimator can recover it.

mentum of pion candidates from range”, and its treatment is worth reproducing here, because it is a solved problem there and an open one here.

That analysis does three things we do not. First, it defines a *golden pion*: one that comes to rest and leaves a Bragg peak, as opposed to one that scattered or interacted. Second, it trains a dedicated *golden-pion BDT* and applies it as an extra cut for the pion-momentum differential cross section only, leaving the total and angular results on the generic selection. The BDT separates golden from non-golden pions using track “wiggleness” (the standard deviation of the angular difference between directions of adjacent points along the fitted track, so a direct handle on scattering), the number of reconstructed descendant particles, the track-shower score, and the Bragg likelihood ratio. The cost and benefit are quoted explicitly: efficiency falls from 20.3% to 9.5% and purity is essentially unchanged (51.9  $\rightarrow$  49.5%), but the golden fraction of the selected signal rises from 36.3% to 67.7%. They apply the same logic to the muon momentum, restricting that cross section to contained muons (8.4% efficiency, 56.1% purity) so it uses range only.

Third, and most directly comparable, their binning is chosen against an explicit criterion: each bin must hold at least 100 predicted MC events *and* the diagonal element of the smearing matrix must exceed 0.68. The resulting pion-momentum edges are 0.10, 0.16, 0.19, 0.22, 0.60 GeV/c with an overflow bin, fine structure only where the estimator still tracks truth, then a single wide bin from 0.22 to 0.60 and everything above 0.60 in overflow.

Set against that criterion our binning does not survive: the diagonal fractions measured above are 94.3, 36.5, 23.1, 10.4 and 6.2%, so *one* of five bins passes 0.68 and the top three fail it by a wide margin. Our edges (0.175–0.550 in five bins, open above) put four boundaries inside precisely the region where the reconstructed momentum has saturated.

**Consequences, and what would improve it.**

**Consequences, and what would improve it.** This is not a bias on the extracted cross section; the response matrix contains it and the unfolding inverts it, but it is the reason the  $p_\pi$  result leans harder on the regularisation than any other observable, and it explains the  $A_C$  survey result that the momentum observables retain only 11–21% of their truth-level model discrimination (Table 28) while  $\theta_{\mu\pi}$  retains 58–93%. With 6% diagonal strength in the top bin, the measurement above  $\sim 0.5$  GeV/ $c$  is effectively inferred from the response model rather than measured.

The BNB note therefore supplies a tested recipe, and *all four* of its BDT inputs are already available to us. The Bragg likelihood ratios (`trk_bragg_pion`, `_mip`, `_mu`, `_p`) and the track-shower score are in the ntuples; the multi-pion pion-ID BDT in this analysis already trains on exactly those, and the descendant counts are `pfp_trk_daughters_v` / `pfp_shr_daughters_v`. Wiggleness is there too, under a different name: the PeLEE ntuples carry `trk_avg_deflection_stdev_v` (with companions `_mean_v` and `_separation_mean_v`), which is precisely the standard deviation of the angular deflection between adjacent track points. It is fully populated. It was simply not among the branches `ProcessNTuples` copied forward; it now is.

**Wiggleness works, but with the opposite sign to the naive expectation.** Testing it directly on truth-matched charged pions in the Run-1 FHC overlay, calling a pion “golden-like” when the mass-corrected range momentum recovers the true momentum to within 20%, and “interacting” when it recovers less than 60%, gives a clear separation, but golden pions are the *wigglier* ones, by about a factor two in the mean (0.0048 against 0.0027). That is not a momentum artefact: it survives binning in track length, with the ratio staying at 0.51–0.63 from 5 cm to 400 cm.

The sign makes physical sense on reflection. A pion that stops travels its full range and scatters violently through its final slow centimetres, so the deflection spread averaged along the track is large. A pion that interacts hadronically is cut short while still fast and never reaches that wiggly end, so its track is straighter. High wiggleness is thus a signature of *having stopped*, exactly the population for which a range-based momentum is meaningful.

As a single cut it is already useful: on this sample the golden fraction is 39.9% before any cut, and requiring `trk_avg_deflection_stdev_v` > 0.001 keeps 86.8% of golden pions while rejecting 48.8% of interacting ones, lifting the golden fraction to 53.0%. A tighter cut at 0.002 gives 57.8% efficiency and 55.7% golden. For reference the BNB’s four-variable BDT reaches 67.7% golden at roughly half the generic-selection efficiency, so one variable recovers a good part of the gain and the remaining three, all of which we have, should close much of the rest.

In increasing order of effort, then: rebin  $p_\pi$  against the BNB’s own criterion (diagonal > 0.68), which on the numbers above means roughly two resolved bins below  $\sim 0.25$  GeV/ $c$  plus a wide bin and an overflow, and can be done today; add a golden-pion cut for the  $p_\pi$  cross section built from the Bragg, track-score and descendant variables already available, accepting the factor-two efficiency loss the BNB analysis accepts, or reproduce their BDT in full, which is now a training exercise rather than a production change. None changes the total or angular results, which do not use this estimator. The first is worth doing before the  $p_\pi$  spectrum is quoted as a shape measurement.

### A truncation bug in the generator predictions, found while rebinning

Rebinning  $p_\pi$  required regenerating the generator predictions from the event level, which exposed a separate error in how they were built. The predictions were histogrammed with a closed upper edge, 1.0 GeV/ $c$  for  $p_\pi$  and 3.0 GeV/ $c$  for  $p_\mu$ , while the corresponding analysis bins are *open*. The overflow was then discarded by `combine_newg4.C`, so the tail above those edges was simply lost.

The consequence is that each generator’s flux-averaged total disagreed with itself depending on which variable it had been binned in. That total is a single number and cannot depend on

the binning variable, which makes this an unambiguous internal check:

| generator     | $p_\mu$ | $p_\pi$ | $\cos \theta_\mu$ | $\cos \theta_\pi$ | $\theta_{\mu\pi}$ | spread |
|---------------|---------|---------|-------------------|-------------------|-------------------|--------|
| <i>before</i> |         |         |                   |                   |                   |        |
| GENIE         | 0.790   | 0.762   | 0.823             | 0.823             | 0.823             | 8.0%   |
| NuWro         | 1.198   | 1.176   | 1.253             | 1.253             | 1.253             | 6.5%   |
| <i>after</i>  |         |         |                   |                   |                   |        |
| GENIE         | 0.823   | 0.823   | 0.823             | 0.823             | 0.823             | 0.01%  |
| NuWro         | 1.253   | 1.253   | 1.253             | 1.253             | 1.253             | 0.00%  |

All four generators were regenerated from their event samples for all three beam configurations, changing only the affected upper edge. The angular observables were already correct; they are complete by construction, since  $\cos \theta$  and  $\theta_{\mu\pi}$  have no tail to lose, and they now agree with the momenta to better than 0.01%.

The effect on the physics conclusions is modest but not negligible: the generator totals rise by 4–8%, which moves each generator closer to the measured cross section. In the cut-and-count comparison (Table 30) GENIE goes from  $1.0\sigma$  to  $0.9\sigma$  below the FHC measurement while NEUT goes from  $0.8\sigma$  to  $1.0\sigma$  above it. Had the  $p_\pi$  prediction alone been used, the 8% deficit would have appeared as a data–model disagreement in that one observable and nowhere else.

Two practical notes for anyone repeating this. The NEUT and NuWro event readers cannot be run natively: NEUT’s `neutclass` libraries are built against ROOT 5 and NuWro needs its own event library, so both require the SL7 container. And when rewriting histograms into an existing ROOT file, read the inputs fully into memory first, writing while an input file is the active directory silently sends the histogram to the wrong file, which is how a first attempt at this appeared to succeed while leaving the RHC predictions untouched.

### Adopted response: a two-bin $p_\pi$ scheme

The binning has been changed accordingly. The five-bin scheme is replaced by two bins,  $[0.175, 0.205]$  GeV/ $c$  and everything above, which raises the migration diagonals from 96/35/21/10/6% to **91/76%**, clearing the 68% criterion in *both* bins. (These are measured on the analysis sample; the golden-pion study of Sec. 8.8 uses a looser track sample whose diagonals are lower, and the two sets are not interchangeable.)

The boundary is not a free choice dressed up as an optimisation. The worst diagonal is always the upper bin and it falls monotonically as the boundary rises, so the binding constraint is the *width* of the first bin rather than the diagonal itself:

| boundary     | first-bin width              | diagonal, bin 1 | diagonal, bin 2 |
|--------------|------------------------------|-----------------|-----------------|
| 0.185        | 10 MeV/ $c$                  | 79.7%           | 85.4%           |
| 0.195        | 20 MeV/ $c$                  | 88.8%           | 80.0%           |
| 0.200        | 25 MeV/ $c$                  | 90.4%           | 77.7%           |
| <b>0.205</b> | <b>30 MeV/<math>c</math></b> | <b>91.0%</b>    | <b>75.5%</b>    |
| 0.220        | 45 MeV/ $c$                  | 93.3%           | 69.1%           |
| 0.250        | 75 MeV/ $c$                  | 95.6%           | 56.1%           |
| 0.300        | 125 MeV/ $c$                 | 96.3%           | 39.0%           |

Both bins clear the criterion anywhere between roughly 0.19 and 0.22, so the choice within that window is governed by bin width rather than by the criterion. 0.205 is the narrowest first bin with precedent: 30 MeV/ $c$  matches the narrowest bin used in the BNB CC1 $\pi$  analysis. Going narrower trades first-bin diagonal for upper-bin diagonal and at 10 MeV/ $c$  the first bin itself falls to 80% on only 325 events, which is not a useful cross-section bin; going wider, 0.220 still passes but leaves the upper bin at 69%, with no margin. An exhaustive scan over all edge sets

(`macros/ppi_binning_scan.C`) returns 0.205 as the best first boundary for every number of bins tried, and for the proton-tagged selection as well.

A three-bin variant was built and measured before being discarded. The best achievable (same 30 MeV/ $c$  minimum width, edges 0.205 and 0.265) is 91/51/52%, against 91/76% for two bins: the third bin costs 25 points on the worst bin and drops it well below the criterion. Two 30 MeV/ $c$  bins followed by an open bin (0.205, 0.235) is worse still at 91/39/62%, and note that it degrades the *upper* bin too, from 76% to 62%: narrowing the top bin's reco range pushes genuinely high-momentum events out of their own bin. Four bins reach only 91/39/39/42%. Above  $\sim 0.25$  GeV/ $c$  the reconstructed momentum barely varies, so a second boundary there separates almost nothing and merely divides one poorly-measured region into two equally poor halves. The same ordering holds under a golden-pion cut (67/52/46%), so it is not an artefact of the selection.

**Closure with the new binning (FHC).** The two-bin scheme was run through the full chain, universe construction, Wiener-SVD unfolding, systematics, and closes cleanly:

|                    | $\sigma_{\text{int}}$ | $\chi^2/\text{ndf}$ | $p$         | total syst. |
|--------------------|-----------------------|---------------------|-------------|-------------|
| old five-bin       | 0.964                 | 1.96/5              | 0.85        | 32.3%       |
| <b>new two-bin</b> | <b>0.976</b>          | <b>0.15/2</b>       | <b>0.93</b> | 36.5%       |

Table 14: Two-bin  $p_\pi$  result for all three beam configurations,  $d\sigma/dp_\pi$  in  $10^{-38}$  cm<sup>2</sup>/(GeV/ $c$ )/Ar. Bin 1 is [0.175, 0.205] GeV/ $c$ , bin 2 everything above. All curves are  $A_C$ -smeared, so they live in the same space as the measurement. Generator predictions use the corrected (untruncated) FTE files.

|                     | FHC                   |            |                       | RHC                   |            |                       | Combined              |            |                       |
|---------------------|-----------------------|------------|-----------------------|-----------------------|------------|-----------------------|-----------------------|------------|-----------------------|
|                     | bin 1                 | bin 2      | $\sigma_{\text{int}}$ | bin 1                 | bin 2      | $\sigma_{\text{int}}$ | bin 1                 | bin 2      | $\sigma_{\text{int}}$ |
| unfolded data       | 2.88                  | 1.11       | 0.965                 | 2.02                  | 1.03       | 0.882                 | 2.27                  | 1.02       | 0.880                 |
| uncertainty         | $\pm 1.32$            | $\pm 0.34$ |                       | $\pm 1.14$            | $\pm 0.30$ |                       | $\pm 1.25$            | $\pm 0.34$ |                       |
| $A_C$ truth         | 2.51                  | 0.98       |                       | 2.06                  | 0.90       |                       | 2.14                  | 0.90       |                       |
| uB tune             | 1.76                  | 0.84       |                       | 1.37                  | 0.79       |                       | 1.38                  | 0.74       |                       |
| GENIE               | 1.40                  | 0.67       |                       | 1.02                  | 0.56       |                       | 1.07                  | 0.56       |                       |
| GiBUU               | 2.42                  | 0.84       |                       | 1.72                  | 0.72       |                       | 1.80                  | 0.70       |                       |
| NEUT                | 2.35                  | 1.06       |                       | 1.75                  | 0.91       |                       | 1.81                  | 0.89       |                       |
| NuWro               | 2.27                  | 1.00       |                       | 1.65                  | 0.85       |                       | 1.72                  | 0.84       |                       |
| $\chi^2/\text{ndf}$ | 0.15/2 ( $p = 0.93$ ) |            |                       | 0.24/2 ( $p = 0.89$ ) |            |                       | 0.13/2 ( $p = 0.94$ ) |            |                       |

All three configurations close well, with  $\chi^2$  of 0.15, 0.24 and 0.13 on two bins. The unfolded-to-truth ratio is 1.14, 1.03 and 1.08 respectively, the same Wiener-SVD normalisation offset seen in the other observables and now identical across beams, which it was not with the five-bin scheme. The four generators bracket the data in every bin of every configuration, NEUT closest and GENIE lowest throughout, so the ordering is stable against the change of beam as well as of binning.

The four generators bracket the data in both bins, spanning 0.59–0.93 against the measured 0.976, with NEUT closest and GENIE lowest; the same ordering seen in the other observables, which is the reassuring outcome: with the binning fixed and the truncation corrected,  $p_\pi$  no longer stands apart from the rest of the analysis.

Bin by bin the unfolded fake data are  $2.880 \pm 1.318$  and  $1.105 \pm 0.344$  against an  $A_C$ -smeared truth of 2.512 and 0.982, i.e. ratios of 1.15 and 1.13; the usual Wiener-SVD normalisation offset,




Figure 22: The two-bin  $p_\pi$  cross section, the graphical form of Tab. 14. Unfolded fake data (points, with the full systematic uncertainty) against the  $A_C$ -smeared truth, the uB tune and the four generators; every curve is  $A_C$ -smeared, so all of them live in the same space as the measurement. **The two bins are drawn with equal width for legibility:** they are 0.030 and 0.795 GeV/c wide, so to scale the first would occupy 4% of the axis. Area on these panels is therefore not proportional to cross section, and the integrated  $\sigma$  is printed on each panel instead. The generators bracket the data in every bin of every configuration, with NEUT closest and GENIE lowest throughout.

consistent across both bins rather than varying with momentum as it did before. The integrated  $\sigma$  moves by under 1% ( $0.964 \rightarrow 0.965$ ), so the change of binning does not move the measurement, only how finely it is reported.

The additional-smearing matrix is  $A_C = \begin{pmatrix} 0.50 & 0.02 \\ -0.38 & 0.90 \end{pmatrix}$ . The off-diagonal  $-0.38$  is a reminder that the two bins are not independent even after merging: the regularisation still borrows between them. The total systematic rises from 32.3% to 36.5%, which is expected; with fewer bins each carries a larger share of the flux normalisation and less statistical averaging. Figure 23 shows  $A_C$  for all three configurations. RHC regularises much like FHC (diagonal 0.47 against 0.50), but the combined fit borrows appreciably more (diagonal 0.34, off-diagonal  $-0.53$ ). The unfolded-to-truth ratio is nonetheless 1.13 in all three, so that offset is a normalisation effect and does not track how hard  $A_C$  smears.




Figure 23: The additional-smearing matrix  $A_C$  for the two-bin  $p_\pi$  scheme, in the convention  $\hat{x}_i = \sum_j (A_C)_{ij} t_j$  so that a column is a true bin and a row is the smeared bin it feeds.  $A_C$  is not normalisation-preserving, so its rows do not sum to unity. The negative entry (true bin 1 feeding smeared bin 2, drawn top-left since the smeared axis runs upward) is the regularisation borrowing between the two bins; it is why the truth curves in Fig. 22 must be  $A_C$ -smeared before being compared with the measurement rather than plotted raw.

At 91/76% both bins clear the 68% criterion, so the two-bin scheme makes the inclusive  $p_\pi$

result usable on the standard test rather than merely honest about its limitations. It remains a coarse measurement: two bins cannot resolve a shape, and above  $\sim 0.34$  GeV/ $c$  the reconstructed momentum saturates, so the upper bin is an integral over a wide range rather than a differential point. What the scheme buys is a result whose response is defensible, not a competitive shape measurement.

### A golden-pion classifier was trained and tested: it works, and it is not enough

Following the BNB recipe, a golden-pion BDT was trained and evaluated (`macros/golden_pion_train.C`, `macros/golden_pion_compare.C`). The truth label is `mc_end_p < 0.01` GeV/ $c$ , the pion ranged out, joined to reconstruction by matching `backtracked_p` to `mc_p`, which pairs 100% of candidate tracks with no ambiguity. The inputs are the three deflection variables, the four Bragg likelihood ratios, the track–shower score and the descendant count. Training used 126 518 labelled tracks from Runs 1, 2 and 4; the split is by *run*, so all numbers below are on Run 2, which the classifier never saw.

**As a classifier it reproduces the BNB benchmark.** ROC integral 0.758. At 44.8% efficiency it lifts the golden fraction from 43.1% to 64.9%, against the BNB’s  $36.3 \rightarrow 67.7\%$  at roughly 47% efficiency. The variable ranking is dominated by the deflection variables (`wmean`, then `wstd`), with the Bragg ratios contributing essentially nothing, worth noting, since Bragg was expected to carry the stopping signature.

**But it does not fix the binning, and cannot.** Table 15 gives the migration diagonal in the five analysis bins as the cut is tightened. The first bin is already fine and stays fine; no other bin reaches 0.68 at any working point, even at 15% efficiency.

Table 15: Golden-pion BDT: efficiency, achieved golden fraction, and the  $p_\pi$  migration diagonal per true bin, on the held-out run. The BNB criterion for a usable bin is a diagonal above 0.68.

| BDT cut                                 | eff.  | golden frac. | b1   | b2   | b3   | b4   | b5   |
|-----------------------------------------|-------|--------------|------|------|------|------|------|
| none                                    | 100%  | 43.1%        | 99.2 | 23.7 | 13.2 | 6.4  | 5.5  |
| $> -0.05$                               | 57.0% | 58.1%        | 99.1 | 29.4 | 19.7 | 11.2 | 7.1  |
| $> 0.00$                                | 44.8% | 64.9%        | 99.1 | 33.1 | 24.5 | 15.4 | 9.1  |
| $> +0.10$                               | 24.6% | 79.4%        | 99.1 | 42.9 | 36.8 | 29.3 | 15.7 |
| $> +0.15$                               | 15.2% | 84.5%        | 99.2 | 46.9 | 42.2 | 36.1 | 19.5 |
| <i>perfect (truth) golden selection</i> |       |              | 98.9 | 36.1 | 24.1 | 14.2 | 8.9  |

**These diagonals are measured on a different sample from the analysis ones, and the two must not be mixed.** The table above is built on the labelled *track* sample: every truth-matched pion track passing loose quality requirements (purity  $> 0.5$ , track score  $> 0.5$ , length  $> 5$  cm), from events of any topology, 126 518 tracks in total. The migration diagonals quoted elsewhere in this note, 96/35/21/10/6% for five bins and 91/76% for two, are measured on the *analysis* sample: events passing the full  $CC1\mu1\pi^\pm$  selection and satisfying the signal definition, about 10 700 per beam mode. The analysis sample has the better response, because the selection removes exactly the events whose pion is poorly reconstructed. On the track sample the same five-bin scheme reads 99/24/13/6/6% and the two-bin scheme 98/62%.

An earlier draft of this note quoted those track-sample figures as though they were the analysis response, and concluded from the 62% that the inclusive  $p_\pi$  measurement failed the usable-bin criterion. It does not: on the analysis sample the upper bin sits at 76%. The numbers

in this table are correct for what they measure, and the conclusions drawn *within* it, which compare rows of the same sample, are unaffected.

The last row is the decisive one, and it must be read against the *efficiency* column rather than against the tightest cut. Perfect truth-level golden selection keeps 43.1% of tracks and gives 36.1, 24.1, 14.2, 8.9% in bins 2–5; the classifier at the matching 44.8% efficiency gives 33.1, 24.5, 15.4, 9.1%. They agree bin by bin. The classifier is therefore already performing as well as perfect knowledge of which pions stopped, and no better classifier exists to be built.

**Multiple Coulomb scattering was tried as a pion momentum estimator, and as a classifier input.** Range saturates because the pion interacts before stopping, which truncates the track. Multiple-Coulomb-scattering (MCS) momentum is inferred from the scattering along whatever track exists and is not truncated by an interaction, so it is the natural candidate to replace range. The CC0 $\pi$  analysis uses exactly this pairing, range for contained tracks and MCS for exiting ones. Measured on 122 878 truth-matched pion tracks (`trk_mcs_muon_mom_v`, filled for 99.9% of them):

| true $p_\pi$ [GeV/c] | $\langle \text{len} \rangle$ | $\langle \text{range/true} \rangle$ | RMS  | $\langle \text{MCS/true} \rangle$ | RMS  |
|----------------------|------------------------------|-------------------------------------|------|-----------------------------------|------|
| 0.175–0.250          | 25 cm                        | 0.834                               | 0.16 | 4.335                             | 14.1 |
| 0.320–0.420          | 46 cm                        | 0.629                               | 0.23 | 2.438                             | 7.2  |
| 0.550–0.800          | 71 cm                        | 0.446                               | 0.18 | 1.423                             | 3.5  |
| 0.800–1.500          | 84 cm                        | <b>0.319</b>                        | 0.16 | <b>1.011</b>                      | 2.1  |

The trade is stark. In the top bin range is biased to 0.319, which is the saturation, while MCS is essentially unbiased at 1.011; but MCS carries a 210% RMS there and 1400% in the lowest bin. It is unusable as a per-event estimator. The reason is structural and explains why the CC0 $\pi$  recipe does not transfer: they apply MCS to *exiting* tracks, which are long by construction, whereas our problematic pions are contained but interacting, 25–84 cm long, and MCS needs of order a metre.

The same note uses the *disagreement* between the two estimators as a track-quality BDT input rather than as an estimator. That is well motivated here: a pion that interacts has a truncated range but an untruncated MCS, so the fractional difference  $(p_{\text{range}} - p_{\text{MCS}})/p_{\text{range}}$  should flag precisely the events that break the response. Adding it to the golden-pion classifier, it ranks *second of ten* in importance, behind only the mean deflection. It nonetheless changes nothing:

| classifier inputs         | ROC integral | diagonals at $\sim 44\%$ efficiency |
|---------------------------|--------------|-------------------------------------|
| baseline (nine variables) | 0.758        | 99.1/33.1/24.5/15.4/9.1             |
| + range–MCS difference    | 0.757        | 99.0/34.8/24.8/14.8/9.1             |

The interpretation is that the new variable is highly informative but not *additionally* informative: it is strongly correlated with the deflection variables already present, since both measure whether the track scattered heavily and ended abruptly. High importance rank with unchanged ROC is the signature of a redundant input. This is an independent confirmation of the ceiling argument above: the classifier already extracts what golden-ness has to offer, and the limit is the physics of the label rather than the quality of the classifier.

One tempting variant was deliberately not pursued. Retraining on the label “range is accurate”,  $|p_{\text{corr}}/p_{\text{true}} - 1| < 0.2$ , would target the quantity we actually care about and would certainly raise the diagonals. It would also be selecting on the answer: keeping events where the estimator happens to be right sculpts the response by construction and inflates the diagonal without measuring anything, the same trap documented for the tight BDT working point above. The golden label is used precisely because it is a physical property of the pion, independent of how well it was reconstructed.



**A caveat on the tighter working points.** The  $> 0.15$  row looks better than perfect truth selection (46.9, 42.2, 36.1, 19.5%), which is impossible for an honest classifier and is worth understanding rather than quoting. It arises because a hard cut on the classifier preferentially keeps *long* tracks: inside the top true bin, the surviving tracks have a mean length of 120.8 cm against 78.2 cm for all pions, and a mean reconstructed momentum of 0.406 against 0.307 GeV/ $c$ . The cut is thus partly a cut on the reconstructed momentum itself, which raises the diagonal by construction while sculpting the spectrum. The apparent gain is circular and the tight working points should not be used. This is the same failure mode anticipated for track length as a direct input; it enters here through the deflection variables, which correlate with length, even though length itself was excluded from training.

**Why, and what is left.** Golden pions still reconstruct at  $\langle p^{\text{corr}}/p^{\text{true}} \rangle = 0.73$  on this sample. Interaction is therefore not the only problem: the momentum conversion is itself biased. A single flat rescaling does *not* repair it, applying the best global factor drops the first bin from 98.9 to 49.1% and leaves 0/5 bins passing; because the bias is momentum-dependent, exactly as the saturation curve of Figure 21 shows.

Coarser binning helps but does not rescue it either. Collapsing to two bins (0.175, 0.230,  $\infty$ ) gives 98.7 and 58.3% with perfect golden selection: better, still short of 0.68. Only a single bin; that is, no differential measurement at all, trivially passes.

**Conclusion.** The comparison is worth stating plainly rather than filing as a partial success. The classifier does what it was built to do and matches the published benchmark, but the  $p_\pi$  differential measurement above  $\sim 0.23$  GeV/ $c$  cannot be made to meet the BNB’s own binning criterion with a range-based estimator, with or without a golden-pion cut. The BNB analysis passes that criterion because its resolved bins (0.10–0.16, 0.16–0.19, 0.19–0.22) all sit *below* our 0.175 GeV/ $c$  threshold, in the region where range still tracks momentum; above 0.22 it uses a single wide bin and an overflow. The realistic options for this analysis are therefore to lower the pion threshold toward 0.10 GeV/ $c$  and resolve the region where the estimator works, to adopt a two-bin-plus-overflow  $p_\pi$  result and say so, or to replace the range-based estimator altogether. The golden-pion BDT is a useful ingredient in any of those, but it is not by itself the fix.

**Planned change of scope for  $p_\pi$ .** On the strength of the numbers above, the current intention is to *withdraw the pion-momentum differential cross section from the inclusive selection* and to quote it instead on the *proton-tagged* subsample. Both candidate routes were tried; as the two following paragraphs show, the golden-pion classifier does not repair the response even at perfect truth-level selection, whereas the proton tag does not change the response appreciably either way. At the two-bin scheme the inclusive selection already satisfies the binning criterion, so  $p_\pi$  need not be withdrawn from it; what it cannot support is the five-bin *shape* measurement originally planned. The other inclusive observables are unaffected: the angular variables and  $p_\mu$  retain their shape information, and the total cross section does not use this estimator at all.

**The proton-tagged route, now measured in all three configurations.** Both routes have now been quantified. The golden-pion route is above; the proton-tagged one has been run for FHC, RHC and the combined exposure. The expectation stated in earlier drafts, that requiring a tagged proton “does nothing directly to the pion reconstruction”, turns out to be *correct*: at the same two-bin edges the proton-tagged response is indistinguishable from the inclusive one, marginally worse rather than better. Applying the same migration-diagonal test gives



| selection            | config. | $N_{\text{sig}}$ bin 1 | bin 1 [0.175, 0.205] | bin 2 (> 0.205) | both > 0.68? |
|----------------------|---------|------------------------|----------------------|-----------------|--------------|
| inclusive CC1mu1piXp | FHC     | 1056                   | 91.0%                | 75.5%           | yes          |
| <b>proton-tagged</b> | FHC     | 469                    | <b>88.9%</b>         | <b>74.2%</b>    | <b>yes</b>   |
| CC1mu1pi1p           | RHC     | 621                    | <b>89.4%</b>         | <b>75.1%</b>    | <b>yes</b>   |
|                      | comb.   | 1090                   | <b>89.2%</b>         | <b>74.7%</b>    | <b>yes</b>   |

measured on the numuMC overlays named in the corresponding `file_properties` lists, selected signal, with the reconstructed momentum mass-corrected exactly as the bin config does it. All four rows clear the criterion in both bins, and the proton-tagged values sit about two points *below* the inclusive ones rather than above them. The three proton-tagged configurations agree to better than a percentage point, so the response is a property of the selection rather than of a particular exposure, but the proton tag does not improve it.

That is the expected result on reflection, and it corrects a claim made in an earlier draft of this note. The pion-momentum response is set by whether the pion ranges out or interacts hadronically before stopping, which is a property of the pion’s own passage through the argon. Requiring a reconstructed proton at the vertex changes the event sample, the purity and the efficiency, but it does not change what happens to the pion downstream, so there is no mechanism by which it would sharpen the momentum response. The measurement now says so directly.

The consequence is that the proton-tagged  $p_\pi$  measurement stands on its own merits, as the pion-momentum cross section of an exclusive final state that also yields  $W$  and TKI, and not as a repair of the inclusive one. The inclusive  $p_\pi$  at two bins is equally defensible on the response test.

Table 16: Two-bin  $p_\pi$  on the proton-tagged CC1mu1pi1p selection, all three beam configurations.  $d\sigma/dp_\pi$  in  $10^{-38} \text{ cm}^2/(\text{GeV}/c)/\text{Ar}$ ; bin 1 is [0.175, 0.205] GeV/c, bin 2 everything above. All curves are  $A_C$ -smeared. The proton-tagged `xsec` configs declare only the tune and the fake data, so no generator rows appear.

|                       | FHC                   |             | RHC                   |             | Combined              |             |
|-----------------------|-----------------------|-------------|-----------------------|-------------|-----------------------|-------------|
|                       | bin 1                 | bin 2       | bin 1                 | bin 2       | bin 1                 | bin 2       |
| unfolded data         | 1.554                 | 0.691       | 2.448                 | 0.533       | 2.358                 | 0.589       |
| uncertainty           | $\pm 0.896$           | $\pm 0.243$ | $\pm 0.780$           | $\pm 0.202$ | $\pm 0.832$           | $\pm 0.212$ |
| $A_C$ truth           | 1.450                 | 0.598       | 1.300                 | 0.522       | 1.537                 | 0.544       |
| uB tune               | 0.926                 | 0.497       | 0.799                 | 0.437       | 1.032                 | 0.445       |
| unfolded/truth        | 1.07                  | 1.16        | 1.88                  | 1.02        | 1.53                  | 1.08        |
| $\sigma_{\text{int}}$ | 0.596                 |             | 0.497                 |             | 0.539                 |             |
| $\chi^2/\text{ndf}$   | 0.15/2 ( $p = 0.93$ ) |             | 2.36/2 ( $p = 0.31$ ) |             | 1.01/2 ( $p = 0.60$ ) |             |

All three extractions close, with  $\chi^2$  of 0.15, 2.37 and 1.01 on two bins ( $p = 0.93, 0.31, 0.60$ ). The  $p_\pi$  differential cross section satisfies the field’s binning criterion in every bin of every beam configuration here, as it does on the inclusive selection; the two are equivalent on that test, and the proton-tagged version is reported because it belongs to the exclusive final state, not because it is required to make  $p_\pi$  measurable.

**One number that needs stating plainly.** The RHC first bin unfolds to 1.88 times its  $A_C$ -smeared truth, against 1.07 for FHC and 1.53 for combined. Quoted as a ratio that looks alarming, but the denominator is small and the absolute discrepancy is  $2.448 - 1.300 = 1.15$  against an uncertainty of 0.78, i.e.  $1.5\sigma$ ; the closure  $\chi^2$  of  $2.37/2$  ( $p = 0.31$ ) is unremarkable. The combined value sits between the two, as it must, being dominated by neither. The honest reading is an upward fluctuation in the least-populated bin of this fake-data throw rather than a




Figure 24: The two-bin  $p_\pi$  cross section on the proton-tagged `CC1mu1pi1p` selection, the graphical form of Tab. 16. Same edges as the inclusive result of Fig. 22 so the two are directly comparable, and the bins are again drawn equal-width.

defect in the extraction, but it is the one place in the proton-tagged set where the central value is not close to truth, and it should be watched when the throw is repeated.

One caveat remains. The split at 0.205 was inherited from the inclusive optimisation rather than re-derived for this selection, so it is a working point that passes comfortably in all three configurations, not necessarily the best available here. A dedicated scan would need its own universe construction per configuration ( $\sim 1\text{--}2$  h each) and is not expected to change the conclusion, since both bins clear the criterion with margin.

## 8.9 Model comparison: $\chi^2$ against truth, tune and generators

Table 17 collects the bin-by-bin  $\chi^2$  of the unfolded fake data against each prediction, for every observable and every beam configuration. All predictions are  $A_C$ -smeared, so the comparison is made in the measurement space, and the full covariance is used. The *truth* column is the closure test: it is small throughout, confirming the chain, and is visibly larger for RHC than for FHC, consistent with the poorer RHC angular response noted in §9.2.

**The  $\cos\theta_\mu$   $\chi^2$  is largely an artefact of the variable.** One entry stands out: in  $\cos\theta_\mu$  the NEUT and NuWro  $\chi^2$  reach 26.3/5 and 26.0/5 for FHC (and 18.4, 17.4 for RHC), an order of magnitude above every other observable. Taken at face value this would be strong evidence against those two generators. It is not. Repeating the identical comparison in  $\theta_\mu$ ; the same events, the same data, the same bins mapped through arccos, gives 2.8/5 and 2.9/5, a factor  $\sim 9$  smaller (Table 18). The mechanism is the  $A_C$  suppression discussed in §9.2: in  $\cos\theta_\mu$  the additional smearing suppresses the sharply peaked external generators by a factor  $\sim 0.33$  while leaving the smoother MicroBooNE tune much less affected, so NEUT and NuWro are pushed away from the data and the  $\chi^2$  inflates. In  $\theta_\mu$ , where the suppression is  $\sim 0.76$ , the same models agree well. The pion angle, which is not sharply peaked, is essentially unchanged between the two parameterisations. This is a further argument for quoting the angular cross sections in  $\theta$  rather than  $\cos\theta$ , and a caution against reading model discrimination directly off a  $\cos\theta_\mu$   $\chi^2$ .

Table 17:  $\chi^2/\text{ndf}$  of the unfolded fake data against each  $A_C$ -smeared prediction, for the inclusive selection. “truth” is the fake-data truth (the closure test).

| Beam | Observable        | truth  | uB tune | GENIE   | GiBUU   | NEUT    | NuWro   |
|------|-------------------|--------|---------|---------|---------|---------|---------|
| FHC  | $p_\mu$           | 2.19/7 | 3.67/7  | 5.52/7  | 3.23/7  | 3.25/7  | 2.20/7  |
| FHC  | $p_\pi$           | 0.15/2 | 0.93/2  | 2.01/2  | 0.60/2  | 0.16/2  | 0.23/2  |
| FHC  | $\cos \theta_\mu$ | 0.59/5 | 4.41/5  | 11.27/5 | 15.68/5 | 29.78/5 | 29.10/5 |
| FHC  | $\cos \theta_\pi$ | 0.56/4 | 2.40/4  | 4.70/4  | 2.27/4  | 2.31/4  | 2.71/4  |
| FHC  | $\theta_{\mu\pi}$ | 1.83/5 | 2.26/5  | 4.06/5  | 3.45/5  | 0.85/5  | 0.59/5  |
| FHC  | $\theta_\mu$      | 2.17/5 | 5.14/5  | 11.30/5 | 17.91/5 | 40.60/5 | 38.62/5 |
| RHC  | $p_\mu$           | 2.93/7 | 9.73/7  | 11.28/7 | 9.09/7  | 8.62/7  | 8.38/7  |
| RHC  | $p_\pi$           | 0.24/2 | 0.74/2  | 2.56/2  | 1.12/2  | 0.18/2  | 0.37/2  |
| RHC  | $\cos \theta_\mu$ | 2.99/5 | 4.79/5  | 5.31/5  | 6.21/5  | 12.95/5 | 12.16/5 |
| RHC  | $\cos \theta_\pi$ | 1.64/4 | 3.12/4  | 5.70/4  | 2.96/4  | 2.92/4  | 4.38/4  |
| RHC  | $\theta_{\mu\pi}$ | 4.57/5 | 4.68/5  | 7.46/5  | 6.49/5  | 4.40/5  | 5.57/5  |
| RHC  | $\theta_\mu$      | 3.44/5 | 4.11/5  | 8.03/5  | 12.69/5 | 34.23/5 | 32.11/5 |
| Comb | $p_\mu$           | 3.27/7 | 4.78/7  | 6.41/7  | 4.71/7  | 4.71/7  | 4.01/7  |
| Comb | $p_\pi$           | 0.13/2 | 0.80/2  | 1.96/2  | 0.88/2  | 0.19/2  | 0.32/2  |
| Comb | $\cos \theta_\mu$ | 3.08/5 | 6.22/5  | 8.69/5  | 10.56/5 | 19.49/5 | 19.63/5 |
| Comb | $\cos \theta_\pi$ | 0.43/4 | 2.15/4  | 4.11/4  | 1.96/4  | 2.15/4  | 3.32/4  |
| Comb | $\theta_{\mu\pi}$ | 2.31/5 | 2.25/5  | 4.04/5  | 4.34/5  | 1.52/5  | 1.34/5  |
| Comb | $\theta_\mu$      | 2.26/5 | 4.10/5  | 6.04/5  | 6.23/5  | 12.79/5 | 13.14/5 |

Table 18: The same comparison for the muon and pion angle in both parameterisations. The  $\theta$  bin edges are exactly arccos of the  $\cos \theta$  edges, so the phase space is identical; only the variable differs. *With the width-aware regularisation the two parameterisations now agree:* both disfavour NEUT and NuWro strongly in every beam configuration. Previously FHC  $\theta_\mu$  returned  $\chi^2 \approx 2.8\text{--}2.9/5$  for those models against  $\approx 26/5$  in  $\cos \theta_\mu$ , which was the rank-1 artefact, not a property of the variable.

| Beam | Variable          | truth  | uB tune | GENIE   | GiBUU   | NEUT    | NuWro   |
|------|-------------------|--------|---------|---------|---------|---------|---------|
| FHC  | $\cos \theta_\mu$ | 0.59/5 | 4.41/5  | 11.27/5 | 15.68/5 | 29.78/5 | 29.10/5 |
| FHC  | $\theta_\mu$      | 2.17/5 | 5.14/5  | 11.30/5 | 17.91/5 | 40.60/5 | 38.62/5 |
| RHC  | $\cos \theta_\mu$ | 2.99/5 | 4.79/5  | 5.31/5  | 6.21/5  | 12.95/5 | 12.16/5 |
| RHC  | $\theta_\mu$      | 3.44/5 | 4.11/5  | 8.03/5  | 12.69/5 | 34.23/5 | 32.11/5 |
| Comb | $\cos \theta_\mu$ | 3.08/5 | 6.22/5  | 8.69/5  | 10.56/5 | 19.49/5 | 19.63/5 |
| Comb | $\theta_\mu$      | 2.26/5 | 4.10/5  | 6.04/5  | 6.23/5  | 12.79/5 | 13.14/5 |
| FHC  | $\cos \theta_\pi$ | 0.56/4 | 2.40/4  | 4.70/4  | 2.27/4  | 2.31/4  | 2.71/4  |
| FHC  | $\theta_\pi$      | 0.52/4 | 1.92/4  | 4.31/4  | 1.86/4  | 2.63/4  | 3.56/4  |

## 8.10 Proton-tagged $\chi^2$ and the $A_C$ correlation

### Withdrawn pending re-extraction.

Every  $\chi^2$  in this subsection compares a measurement of the transverse variables taken about the *detector  $z$  axis* with generator predictions taken about the *neutrino direction* (Sec. 2.2). They are not the same variables, and the numbers below therefore do not measure model agreement.

The size of the mismatch is not subtle. Under the two conventions the true  $\delta p_T$  distribution over the current bin edges runs 20.7/17.1/14.7/15.2/16.1/16.3% about  $z$  against 80.8/10.8/4.5/2.5/1.2/0.2% about the neutrino; the generator predictions sit at 77.3/12.3/5.1/3.2/1.9/0.3% (GENIE) and 80.9/10.1/4.4/2.7/1.6/0.3% (NEUT), i.e. they agree closely with the beam-frame truth and not at all with what was measured against them. The large  $\chi^2$  values below are that definitional mismatch, not physics. That the generators reproduce the beam-frame truth this well is also a useful independent check of the correction itself.

The bin edges are affected as well as the  $\chi^2$ : they were chosen as equal-population quantiles of the uncorrected variable, which is why the old occupancies above are flat. They are being re-derived along with the re-extraction, and the conclusion drawn here — that the  $\chi^2$  ordering tracks  $|A_C - 1|$  — must be re-established from scratch afterwards rather than assumed to survive.

Table 19 repeats the comparison for the six proton-tagged observables (FHC), with the measured additional-smearing factor of §10.2 in the last column. Ordering the rows by  $|A_C - 1|$  makes an unambiguous pattern visible: the generator  $\chi^2$  grows monotonically with the distance of  $A_C$  from unity, and does so in the same direction for every model.

| Observable       | $A_C$ | $ A_C - 1 $ | typical generator $\chi^2/6$ |
|------------------|-------|-------------|------------------------------|
| $W_{\text{had}}$ | 0.69  | 0.31        | 4.1                          |
| $W_{\pi p}$      | 0.77  | 0.23        | 1.6                          |
| $\delta\alpha_T$ | 0.71  | 0.29        | 7.1                          |
| $\delta\phi_T$   | 0.80  | 0.20        | 15.5                         |
| $\delta p_T$     | 0.67  | 0.33        | 70                           |
| $p_n$            | 0.69  | 0.31        | 76                           |

*This argument was made with the pre-fix  $A_C$  and does not survive.* With the uniform-grid regularisation the smearing factors ranged from 0.43 to 1.71, and  $p_n$  was *amplified* by 71%, which is not something a physical smearing matrix should do; on that set  $|A_C - 1|$  did order the generator  $\chi^2$ , and the ordering was read as a smearing artefact. With the width-aware regularisation the factors cluster at 0.67–0.80 and  $|A_C - 1|$  is flat to within 0.13 across observables, while the generator  $\chi^2$  still spans 1.6 to 76 per six bins. A flat quantity cannot order a varying one, so the artefact explanation is withdrawn: the  $\chi^2$  ordering of the TKI observables is physics.

That is the expected behaviour rather than a surprise.  $\delta p_T$  and  $p_n$  probe the nuclear initial state and final-state interactions directly, and generators differ most there; the previous  $A_C$  was flattening precisely the observables built to discriminate. The same reassessment applies to  $\cos\theta_\mu$  (§8.9), where reparameterising to  $\theta_\mu$  reduced the NEUT and NuWro  $\chi^2$  by a factor  $\sim 9$  on identical data. The closure column is excellent throughout ( $\leq 0.46/6$ ), so the chain itself is sound; it is the *model comparison* in  $\delta p_T$  and  $p_n$  that should not be read as discrimination until the binning and response of those two observables are revisited.

Table 19:  $\chi^2/\text{ndf}$  of the unfolded fake data against each  $A_C$ -smeared prediction for the proton-tagged observables (FHC), with the additional-smearing factor  $A_C$ .

| Observable       | truth  | uB tune | GENIE   | GiBUU   | NEUT     | NuWro   | $A_C$ |
|------------------|--------|---------|---------|---------|----------|---------|-------|
| $W_{\text{had}}$ | 0.86/6 | 2.50/6  | 4.51/6  | 3.87/6  | 3.46/6   | 4.71/6  | 0.69  |
| $W_{\pi p}$      | 0.19/6 | 1.65/6  | 2.59/6  | 1.11/6  | 0.85/6   | 1.87/6  | 0.77  |
| $\delta\alpha_T$ | 0.67/6 | 3.46/6  | 7.41/6  | 6.74/6  | 7.27/6   | 6.83/6  | 0.71  |
| $\delta\phi_T$   | 4.02/6 | 5.96/6  | 12.22/6 | 15.40/6 | 17.60/6  | 16.82/6 | 0.80  |
| $\delta p_T$     | 1.87/6 | 6.37/6  | 44.43/6 | 69.31/6 | 91.61/6  | 76.04/6 | 0.67  |
| $p_n$            | 0.19/6 | 4.93/6  | 39.79/6 | 74.32/6 | 108.00/6 | 83.83/6 | 0.69  |

### 8.11 The $W$ binnings are finer than the resolution

$W_{\pi p}$  shows the least model tension of the proton-tagged observables, and it was also the worst-conditioned  $A_C$  in the analysis before the regularisation fix ( $s_2/s_1 = 0.022$ , against 0.2–0.25 for the rest). Those two facts have a common cause, and it is not physics.

The bin edges were chosen for equal population, six bins of roughly equal content, without reference to the resolution. Measured on selected signal, FHC Runs 1/2/4:

|                            | $W_{\pi p}$   | $W_{\text{had}}$ |
|----------------------------|---------------|------------------|
| residual RMS [GeV/ $c^2$ ] | 0.142         | 0.283            |
| robust half-width (16–84%) | 0.101         | 0.177            |
| bias (mean reco – true)    | <b>−0.111</b> | −0.032           |
| narrowest analysis bin     | <b>0.040</b>  | 0.100            |
| widest analysis bin        | 1.430         | 1.300            |
| width ratio                | <b>35.7</b>   | 13.0             |

The second and third  $W_{\pi p}$  bins are 40 MeV/ $c^2$  wide against a resolution of 101–142 MeV/ $c^2$ : they are two to three times narrower than the detector can resolve. The bias alone, −111 MeV/ $c^2$ , is larger than three of the six bin widths. The migration diagonals follow directly, 87.6/22.3/11.7/12.0/10.0/2.3%: only the first bin is measured, and the top bin at 2.3% is essentially unmeasured.

This is the same failure as the five-bin  $p_\pi$  scheme, and for the same underlying reason.  $W_{\pi p}$  is reconstructed from the pion and proton momenta, so it inherits the saturation of the range-based pion momentum described in §8.8; its resolution cannot be better than that of its inputs.

An exhaustive edge scan, the same procedure and criterion used for  $p_\pi$ , gives:

| scheme                     | diagonals                     | verdict            |
|----------------------------|-------------------------------|--------------------|
| current, six bins          | 87.6/22.3/11.7/12.0/10.0/2.3% | 1 of 6 usable      |
| best two bins (split 1.15) | <b>83.2/77.9%</b>             | <b>both usable</b> |
| best three bins            | 83.2/49.7/45.9%               | fails              |
| best four bins             | 83.2/45.0/28.2/28.2%          | fails              |

As with  $p_\pi$ , two bins is not merely the best option but the only one that satisfies the criterion, and the first bin is thin (137 selected signal events, against a floor of 100). The implication for the model comparison is direct: the low  $\chi^2$  currently quoted for  $W_{\pi p}$  reflects a binning that cannot resolve the differences between generators, not agreement between them. The same test should be applied to  $W_{\text{had}}$ , whose upper bins are also narrower than its 177 MeV/ $c^2$  half-width, before either  $W$  result is quoted as a shape measurement.

**Adopted scheme: two bins split at 1.15 GeV/ $c^2$ .** The boundary is more tightly constrained than it was for  $p_\pi$ , where anything between roughly 0.19 and 0.22 worked. Here only one choice satisfies both the diagonal criterion and the 100-event floor:

| boundary [GeV/ $c^2$ ] | bin 1        | bin 2        | $N(\text{bin 1})$ |                       |
|------------------------|--------------|--------------|-------------------|-----------------------|
| 1.13                   | 76.5%        | 88.0%        | 51                | below the event floor |
| <b>1.15</b>            | <b>83.2%</b> | <b>77.9%</b> | <b>137</b>        | <b>adopted</b>        |
| 1.17                   | 86.8%        | 66.8%        | 266               | bin 2 just fails      |
| 1.19                   | 87.6%        | 56.0%        | 468               | fails                 |
| 1.23                   | 91.7%        | 37.6%        | 1114              | fails                 |

Moving the boundary up buys population in the first bin at the direct expense of the second: the two diagonals trade against each other steeply, and the window in which both exceed 0.68 is only about  $\pm 0.01$  GeV/ $c^2$  wide. That narrowness is itself a consequence of the resolution being comparable to the full width of the peak region.

Two consequences should be carried with the result. First, **the first bin is thin**, 137 selected signal events against a floor of 100, so its statistical uncertainty will be large once systematics are folded in; the  $p_\pi$  first bin, with 469 events, already carries 42–53%. Second, if that proves too sparse in practice, 1.17 is the natural fallback: it doubles the population for a bin-2 diagonal of 66.8%, marginally below threshold, which is a defensible trade to state explicitly rather than a criterion to fail silently.

The scheme is implemented in `configs/ccpi1p_Wpipr_bin_config_2bin.txt`, with the reasoning recorded alongside it in a companion `.README`. At the time of writing the universe construction for the FHC configuration is in progress; the extracted cross section, its closure and the resulting generator  $\chi^2$  are not yet available, and the comparison that matters, whether the apparent  $W_{\pi p}$  agreement survives a binning the detector can actually resolve, remains to be made. The same test has not yet been applied to  $W_{\text{had}}$ .

## 8.12 What limits $W_{\pi p}$ , and what can be done about it

The two-bin  $W_{\pi p}$  scheme of §8.11 was built and run. It does not work, for a reason distinct from the six-bin failure, and the investigation that followed settles what the pion momentum can and cannot deliver.

**The two-bin extraction fails on statistics.** The *response* behaves exactly as designed: the framework returns migration diagonals of 0.802 and 0.773, matching the truth-level scan, and the closure is acceptable at  $\chi^2 = 1.17/2$  ( $p = 0.56$ ). The *extraction* nonetheless carries almost no shape information, because  $A_C$  collapses:

| two-bin scheme                    | $A_C$                                                           | $s_2/s_1$    |
|-----------------------------------|-----------------------------------------------------------------|--------------|
| $W_{\pi p}$ , first bin $N = 137$ | $\begin{pmatrix} 0.018 & -0.410 \\ 0.034 & 0.892 \end{pmatrix}$ | <b>0.031</b> |
| $p_\pi$ proton-tagged, $N = 469$  | $\begin{pmatrix} 0.498 & -0.089 \\ 0.013 & 0.854 \end{pmatrix}$ | 0.577        |
| $p_\pi$ inclusive, $N = 1056$     | $\begin{pmatrix} 0.594 & -0.382 \\ 0.011 & 0.891 \end{pmatrix}$ | 0.528        |

The top-left entry is 0.018:  $A_C$  passes under two per cent of the first bin and takes  $-0.41$  from the second. This is a rank-one collapse of the kind described in §9.2, but its cause is statistical rather than a defective regulariser. With 137 events the first bin has such poor signal-to-noise that the Wiener filter suppresses that mode almost entirely, and the ordering across the three schemes,  $137 \rightarrow 0.031$ ,  $469 \rightarrow 0.577$ ,  $1056 \rightarrow 0.528$ , follows population rather than anything geometric.

$W_{\pi p}$  is therefore caught between two requirements. Bins narrow enough to satisfy the response criterion leave the first bin too sparse to unfold; bins wide enough for a stable unfold (1.19 gives 468 events, matching the  $p_\pi$  case that works) fail the criterion at 56%.

**No binning escapes it.** The conclusion does not rest on the one scheme that was tried. Scanning *every* two- and three-bin scheme with edges between 1.12 and 1.70 GeV/ $c^2$  (FHC Run1+2+4, 3505 selected signal events) finds none that keeps both a migration diagonal above 0.68 and adequate statistics in every bin:

| two-bin edge  | $N_1/N_2$ | diag <sub>1</sub> / diag <sub>2</sub> |                      |
|---------------|-----------|---------------------------------------|----------------------|
| 1.15          | 137/3367  | 0.832/0.779                           | first bin too sparse |
| 1.18          | 351/3153  | 0.866/0.611                           |                      |
| 1.24          | 1298/2206 | 0.930/0.347                           |                      |
| 1.27 (median) | 1744/1760 | 0.951/ <b>0.261</b>                   | upper bin unusable   |
| 1.45          | 2966/538  | 0.993/0.030                           |                      |

The trend exposes the mechanism. Reconstructed  $W_{\pi p}$  is biased about 111 MeV low, inheriting the saturation of the range-based pion momentum (§8.8), so raising the edge far enough to balance the statistics drains the upper bin by downward migration: diag<sub>1</sub> climbs from 0.786 to 0.999 across the scan while diag<sub>2</sub> collapses from 0.927 to 0.004. There is no edge at which both bins are simultaneously populated and diagonal, and no three-bin combination does better — the third-bin diagonal never exceeds 0.347.

**At the current exposure  $W_{\pi p}$  does not support a shape measurement** and is quoted as an integrated cross section unless the pion momentum improves. That integrated number is a cut-and-count measurement rather than an unfolded one: with a single bin there is nothing to unfold, since the response is  $1 \times 1$ ,  $A_C$  is trivially unity and the regularisation has no effect. It is reported through the total flux-averaged cross-section path of §9.1, and integrated over the full range it is the same quantity as the total proton-tagged cross section. The corresponding integrated generator predictions are built by `generator_predictions/newg4/make_wpipr_1bin.C`, summing the per-bin cross sections of the six-bin flux-averaged predictions.

**The limitation is the pion momentum, and only the pion momentum.** Correlating the  $W_{\pi p}$  residual against each input residual isolates the cause:

| input                | $r$           | $W$ residual RMS when...           |              |
|----------------------|---------------|------------------------------------|--------------|
| pion momentum        | <b>+0.726</b> | all events                         | 0.142        |
| proton momentum      | +0.012        | pion momentum well reconstructed   | <b>0.061</b> |
| pion $\cos \theta$   | -0.029        | proton momentum well reconstructed | 0.130        |
| proton $\cos \theta$ | -0.032        | both well reconstructed            | 0.038        |

Fixing the pion momentum alone takes the resolution from 0.142 to 0.061, a factor 2.3; fixing the proton buys almost nothing, and the angles are irrelevant. A single defect propagates through the whole analysis: the hadronic interaction that truncates the pion track forces  $p_\pi$  to two bins, and forces  $W_{\pi p}$  out of a shape measurement altogether.

**Reparameterising to  $W^2$  does not help.** A monotonic transformation with mapped edges leaves the migration matrix unchanged, which was verified directly: splitting at 1.15 GeV/ $c^2$  in  $W$  and at  $1.15^2 = 1.3225$  GeV<sup>2</sup>/ $c^4$  in  $W^2$  gives diagonals of 83.2% and 77.9% in both cases, on identical events. Resolution and bin width scale together,  $\sigma(W^2) = 2W\sigma(W)$ , so  $\sigma/\text{width}$  is invariant at about 2 either way. This is the same lesson as  $\theta_\mu$  versus  $\cos \theta_\mu$  (§9.2).

### 8.13 Survey of pion momentum estimators without a magnetic field

Four approaches were measured on truth-matched pion tracks. All are limited by the same mechanism, that each measures what the pion did *before* it interacted:

| estimator             | $\langle \text{reco/true} \rangle$ , top bin | RMS  | note                                     |
|-----------------------|----------------------------------------------|------|------------------------------------------|
| range (analysis)      | 0.319                                        | 0.16 | saturates near 0.34 GeV/ $c$             |
| calorimetric energy   | 0.383                                        | 0.31 | also truncated, and $2\times$ noisier    |
| MCS                   | 1.011                                        | 2.1  | unbiased, unusably noisy on short tracks |
| golden-pion selection | 0.327                                        | 0.18 | no gain above 0.8 GeV/ $c$               |

**Calorimetry does not escape the truncation.** Deposited charge stops at the interaction point just as range does, so  $\langle \text{calo/true} \rangle$  falls from 1.136 to 0.383 across the momentum range while the RMS is three to five times that of range (1.007 to 0.313, against 0.157 to 0.158). It is the same measurement with more noise.

**The golden-pion selection is weaker than previously described.** Splitting the sample by the truth label shows that golden pions are not well reconstructed either, and that their advantage disappears exactly where it is needed:

| true $p_\pi$ [GeV/ $c$ ] | golden fraction | golden $\langle \text{range/true} \rangle$ | non-golden   |
|--------------------------|-----------------|--------------------------------------------|--------------|
| 0.175–0.250              | 77.8%           | 0.873                                      | 0.699        |
| 0.320–0.420              | 44.1%           | 0.705                                      | 0.570        |
| 0.550–0.800              | 21.4%           | 0.507                                      | 0.429        |
| 0.800–1.500              | 17.2%           | <b>0.327</b>                               | <b>0.317</b> |

Above 0.8 GeV/ $c$  the golden and non-golden values are identical to within one per cent. The reason is physical: a 1 GeV/ $c$  pion has a CSDA range of several metres in argon and cannot stop inside a 2.5 m TPC unless it has already lost most of its energy through processes that truncate or break the track. “Golden” at high momentum therefore selects a pathological subsample rather than a clean one, which is why the classifier of §8.8 reaches a ceiling: above  $\sim 0.5$  GeV/ $c$  there is no clean subsample to find.

**A multivariate regression fixes the bias but redistributes the resolution.** A BDTG regression on  $p_\pi^{\text{true}}$ , using range, MCS, calorimetry, length, the three deflection variables, the four Bragg likelihoods, track score and daughter count, was trained on Runs 1 and 4 and evaluated on the held-out Run 2. It largely removes the saturation in the mean,  $\langle \text{pred/true} \rangle$  running 1.196 to 0.722 against 0.835 to 0.319 for range, at comparable RMS. Rebuilding  $W_{\pi p}$  with it:

| pion estimator         | $W$ bias       | $W$ RMS      |
|------------------------|----------------|--------------|
| range (analysis)       | −0.127         | 0.144        |
| BDTG regression        | − <b>0.005</b> | 0.119        |
| true $p_\pi$ (ceiling) | +0.003         | <b>0.046</b> |

The bias falls by a factor of 25, which matters independently of binning: a  $-127 \text{ MeV}/c^2$  shift distorts the whole  $W$  distribution. The resolution improves only from 0.144 to 0.119 against a ceiling of 0.046, so the regression recovers about a quarter of what is available.

Its effect on the binning is a redistribution rather than a rescue. The six-bin diagonals move from 98/30/19/19/13/6% to 63/38/33/43/52/56%: the top bin improves nearly tenfold, but the first bin degrades from 98% to 63%, and in the two-bin scheme the worst bin falls from 73% to 58%. That degradation is regression to the mean, the estimator being pulled toward the sample centre, and it damages the low- $W$  region where the resonance physics sits. The same pattern appears in  $p_\pi$  itself, where the first bin falls from 99.2% to 60.6% while the top bin rises from 0.5% to 33.2%.

Two objections stand against adopting it. The estimator would be MC-trained and sits upstream of everything, so its model dependence becomes a systematic on every result rather than on one cut. And the trade it offers moves measurability out of the region of interest. It is not adopted here.



**What would actually work.** The ceiling measurements above bound what is available: a perfect pion momentum would give  $W_{\pi p}$  a resolution of 0.046, a factor of three better than the regression achieves and enough to support several bins. The information is carried away by the interaction products, so the approach that attacks the cause is to recover their energy: the pion and its daughters treated as a mini-calorimeter, using the per-PFP calorimetric energies and daughter links already present in the input files. For the proton-tagged sample there is a second, independent handle in transverse kinematic imbalance, where the muon and proton constrain the pion transverse momentum without using the pion’s own reconstruction at all, at the cost of Fermi-motion smearing of order 250 MeV/ $c$ . Neither has been tried.

**Recovering the interaction products does not work either.** The ceiling measurements above locate the missing information in the secondaries produced when the pion interacts, so the natural remedy is to treat the pion and its products as a single calorimeter. This was tested. PeLEE carries no parent index, so association is geometric, by PFP start point relative to the pion track end, optionally restricted to the generation immediately below the pion in the Pandora hierarchy. The hierarchy restriction matters: at a 40 cm radius the unrestricted sum collects 0.371 GeV in the lowest momentum bin, where only 0.034 GeV is missing, so it is dominated by unrelated activity.

With the hierarchy cut applied, on Run 1:

| true $p_\pi$ [GeV/ $c$ ] | $\langle \text{KE missing} \rangle$ | $\langle \sum E \text{ of daughters within } R \rangle$ [GeV] |       |       |              |
|--------------------------|-------------------------------------|---------------------------------------------------------------|-------|-------|--------------|
|                          |                                     | 5 cm                                                          | 10 cm | 20 cm | 40 cm        |
| 0.175–0.420              | 0.034                               | 0.023                                                         | 0.026 | 0.029 | 0.037        |
| 0.420–0.800              | 0.246                               | 0.040                                                         | 0.048 | 0.055 | 0.062        |
| 0.800–1.500              | <b>0.659</b>                        | 0.056                                                         | 0.074 | 0.091 | <b>0.104</b> |

“KE missing” is the pion’s true kinetic energy less the calorimetric energy of its own track. In the top momentum bin 659 MeV is unaccounted for and the visible daughters supply 104 MeV, of which about 37 MeV is the baseline seen even in the lowest bin where almost nothing is missing. Genuine recovery is therefore of order 10% at high momentum and 25% in the middle bin.

*It also does not change the binning assessment*, which was tested directly rather than inferred from the bias. Adding the daughter energy improves  $\langle \text{reco/true} \rangle$  everywhere, the top bin moving from 0.373 to 0.470 and the first bin becoming nearly unbiased at 0.953, and it roughly triples the upper migration diagonals. Neither is enough: the best upper diagonal rises from 5.5% to 15.7%, against the 68% required. In the adopted two-bin scheme it is a wash and arguably a loss, the worst bin improving from 61.4% to 71.0% while the first degrades from 97.7% to 82.4%, the same regression-to-the-mean behaviour seen with the multivariate estimator. Since both bins already clear the criterion with plain range, nothing is rescued and the daughter association would be imported as a new systematic for no gain. **The two-bin choice for  $p_\pi$  is therefore robust against every estimator tested**, which is a stronger statement than the binning study alone could make.

That recovery is far too little to change the measurement, and the reason is physical rather than algorithmic. Pion absorption on argon,  $\pi A \rightarrow NN \dots$ , deposits much of the energy in nucleons, roughly half of which are neutrons and therefore invisible in liquid argon; the charged products are frequently short and fall below the threshold to be reconstructed as separate particles. The energy is not merely unassociated, it is largely undetected. Applying the daughter sum would move  $\langle \text{reco/true} \rangle$  in the top bin from 0.32 to about 0.42, still less than half of truth.

**Where this leaves the pion momentum.** Five approaches have now been measured: range, calorimetry, MCS, golden-pion tagging and a multivariate regression, together with recovery of the interaction products. Each is limited by the same fact, that a charged pion which interacts before stopping destroys the information a non-magnetised detector uses to infer its momentum,

and that a substantial part of the energy leaves the detector unseen. The regression is the only one that materially improves anything, removing the  $W_{\pi p}$  bias, and it does so at the cost of model dependence and of the low- $W$  region.

One avenue remains untried and is independent of the pion’s own reconstruction: for the proton-tagged sample, transverse kinematic imbalance constrains the pion transverse momentum from the muon and proton alone. It is limited by Fermi motion, of order 250 MeV/ $c$ , which is comparable to the present errors rather than better than them, so it is worth measuring but should not be expected to transform the measurement. Failing that, the honest conclusion is that  $p_\pi$  is a two-bin measurement and  $W_{\pi p}$  an integrated one at this exposure, and that both are limited by physics rather than by analysis choices.

## 8.14 Cross-check methods

Seven additional methods are run in parallel:

Table 20: Unfolding methods.

| Label                     | Description                                        |
|---------------------------|----------------------------------------------------|
| <code>wiener_svd</code>   | Wiener-SVD (primary result)                        |
| <code>bin_by_bin</code>   | Efficiency correction only; no migration unfolding |
| <code>ibu</code>          | D’Agostini iterative Bayesian, 4 iterations        |
| <code>tikhonov</code>     | Tikhonov-regularised least-squares, $\tau = 1.0$   |
| <code>roo_bayes</code>    | RooUnfoldBayes, 4 iterations                       |
| <code>roo_svd</code>      | RooUnfoldSvd                                       |
| <code>roo_invert</code>   | RooUnfoldInvert (direct matrix inversion)          |
| <code>roo_binbybin</code> | RooUnfoldBinByBin                                  |

The D’Agostini iterative-Bayesian method provides the principal quantitative cross-check of the Wiener-SVD result. Because it does not apply the Wiener-SVD additional-smearing matrix  $A_C$ , its unfolded total is free of the (non-norm-preserving)  $A_C$  suppression and is stable against the iteration count. Table 21 compares the two integrated totals for all three configurations: the D’Agostini result sits above the Wiener-SVD total by  $\sim 23\%$  (FHC) and  $\sim 37\text{--}38\%$  (RHC, Combined), equivalently  $A_C$  suppresses the quoted Wiener-SVD integral by  $\sim 19\%$  (FHC) and  $\sim 27\%$  (RHC/Combined), the larger suppression reflecting the coarser RHC/combined smearing, and it varies by  $< 1\%$  across 1, 2 and 4 iterations. For FHC the D’Agostini per-observable truth is flat to 1% (1.111–1.122), directly confirming that the observable-to-observable spread in the Wiener-SVD integrated cross sections (Table 25) is the  $A_C$  smearing rather than a physical or acceptance effect.

Table 21: Wiener-SVD vs. D’Agostini (iterative Bayesian, 4 iterations) integrated cross section, averaged over the five observables (fake-data truth,  $10^{-38}$  cm<sup>2</sup>/Ar); both columns recomputed with the corrected MC POT normalisation. D’Agostini carries no  $A_C$  smearing, so it recovers a 25–29% higher total. Crucially the ratio is now *consistent* across the three configurations (1.25–1.29), whereas with the earlier, incorrect normalisation the RHC and combined ratios (1.37, 1.38) sat well above FHC (1.23); the inconsistency was an artefact of the inflated RHC/combined Wiener-SVD values, not of the unfolding methods. D’Agostini is also nearly flat across observables (spread 1.2% FHC, 2.2% RHC, 1.5% combined), in contrast to the Wiener-SVD spread, confirming that the observable-to-observable variation of  $\sigma_{\text{int}}$  in Table 25 is the  $A_C$  smearing rather than a physical or normalisation effect.

| Config   | $\langle\sigma\rangle_{\text{WSVD}}$ | $\langle\sigma\rangle_{\text{DAg}}$ | DAg/WSVD | DAg 1→4 iter |
|----------|--------------------------------------|-------------------------------------|----------|--------------|
| FHC      | 0.923                                | 1.170                               | 1.27     | < 1%         |
| RHC      | 0.836                                | 1.041                               | 1.25     | < 1%         |
| Combined | 0.852                                | 1.102                               | 1.29     | < 1%         |

## 9 Full-exposure results: FHC, RHC, and combined

This is the primary result of the analysis: the CC1 $\mu$ 1 $\pi^\pm$  cross section extracted over the full available exposure in forward horn current (FHC), reverse horn current (RHC), and their combination, all on Poisson-thrown central-value fake data. The FHC measurement uses Runs 1, 2, 4 and 5 (data POT  $8.857 \times 10^{20}$ ); RHC uses Runs 1, 2, 3, 4a, 4b, 4c ( $1.108 \times 10^{21}$ ); the combined measurement uses all fourteen overlay files at the joint POT  $1.994 \times 10^{21}$ . All three close cleanly against the  $A_C$ -smeared truth for every observable (Tables 22–24); the differential results are shown in Figs. 25–27.

Table 22: Full-exposure FHC ({Run 1,2,4,5}, data POT  $8.857 \times 10^{20}$ , native Run-4 FHC detector variations): reco-level and unfolded flux systematic, unfolding amplification, and fake-data closure  $\chi^2/\text{ndf}$  (with  $p$ -value).

| Observable        | Flux (reco) | Flux (unfolded) | Amplification | Closure $\chi^2/\text{ndf}$ |
|-------------------|-------------|-----------------|---------------|-----------------------------|
| $p_\mu$           | 17.1%       | 23.9%           | 1.40×         | 0.55/7 ( $p = 1.00$ )       |
| $p_\pi$           | 16.9%       | 23.9%           | 1.41×         | 0.29/5 ( $p = 1.00$ )       |
| $\cos \theta_\mu$ | 16.7%       | 23.9%           | 1.43×         | 0.93/5 ( $p = 0.97$ )       |
| $\cos \theta_\pi$ | 16.6%       | 27.7%           | 1.67×         | 0.10/4 ( $p = 1.00$ )       |
| $\theta_{\mu\pi}$ | 16.8%       | 22.3%           | 1.33×         | 1.84/5 ( $p = 0.87$ )       |

We extend the measurement to the reverse-horn-current beam as both a standalone measurement and a combined  $\nu_\mu + \bar{\nu}_\mu$  result. The RHC *flux* is  $\bar{\nu}_\mu$ -dominant (54.6%  $\bar{\nu}_\mu$ , versus FHC’s 35%), with authoritative > 60 MeV integrals  $\Phi_{\nu_\mu}^{\text{RHC}} = 2.923 \times 10^{-10}$ ,  $\Phi_{\bar{\nu}_\mu}^{\text{RHC}} = 3.523 \times 10^{-10}$ , total  $6.446 \times 10^{-10}$  cm<sup>2</sup>POT<sup>−1</sup> (the FHC-convention computation reproduces the authoritative FHC values exactly). The *measured event rate*, however, remains  $\nu_\mu$ -dominant (76%  $\nu_\mu$ , 20%  $\bar{\nu}_\mu$ ) because the  $\nu_\mu$  cross section is  $\sim 2\text{--}3\times$  larger than the  $\bar{\nu}_\mu$  one. The RHC overlays (Runs 1, 2, 3, 4a, 4b, 4c) are all verified as genuine  $\nu_\mu$  overlays and processed,  $4.05 \times 10^6$  events, combined MC POT  $1.546 \times 10^{22}$ , mutually consistent ( $2.62 \times 10^{-16}$  events/POT) with the PPF weights, covering the full RHC exposure (Run 3, the largest period at  $5.52 \times 10^{21}$  MC POT /  $1.44 \times 10^6$  events, was added last). Native Run-4 RHC detector-variation samples (the standard eight-knob set: light-yield down/Rayleigh, recombination, space charge, and the four wire-modification variations) are now used for the RHC detector systematic, verified as genuine  $\nu_\mu$  overlays with a uniform  $3.80 \times 10^{-16}$  events/POT across all knobs; they replace the Run-1 FHC stand-in used previously. Native RHC dirt is still taken from FHC. The standalone result below uses

Poisson-thrown central-value fake data at the RHC data POT  $1.108 \times 10^{21}$ . The generator predictions are re-averaged to the RHC flux composition; the combined FHC+RHC predictions are the fluence-weighted combination of the two horn modes (0.458 FHC / 0.542 RHC, each mode weighted by  $\Phi_{\text{tot}} \times \text{POT}$ ).

The combined FHC+RHC measurement uses the summed 14-file response over the full exposure ( $5.6 \times 10^6$  events, combined data POT  $D_{\text{comb}} = 1.994 \times 10^{21} = D_{\text{FHC}} + D_{\text{RHC}}$ ), with each horn mode normalised to its *own* data exposure. This is a genuine subtlety: the detector-variation covariance is absolute (POT-scaled reco counts), so a single POT scale applied to all files would weight the modes by their Monte-Carlo POT (0.62:1 FHC:RHC) rather than by data exposure (0.80:1). Each mode’s numuMC and detVar samples therefore carry a per-mode `summed_pot` (FHC  $2.158 \times 10^{22}$ , RHC  $2.782 \times 10^{22}$ ,  $= S_{\text{mode}} \times D_{\text{comb}}/D_{\text{mode}}$ ) so the framework’s  $D_{\text{comb}}/\text{summed\_pot}$  scaling reduces to  $D_{\text{FHC}}/S_{\text{FHC}}$  and  $D_{\text{RHC}}/S_{\text{RHC}}$ ; the fake data is Poisson-thrown with the matching per-mode rate.

Contrary to an earlier concern, the joint *flux systematic* is correct as built. All fourteen numuMC files carry the *same* 600 PPFX multisim universes, which the universe maker accumulates index-matched across both horn modes in a single chain, so universe  $u$  is the same hadron-production draw in FHC and RHC. Per-universe summing therefore propagates one parameter draw through both beam configurations and lets each mode’s flux response set the result, capturing the physical (not unit) cross-horn correlation automatically; this is the rigorously correct multisim treatment, superior to any hand-built block covariance. The joint result was in fact blocked by a different, purely technical issue: a framework guard that forbade a second detector-variation file of the same type (needed to carry Run-4 FHC *and* Run-4 RHC as `detVarCV`). That guard now accumulates the per-mode detVar samples exactly as it already did for the alternate-CV samples, matching its own in-code TODO. With both in place the combined closure passes for every observable (Table 24), with an unfolded flux systematic of 22–27% that sits between the standalone FHC and RHC values, as expected for the fluence-weighted mix.

The standalone RHC extraction (native Run-4 RHC detector variations) closes cleanly for every observable (Table 23): the  $A_C$ -smeared truth is recovered with  $\chi^2/\text{ndf} < 0.3$  and  $p > 0.94$  throughout. Two features stand out. First, the reco-level flux systematic is  $\sim 15\%$ , genuinely below the FHC  $\sim 17\%$  floor; the  $\bar{\nu}_\mu$ -dominant flux carries a smaller PPFX hadron-production uncertainty. Second, the unfolding amplification ( $1.29\text{--}1.99\times$ ) mirrors FHC, so the unfolded flux term (19.5–28%) is comparable, again data-statistics limited.

Table 23: Standalone RHC ( $\bar{\nu}_\mu$ -dominant) extraction, full exposure  $1.108 \times 10^{21}$  with native Run-4 RHC detector variations: reco-level and unfolded flux systematic, unfolding amplification, and fake-data closure  $\chi^2/\text{ndf}$  (with  $p$ -value).

| Observable        | Flux (reco) | Flux (unfolded) | Amplification | Closure $\chi^2/\text{ndf}$ |
|-------------------|-------------|-----------------|---------------|-----------------------------|
| $p_\mu$           | 15.4%       | 27.5%           | $1.78\times$  | 1.37/7 ( $p = 0.99$ )       |
| $p_\pi$           | 15.8%       | 23.6%           | $1.49\times$  | 0.21/5 ( $p = 1.00$ )       |
| $\cos \theta_\mu$ | 15.0%       | 26.5%           | $1.76\times$  | 0.43/5 ( $p = 0.99$ )       |
| $\cos \theta_\pi$ | 14.9%       | 28.5%           | $1.91\times$  | 1.03/4 ( $p = 0.91$ )       |
| $\theta_{\mu\pi}$ | 15.1%       | 20.4%           | $1.35\times$  | 0.34/5 ( $p = 1.00$ )       |

Table 24: Combined FHC+RHC ( $\nu_\mu + \bar{\nu}_\mu$ ) extraction over the full joint exposure  $D_{\text{comb}} = 1.994 \times 10^{21}$ , per-mode normalised, with native Run-4 FHC+RHC detector variations (accumulated per mode): reco-level and unfolded flux systematic, unfolding amplification, and fake-data closure  $\chi^2/\text{ndf}$  (with  $p$ -value). The unfolded flux term (22–27%) lies between the standalone FHC and RHC values.

| Observable        | Flux (reco) | Flux (unfolded) | Amplification | Closure $\chi^2/\text{ndf}$ |
|-------------------|-------------|-----------------|---------------|-----------------------------|
| $p_\mu$           | 16.2%       | 22.5%           | $1.39\times$  | 0.58/7 ( $p = 1.00$ )       |
| $p_\pi$           | 16.3%       | 23.9%           | $1.47\times$  | 1.27/5 ( $p = 0.94$ )       |
| $\cos\theta_\mu$  | 15.8%       | 25.0%           | $1.58\times$  | 0.28/5 ( $p = 1.00$ )       |
| $\cos\theta_\pi$  | 15.7%       | 27.0%           | $1.72\times$  | 1.19/4 ( $p = 0.88$ )       |
| $\theta_{\mu\pi}$ | 15.9%       | 26.0%           | $1.64\times$  | 0.43/5 ( $p = 0.99$ )       |




Figure 25: Differential cross sections  $d\sigma/dx$  for all five observables on the current Poisson-thrown central-value fake data, FHC full exposure: unfolded fake data (black points),  $A_C$ -smearred fake-data truth (grey dashed), the MicroBooNE tune (black solid), and the four generator predictions (GENIE, GiBUU, NEUT, NuWro). The unfolded points recover the injected truth for every observable, the same closure summarised in Tables 23–24.




Figure 26: Standalone RHC ( $\bar{\nu}_\mu$ -dominant) differential cross sections on the current Poisson-thrown fake data, with the RHC-flux-averaged generator predictions. Same layout as Fig. 25.




Figure 27: Combined FHC+RHC ( $\nu_\mu + \bar{\nu}_\mu$ ) differential cross sections on the current per-mode-normalised fake data, with the fluence-weighted combined generator predictions.

Table 25: Integrated (unfolded) cross section  $\sigma_{\text{int}}$  over the measured phase space, for each observable and configuration (current fake data, coarsened  $\cos\theta_\pi/\theta_{\mu\pi}$  binning, per-configuration flux;  $10^{-38}$  cm<sup>2</sup>/Ar). The value differs between observables because the Wiener-SVD additional-smearing  $A_C$  is not norm-preserving; the D’Agostini cross-check (Sec. 8.14) recovers a flatter,  $\sim 20\%$  higher total, confirming the spread is a smearing effect. All entries were recomputed in a single pass with the current code and a forced rebuild of the cached universes. The MC POT normalization scales each ntuple file by its own stored POT ( $D_{\text{run}}/\text{POT}_{\text{file}}$ ): the processing writes the *total* run MC POT into every file of a run, so for the multi-file RHC runs (Run 3 split across 5 ntuple files, Run 4 across 3) the per-file scaling summed over the splits gives the correct total. An intermediate version of this analysis instead divided by the *sum* of the per-file POT values; because those values are already run totals, that over-counted the denominator by the split multiplicity and scaled the RHC/Combined simulation a factor of 5 (Run 3) and 3 (Run 4) too low, which under-subtracted the background and inflated the RHC/Combined cross sections. That change has been reverted and the RHC/Combined columns here are the corrected values, roughly 10–25% below the intermediate ones; FHC (one MC file per run) is unaffected by either convention. The last two rows give the alternative angular parameterisation  $\theta_\mu, \theta_\pi$  (Sec. 9.2), now complete for all three configurations. (Values are fake-data closures; the signal is  $\nu_\mu + \bar{\nu}_\mu$ , so the RHC flux-average is not expected to lie far below FHC.)

| Observable        | FHC   | RHC   | Combined |
|-------------------|-------|-------|----------|
| $p_\mu$           | 0.956 | 0.717 | 0.736    |
| $p_\pi$           | 0.965 | 0.882 | 0.880    |
| $\cos\theta_\mu$  | 0.966 | 0.757 | 0.636    |
| $\cos\theta_\pi$  | 0.829 | 0.766 | 0.729    |
| $\theta_{\mu\pi}$ | 0.987 | 0.889 | 0.931    |
| $\theta_\mu$      | 0.861 | 0.781 | 0.650    |
| $\theta_\pi$      | 0.878 | 0.739 | 0.748    |

**Uncertainties shown.** Every differential cross section and every integrated value quoted in this note carries *both* statistical and systematic uncertainties. The error bars on the unfolded points are the square root of the diagonal of the *total* covariance matrix, propagated through the unfolding: flux (PPFX multisims), cross-section (GENIE multisims and unisims), detector variations, hadron re-interaction, POT counting and target number, together with the data, MC and EXT statistical terms. For  $p_\mu$  in FHC the bin-averaged total is 32.9% (Table 31), dominated by flux (24.6%), with detector 12.3%, data statistics 11.6% and cross-section 9.2%. The cut-and-count total cross sections of §9.1 quote the same combination, written there as systematic  $\oplus$  data-statistical (FHC: 25.5%  $\oplus$  4.3%).

## 9.1 Total flux-averaged cross section (cut-and-count)

As a check independent of the unfolding, the total flux-averaged cross section is also extracted directly from the selected event counts,

$$\sigma = \frac{N_{\text{sel}} - N_{\text{bkg}}}{\epsilon \Phi N_{\text{Ar}}}, \quad (13)$$

where  $N_{\text{sel}}$  is the number of selected events,  $N_{\text{bkg}}$  the predicted background ( $\nu$ -background + EXT + dirt),  $\epsilon = N_{\text{sel}}^{\text{sig}}/N_{\text{gen}}^{\text{sig}}$  the selection efficiency in the measured phase space ( $\simeq 17.5\%$  for all configurations),  $\Phi$  the integrated flux ( $\Phi = \text{POT} \times \Phi_{\text{POT}}$ , Table 3) and  $N_{\text{Ar}} = 8.71 \times 10^{29}$  the number of argon nuclei in the fiducial volume (the dead  $z$  region removed), giving the same normalisation  $N_{\text{Ar}} \Phi/10^{38}$  used for the differential result. This uses no matrix inversion; by avoiding the Wiener-SVD additional smearing it recovers the true (un-suppressed) total, consistent with the D’Agostini cross-check (Sec. 8.14) and lying above the smearing-suppressed  $\sigma_{\text{int}}$



of Table 25, most visibly for RHC. The uncertainty is the quadrature of the systematic breakdown (Tables 31–33, flux-dominated) and the cut-and-count data-statistical term  $\sqrt{N_{\text{sel}}}/(N_{\text{sel}} - N_{\text{bkg}})$ , which is smaller (3–4%) than the unfolded data-statistical term because no regularisation amplifies it.

## 9.2 Angular variable: $\cos\theta_\mu$ versus $\theta_\mu$

The muon-angle distribution is measured in both  $\cos\theta_\mu$  and  $\theta_\mu$ . The two carry identical information; the  $\theta_\mu$  bin edges are exactly arccos of the  $\cos\theta_\mu$  edges, so the phase space is partitioned identically, but they behave very differently under the Wiener–SVD additional smearing  $A_C$ , and the choice matters for how the measurement compares to models.

In  $\cos\theta_\mu$  the signal is sharply forward-peaked: the last two bins ( $\cos\theta_\mu > 0.8$ ) hold most of the rate, and  $A_C$  suppresses the integral of any truth-level prediction by a factor  $\sim 0.33$ , against  $\sim 0.6$  for the momentum observables. The four generators are pushed  $2\text{--}3\times$  below the tune and the data even though their truth-level total cross sections agree to within tens of percent.

An earlier version of this note attributed both effects to the sharpness of the peak,  $A_C$  smearing away an unresolvable feature, and thereby penalising the peaked generators more than the smoother tune. *Both halves of that explanation are wrong*, and are retracted here. Across the full set of observables the  $A_C$  integral factor shows no correlation with peakedness ( $r = -0.17$  on the pre-fix factors):  $W_{\pi p}$  is the sharpest distribution in the analysis (peak-to-minimum ratio 36) yet was suppressed only to 0.78,  $\delta p_T$  is comparatively flat (ratio 10) yet was suppressed to 0.43, and  $p_n$  is flat yet was *enhanced* to 1.71; an enhancement no peak argument can produce. The retraction stands, and the width-aware  $A_C$  makes the point moot: the factors are now uniform across observables regardless of shape. Within  $\cos\theta_\mu$  itself the ordering is backwards: NEUT and NuWro are the *flattest* of the five truth-level curves (peak-to-minimum 3.8 and 4.3, against 16.3 for the tune) and are suppressed the *most* (0.37 and 0.29, against 0.67). What actually drives their suppression is that  $A_C$  pushes their backward bins negative ( $-0.0005$  to  $-0.0212$  in normalised units), eating the integral.  $A_C$  is the Wiener filter acting on the SVD modes of the response: it removes whatever the data cannot statistically constrain, which is set by the conditioning of the response for that observable and binning, not by the shape of the truth distribution, and it is not norm-preserving, so it can suppress or enhance.

### The FHC $\theta_\mu$ collapse, its cause, and its resolution

Reparameterising to  $\theta_\mu$  appeared to remove the problem entirely: the FHC suppression rose to  $\sim 0.76$  and the NEUT/NuWro  $\chi^2$  fell from  $\sim 26/5$  to  $\sim 2.8/5$ , with the generators sitting on top of the data. *That result was an artefact*, and it is worth setting out what the artefact was, because the cause turned out not to be  $\theta_\mu$  at all.

In the FHC  $\theta_\mu$  extraction the unfolded data, the  $A_C$ -smeared truth, the MicroBooNE tune and all four generators shared the same normalised shape to four decimal places (Table 26).  $A_C$  was *rank-1*: writing the four generator predictions as the columns of an input matrix and their  $A_C$ -smeared counterparts as the columns of the output matrix, the inputs were full rank ( $s_2/s_1 = 5.0 \times 10^{-2}$ ) while the outputs gave  $s_2/s_1 = 2.8 \times 10^{-6}$ . Every shape degree of freedom was annihilated, each model was mapped onto one common curve and only its normalisation survived. The agreement with the data was true by construction.

**The cause: the regularisation matrix ignored the bin widths.** Wiener–SVD regularises with a matrix  $C$ ; this analysis uses the second-derivative form. It was built as the plain tridiagonal stencil  $(1, -2, 1)$ , which is a second derivative only on a *uniform* grid. Our binnings are strongly non-uniform, with width ratios up to 14:1 (Table 27), so the operator was not a second derivative at all. It was near-singular, its row sums were exactly the  $10^{-6}$  regularisation floor, and its near-null direction was the constant vector rather than anything physical. The Wiener filter was



Table 26: FHC  $\theta_\mu$  before the fix described below: normalised bin contents. Five independent inputs, the measurement and four generator predictions plus the tune, collapse onto one shape, the signature of a rank-1  $A_C$ .

|                      | bin 1  | bin 2  | bin 3  | bin 4  | bin 5  |
|----------------------|--------|--------|--------|--------|--------|
| Unfolded data        | 0.1485 | 0.3361 | 0.2899 | 0.1934 | 0.0321 |
| $A_C$ -smeared truth | 0.1484 | 0.3359 | 0.2901 | 0.1933 | 0.0323 |
| MicroBooNE tune      | 0.1484 | 0.3359 | 0.2901 | 0.1934 | 0.0322 |
| NEUT                 | 0.1484 | 0.3359 | 0.2901 | 0.1934 | 0.0322 |

then free to retain that direction alone, which is precisely a rank-1  $A_C$  that passes normalisation and discards shape.

That also explains why  $\theta_\mu$  was the worst case rather than a random one: its bin widths span a factor of 10.4, and  $\cos\theta_\mu$ , the next most non-uniform at 14.5, was the other observable whose  $\chi^2$  this note previously attributed to an  $A_C$  artefact. The remaining observables sit at ratios of 3–6 and were far less affected.

**The fix.** Both derivative regularisation matrices now difference the density on the actual grid: each column is divided by its bin width and the stencil uses the true spacings between bin centres, with a one-sided first-derivative stencil at the ends. (A Dirichlet closure was rejected: it asserts the distribution vanishes at the edges, which is false here,  $\cos\theta_\mu$  peaks at its upper edge.) With no widths supplied the code reduces exactly to the previous behaviour. The widths are passed in by the unfolding application, which holds the slice definitions that the cross-section extractor does not.

The effect on the conditioning of  $A_C$ , measured as  $s_2/s_1$  of the smearing matrix itself, is given in Table 27. Every inclusive observable improves and none is degraded except combined  $\cos\theta_\mu$ , marginally.

Table 27:  $A_C$  conditioning  $s_2/s_1$  before and after the regularisation fix, with the bin-width ratio of each observable. A value approaching zero means the smearing matrix has collapsed onto a single mode and the extraction carries no shape information.

| observable                                                                                         | width<br>ratio | FHC          |             | RHC    |       | COMB                                       |
|----------------------------------------------------------------------------------------------------|----------------|--------------|-------------|--------|-------|--------------------------------------------|
|                                                                                                    |                | before       | after       | before | after | before $\rightarrow$ after                 |
| $p_\mu$                                                                                            | 6.3            | 0.34         | 0.90        | 0.31   | 0.94  | 0.19 $\rightarrow$ 0.68                    |
| $\cos\theta_\mu$                                                                                   | 14.5           | 0.46         | 0.57        | 0.45   | 0.55  | 0.42 $\rightarrow$ 0.39                    |
| $\cos\theta_\pi$                                                                                   | 3.6            | 0.67         | 0.71        | 0.55   | 0.79  | 0.55 $\rightarrow$ 0.74                    |
| $\theta_{\mu\pi}$                                                                                  | 3.0            | 0.42         | 0.85        | 0.47   | 0.65  | 0.42 $\rightarrow$ 0.46                    |
| $\theta_\mu$                                                                                       | 10.4           | <b>0.015</b> | <b>0.51</b> | 0.22   | 0.62  | <b>0.045 <math>\rightarrow</math> 0.38</b> |
| $p_\pi$ (2 bins)                                                                                   | —              | 0.46         | 0.53        | 0.41   | 0.44  | 0.29 $\rightarrow$ 0.40                    |
| <i>proton-tagged; only the combined configuration had <math>A_C</math> recorded before the fix</i> |                |              |             |        |       |                                            |
| $W_{\pi p}$                                                                                        | —              |              |             |        |       | <b>0.022 <math>\rightarrow</math> 0.60</b> |
| $W_{\text{had}}$                                                                                   | —              |              |             |        |       | 0.23 $\rightarrow$ 0.86                    |
| $\delta p_T$                                                                                       | —              |              |             |        |       | 0.25 $\rightarrow$ 0.93                    |
| $\delta\alpha_T$                                                                                   | —              |              |             |        |       | 0.24 $\rightarrow$ 0.48                    |
| $\delta\phi_T$                                                                                     | —              |              |             |        |       | <b>0.083 <math>\rightarrow</math> 0.85</b> |
| $p_n$                                                                                              | —              |              |             |        |       | 0.21 $\rightarrow$ 0.88                    |

**$\theta_\mu$  is reinstated, and the validation is decisive.** The fake data are Poisson throws from the MicroBooNE tune, so the tune is the model that *ought* to fit best. Before the fix it did not

stand out, because every model had been mapped onto the same curve; after it, it does:

| FHC $\theta_\mu$ , $\chi^2/5$ bins | before (rank-1)     | after                             |
|------------------------------------|---------------------|-----------------------------------|
| MicroBooNE tune                    | 3.07 ( $p = 0.69$ ) | <b>5.14</b> ( $p = 0.40$ )        |
| GENIE                              | 4.73 ( $p = 0.45$ ) | 11.3 ( $p = 0.046$ )              |
| GiBUU                              | 3.03 ( $p = 0.70$ ) | 17.9 ( $p = 0.003$ )              |
| NuWro                              | 2.90 ( $p = 0.72$ ) | 38.6 ( $p = 2.8 \times 10^{-7}$ ) |
| NEUT                               | 2.82 ( $p = 0.73$ ) | 40.6 ( $p = 1.1 \times 10^{-7}$ ) |

The closure itself is unaffected and remains good,  $\chi^2 = 2.17/5$  ( $p = 0.83$ ). The  $\theta_\mu$  measurement is therefore no longer withdrawn: it carries genuine discriminating power, and on this fake data it correctly identifies the model the data were generated from. What must be withdrawn instead is the earlier *explanation* that  $\theta_\mu$  “fixes” the  $\cos\theta_\mu$  tension; both parameterisations are now properly conditioned and should be read together.

**A correction to the survey below.** The survey of eighteen extractions reported that FHC  $\theta_\mu$  was the only rank-deficient case. That was wrong in two ways. Combined  $\theta_\mu$  was also badly degraded at  $s_2/s_1 = 0.045$  and was not identified as such; and the survey covered only the inclusive extractions, so the proton-tagged channels were never examined. They should have been: combined  $W_{\pi p}$  sat at 0.022, as compromised as FHC  $\theta_\mu$ , and combined  $\delta\phi_T$  at 0.083. All are repaired by the same fix.

### Is any other observable affected? A survey of all eighteen extractions

The rank-1 failure was found by inspection, so the unfolders was modified to write the slice-restricted  $A_C$  into the closure file (`h_A_C`; it had previously only been drawn to a canvas, and the attempt to save it failed silently with “the current directory is not associated with a file”). All eighteen inclusive extractions, six observables  $\times$  three beam configurations, were then re-unfolded and their  $A_C$  examined directly. Table 28 reports, for each, how much of the model discrimination present in the truth-level predictions survives the smearing: the largest pairwise difference between normalised generator shapes before and after  $A_C$ , and their ratio.

*The survey that follows was made before the regularisation fix and its conclusion of uniqueness is superseded; see the correction above.* As measured then, the collapse looked unique to FHC  $\theta_\mu$ : it was the only entry with zero retention, and the only  $A_C$  whose rows are all the same vector (spread 9% across rows, effective rank 1.02, where every other extraction has row-to-row spreads of 170–1050% and effective ranks 1.24–1.91). RHC and combined  $\theta_\mu$  are not affected; the combined case is the weaker of the two (12% retention, effective rank 1.06) and its  $\theta_\mu$  model comparison should be read as a soft test, but it is not degenerate.

Two broader limitations are visible in the same table and should be stated plainly rather than left implicit.

First, the momentum observables retain only 11–21% of their truth-level model discrimination in every beam configuration.  $A_C$  does not destroy the shape information there, but it removes most of it, so a good  $\chi^2$  against  $p_\mu$  or  $p_\pi$  is a weak statement about a model. The angular observables are better:  $\theta_{\mu\pi}$  retains 58–93% and is the healthiest variable in the analysis.

Second, the apparent 120–919% retention in  $\cos\theta_\mu$  is not the good news it appears to be. It arises because  $A_C$  drives the generator predictions to *negative* differential cross sections in the backward bins while leaving the tune positive, for the combined configuration the smeared predictions run from  $-0.43$  (NuWro) to  $+0.51$  (the tune) in the second bin. A negative differential cross section is unphysical, so the large spread reflects a regularisation artefact rather than genuine model separation, and any  $\chi^2$  computed against those curves inherits the problem. This is the same effect that produces the strong  $\cos\theta_\mu$  integral suppression discussed above.

Table 28: Retention of model discrimination through  $A_C$ . “Before” and “after” are the largest pairwise difference between normalised generator shapes, truth-level and  $A_C$ -smeared. A ratio near zero means the predictions have been collapsed onto a common curve and can no longer be told apart by the measurement; a ratio above unity means  $A_C$  has amplified the spread, which for  $\cos \theta_\mu$  happens through unphysical negative excursions rather than genuine discrimination (see text).

| Beam | Observable        | before | after                | retention |
|------|-------------------|--------|----------------------|-----------|
| FHC  | $p_\mu$           | 0.0248 | 0.0052               | 21%       |
| FHC  | $p_\pi$           | 0.0586 | 0.0080               | 14%       |
| FHC  | $\cos \theta_\mu$ | 0.0554 | 0.0724               | 131%      |
| FHC  | $\cos \theta_\pi$ | 0.0589 | 0.0211               | 36%       |
| FHC  | $\theta_{\mu\pi}$ | 0.0422 | 0.0391               | 93%       |
| FHC  | $\theta_\mu$      | 0.0554 | $2.5 \times 10^{-6}$ | <b>0%</b> |
| RHC  | $p_\mu$           | 0.0232 | 0.0036               | 15%       |
| RHC  | $p_\pi$           | 0.0556 | 0.0068               | 12%       |
| RHC  | $\cos \theta_\mu$ | 0.0644 | 0.0771               | 120%      |
| RHC  | $\cos \theta_\pi$ | 0.0629 | 0.0165               | 26%       |
| RHC  | $\theta_{\mu\pi}$ | 0.0505 | 0.0295               | 58%       |
| RHC  | $\theta_\mu$      | 0.0644 | 0.0188               | 29%       |
| COMB | $p_\mu$           | 0.0240 | 0.0028               | 12%       |
| COMB | $p_\pi$           | 0.0571 | 0.0064               | 11%       |
| COMB | $\cos \theta_\mu$ | 0.0598 | 0.5500               | 919%      |
| COMB | $\cos \theta_\pi$ | 0.0609 | 0.0166               | 27%       |
| COMB | $\theta_{\mu\pi}$ | 0.0463 | 0.0360               | 78%       |
| COMB | $\theta_\mu$      | 0.0598 | 0.0070               | 12%       |

Taken together, the earlier framing of this section had the two muon-angle variables the wrong way round.  $\cos\theta_\mu$  was described as the variable  $A_C$  ruins and  $\theta_\mu$  as the fix; in fact FHC  $\theta_\mu$  is the one extraction in which the measurement retains no shape information at all, while  $\cos\theta_\mu$  retains its shape modes but at the cost of unphysical negative excursions. Neither is a clean muon-angle measurement, and both carry the caveats recorded here.

Table 29: Integrated cross section ( $10^{-38} \text{ cm}^2/\text{Ar}$ ) for the unfolded fake data, the MicroBooNE tune and the four generators, in  $\cos\theta_\mu$  and in  $\theta_\mu$ . All entries are  $A_C$ -smeared, i.e. live in the same space as the measurement. “ $A_C$  supp.” is the mean ratio of the  $A_C$ -smeared to the truth-level generator integral; the momentum observable  $p_\mu$  is shown for reference. **The FHC  $\theta_\mu$  row is retained for the record only and must not be used for model comparison:**  $A_C$  is rank-1 for that config, so every entry in the row is the same shape rescaled and the apparent agreement is by construction (Table 26).

| Beam             | Variable         | data             | tune      | GENIE        | GiBUU     | NEUT                | NuWro |
|------------------|------------------|------------------|-----------|--------------|-----------|---------------------|-------|
| FHC              | $\cos\theta_\mu$ | 0.819            | 0.673     | 0.274        | 0.390     | 0.367               | 0.295 |
| FHC              | $\theta_\mu$     | 0.996            | 0.871     | 0.625        | 0.876     | 1.025               | 0.919 |
| RHC              | $\cos\theta_\mu$ | 0.662            | 0.633     | 0.226        | 0.338     | 0.320               | 0.256 |
| RHC              | $\theta_\mu$     | 0.828            | 0.824     | 0.328        | 0.487     | 0.505               | 0.423 |
| $A_C$ supp. FHC: |                  | $\cos\theta_\mu$ | 0.23–0.34 | $\theta_\mu$ | 0.73–0.78 | $(p_\mu$ 0.59–0.67) |       |
| $A_C$ supp. RHC: |                  | $\cos\theta_\mu$ | 0.25–0.36 | $\theta_\mu$ | 0.41–0.52 | $(p_\mu$ 0.69–0.75) |       |




Figure 28: Muon-angle differential cross section in  $\theta_\mu$  (left) and  $\cos\theta_\mu$  (right), for FHC (top) and RHC (bottom). Unfolded fake data (black points), the MicroBooNE tune (black line) and the four generators are all  $A_C$ -smeared. In  $\cos\theta_\mu$  the predictions are strongly suppressed by  $A_C$  and fall well below both the data and the tune; the most suppressed models are those whose backward bins  $A_C$  drives negative. **The FHC  $\theta_\mu$  panel in this figure predates the regularisation fix:**  $A_C$  was rank-1 there, so the data and every prediction are the same curve up to normalisation and their apparent agreement carries no shape information. With the width-aware regularisation  $\theta_\mu$  discriminates normally (Sec. 9.2).

Table 30 compares the result with the four generator predictions, each flux-averaged over the corresponding beam. Because the analysis is blind the “measurement” is here the central-value overlay (the MicroBooNE tune) that the fake data is thrown from; at unblinding  $N_{\text{sel}}$  becomes the real beam-on count. It agrees with all four generators within  $\sim 1\sigma$ : standalone GENIE sits about  $1\sigma$  low, while GiBUU, NEUT and NuWro lie within half a sigma. RHC and the combined total carry the larger uncertainty, driven by the RHC detector systematic.

Table 30: Total flux-averaged cross section from the cut-and-count method (Eq. 13;  $10^{-38} \text{ cm}^2/\text{Ar}$ ), compared with the flux-averaged generator predictions. The measurement uncertainty is the systematic breakdown (flux/detector/cross-section/reinteraction, Tables 31–33)  $\oplus$  cut-and-count data statistics. Values in parentheses give each generator’s deviation from the measurement in units of the measurement uncertainty. The generator predictions are integrated over the full binning; earlier versions of this table used truncated binnings ( $p_\mu$  closed at 3 GeV/ $c$ ,  $p_\pi$  at 1 GeV/ $c$ ) that discarded the tails and made every prediction 4–8% low (§8.8).

| Config   | Measurement     | GENIE                | GiBUU                | NEUT                 | NuWro                |
|----------|-----------------|----------------------|----------------------|----------------------|----------------------|
| FHC      | $1.06 \pm 0.27$ | 0.82 (0.9 $\sigma$ ) | 1.13 (0.3 $\sigma$ ) | 1.32 (1.0 $\sigma$ ) | 1.25 (0.7 $\sigma$ ) |
| RHC      | $0.98 \pm 0.35$ | 0.67 (0.9 $\sigma$ ) | 0.94 (0.1 $\sigma$ ) | 1.11 (0.3 $\sigma$ ) | 1.04 (0.2 $\sigma$ ) |
| Combined | $1.02 \pm 0.30$ | 0.74 (0.9 $\sigma$ ) | 1.03 (0.0 $\sigma$ ) | 1.20 (0.6 $\sigma$ ) | 1.14 (0.4 $\sigma$ ) |

The systematic breakdown by source (bin-averaged fractional uncertainty in cross-section space) is given for all three configurations in Tables 31–33. The flux (PPFX) term dominates throughout, followed by the detector variations; the data-statistical term reflects the fake-data POT of each configuration. The RHC detector term is the largest single-configuration entry, reaching 20.5% on  $p_\pi$ ; it is no longer the  $\cos\theta$  observables that drive it.

Table 31: FHC ({Run 1,2,4,5}) systematic breakdown by source, bin-averaged fractional uncertainty (%).

| Source                | $p_\mu$     | $p_\pi$     | $\cos\theta_\mu$ | $\cos\theta_\pi$ | $\theta_{\mu\pi}$ | $\theta_\mu$ |
|-----------------------|-------------|-------------|------------------|------------------|-------------------|--------------|
| <b>Total</b>          | <b>32.9</b> | <b>38.4</b> | <b>30.8</b>      | <b>33.0</b>      | <b>29.7</b>       | <b>31.4</b>  |
| Cross section (GENIE) | 9.2         | 13.4        | 8.2              | 10.8             | 9.1               | 8.1          |
| Flux (PPFX)           | 24.6        | 26.6        | 25.7             | 26.4             | 25.4              | 25.4         |
| Detector              | 12.3        | 10.9        | 9.6              | 10.1             | 8.0               | 10.4         |
| Reinteraction         | 3.9         | 9.5         | 3.3              | 3.9              | 3.4               | 3.3          |
| MC stat               | 3.8         | 5.1         | 3.0              | 3.4              | 2.4               | 3.5          |
| EXT stat              | 2.8         | 5.1         | 2.9              | 3.8              | 2.3               | 3.4          |
| Data stat             | 11.6        | 15.4        | 9.1              | 10.3             | 7.3               | 10.6         |
| POT + targets         | 2.8         | 3.1         | 2.9              | 3.0              | 2.8               | 2.8          |

Table 32: RHC ({Run 1,2,3,4a,4b,4c}) systematic breakdown by source, bin-averaged fractional uncertainty (%).

| Source                | $p_\mu$     | $p_\pi$     | $\cos\theta_\mu$ | $\cos\theta_\pi$ | $\theta_{\mu\pi}$ | $\theta_\mu$ |
|-----------------------|-------------|-------------|------------------|------------------|-------------------|--------------|
| <b>Total</b>          | <b>38.1</b> | <b>42.8</b> | <b>33.9</b>      | <b>32.1</b>      | <b>29.9</b>       | <b>29.9</b>  |
| Cross section (GENIE) | 10.7        | 14.7        | 9.6              | 9.6              | 9.4               | 8.6          |
| Flux (PPFX)           | 27.1        | 27.4        | 26.1             | 24.6             | 24.1              | 23.4         |
| Detector              | 17.2        | 20.5        | 13.7             | 12.7             | 9.7               | 11.3         |
| Reinteraction         | 4.2         | 9.4         | 3.8              | 3.6              | 3.5               | 3.4          |
| MC stat               | 3.7         | 4.1         | 3.0              | 2.8              | 2.4               | 2.6          |
| EXT stat              | 4.0         | 5.3         | 3.7              | 3.8              | 2.9               | 3.3          |
| Data stat             | 13.7        | 15.2        | 11.1             | 10.2             | 8.8               | 9.7          |
| POT + targets         | 3.3         | 3.5         | 3.3              | 3.0              | 3.0               | 2.9          |

Table 33: Combined FHC+RHC systematic breakdown by source, bin-averaged fractional uncertainty (%).

| Source                | $p_\mu$     | $p_\pi$     | $\cos\theta_\mu$ | $\cos\theta_\pi$ | $\theta_{\mu\pi}$ | $\theta_\mu$ |
|-----------------------|-------------|-------------|------------------|------------------|-------------------|--------------|
| <b>Total</b>          | <b>42.9</b> | <b>44.4</b> | <b>41.6</b>      | <b>36.4</b>      | <b>32.4</b>       | <b>37.0</b>  |
| Cross section (GENIE) | 11.6        | 13.7        | 9.2              | 10.1             | 9.7               | 8.9          |
| Flux (PPFX)           | 27.4        | 26.7        | 29.1             | 25.5             | 25.3              | 26.5         |
| Detector              | 24.6        | 27.9        | 23.0             | 21.3             | 15.4              | 20.3         |
| Reinteraction         | 4.7         | 9.4         | 4.0              | 3.6              | 3.3               | 3.8          |
| MC stat               | 4.4         | 3.3         | 4.1              | 2.3              | 2.0               | 3.1          |
| EXT stat              | 3.8         | 3.7         | 4.4              | 2.7              | 1.7               | 3.3          |
| Data stat             | 14.4        | 10.8        | 13.7             | 7.5              | 6.5               | 10.3         |
| POT + targets         | 3.2         | 3.3         | 3.5              | 3.0              | 3.0               | 3.1          |

### 9.3 Background-control sidebands

Because the analysis is blind, the background model cannot yet be confronted with beam-on data in the signal region. To exercise it before unblinding, four signal-depleted control regions (“sidebands”) are defined by inverting exactly one signal requirement, leaving the rest of the selection intact:

- **CC0 $\pi$** : the muon selection with no reconstructed charged pion (charged-current quasi-elastic /  $2p2h$ -enriched);
- **multi- $\pi$** : two or more charged pions (deep-inelastic / resonant multi-pion);
- **$\pi^0$** : a reconstructed shower present but the  $\pi^0$  shower veto failed (CC  $\pi^0$  / NC  $\pi^0$ );
- **cosmic**: the full signal selection with the muon–pion opening angle inverted ( $\theta_{\mu\pi} > 2.6$ ), enriched in out-of-time cosmic activity.

Each region is evaluated with the same per-run POT scaling and CV event weighting (§3) used for the measurement. Because the current “data” is a Poisson throw of the neutrino Monte Carlo central value only (it contains no EXT or dirt component), the meaningful closure quantity here is the ratio of the pseudo-data to the CV-weighted *neutrino* MC,  $N_{\text{data}}/N_{\nu\text{MC}}$ ; the EXT and dirt yields are shown for scale but enter the background-model test only once real beam-on data are available. Table 34 summarises the four regions for the three beam configurations.

Table 34: Background-control sidebands. “Signal fraction” is the signal contamination of the (CV-weighted neutrino) MC in the region; all four are strongly signal-depleted.  $N_{\text{data}}/N_{\nu\text{MC}}$  is the pseudo-data to neutrino-MC ratio: the high-statistics regions (CC0 $\pi$ ,  $\pi^0$ ) close to unity across FHC/RHC/combined, validating the per-run weighting and POT-scaling machinery; the low-statistics regions (multi- $\pi$ , cosmic) scatter within Poisson expectation ( $\sqrt{N} \approx 6\text{--}10\%$ ). The background-model test against the full stack (with EXT and dirt) is performed at control-region unblinding with beam-on data.

| Sideband     | Signal frac.<br>(of $\nu\text{MC}$ ) | $N_{\text{data}}/N_{\nu\text{MC}}$ |      |          |
|--------------|--------------------------------------|------------------------------------|------|----------|
|              |                                      | FHC                                | RHC  | Combined |
| CC0 $\pi$    | $\sim 10\%$                          | 1.00                               | 1.00 | 1.00     |
| multi- $\pi$ | $\sim 23\%$                          | 1.03                               | 1.11 | 1.08     |
| $\pi^0$      | $\sim 10\%$                          | 0.99                               | 1.02 | 1.01     |
| cosmic       | $\sim 8\%$                           | 0.94                               | 1.24 | 1.10     |

**Yields and composition.** Table 35 gives the full breakdown for the three configurations. The regions differ enormously in size and in what they are sensitive to, which is what makes them complementary rather than redundant:

- **CC0 $\pi$**  is by far the largest ( $\sim 12\text{k}$  pseudo-data events per beam,  $\sim 24\text{k}$  combined). Its neutrino component is dominated by the quasi-elastic and  $2p2h$  modes that feed the signal region only through pion *absorption* in the final state. It is therefore the natural handle on the FSI model that converts CC  $1\pi^\pm$  into CC0 $\pi$  and vice versa; the largest single migration path into and out of the signal definition (§4).
- **$\pi^0$**  is the second largest (6.5–14.7k). It is enriched in CC/NC  $\pi^0$  events, whose showers are precisely what the shower veto (Cut 8) is designed to remove; it therefore tests the veto efficiency and the  $\pi^0$  production normalisation together. These are the events that leak into the signal region when a photon is missed.

- **multi- $\pi$**  is small (127–303 events) but has the highest signal contamination of the four ( $\sim 23\%$ ), because a true multi-pion event with one pion lost is exactly the wrong-topology background the selection fights. Its size is what limits its constraining power at present.
- **cosmic** is the only region where the beam-off (EXT) contribution *dominates* the prediction: 308 EXT against 68 neutrino-MC events for FHC, a ratio of 4.5, against 0.8 in CC0 $\pi$ . It is consequently the region that constrains the cosmic normalisation, and the one whose interpretation changes most at unblinding, since the current pseudo-data contain no EXT component at all.

**What the present numbers do and do not show.** With Poisson-thrown neutrino-MC pseudo-data, the ratios of Table 34 test the *machinery*, per-run POT scaling, CV weighting, the selection logic in each inverted region, and not the background model. That test passes: the two high-statistics regions sit at  $N_{\text{data}}/N_{\nu\text{MC}} = 1.00\text{--}1.02$  in all three configurations, which is a non-trivial check of the per-run normalisation that caused the difficulties described in §B. The apparent deviations in the small regions are statistical: with 69 cosmic-sideband events for FHC, a  $\sqrt{N}$  fluctuation is 12%, so the observed 0.94 and the RHC 1.24 (on 100 events, 10%) are within about one and two standard deviations respectively.

**Use at unblinding.** Once the control regions are opened with beam-on data the same table becomes a genuine background test, and the statistical precision available is set by the yields above: the CC0 $\pi$  and  $\pi^0$  regions would constrain their dominant background normalisations to roughly 1% and 1.5% statistically for the combined exposure, well below the  $\sim 20\text{--}30\%$  cross-section systematic they are intended to test. The cosmic region would constrain the EXT normalisation to  $\sim 8\%$ , comparable to the 3–5% trigger-ratio uncertainty already assigned. The multi- $\pi$  region, at  $\sim 300$  combined events, gives  $\sim 6\%$ , useful as a cross-check but not competitive with the others. The intended procedure is to unblind the control regions first, use them to constrain or reweight the dominant background components, and only then open the signal region.

Table 35: Sideband yields at full exposure. “Signal” is the signal contamination of the neutrino MC, “ $\nu$  bkg” the remaining neutrino MC, EXT the beam-off (cosmic) sample scaled by the trigger ratio, and dirt the out-of-cryostat overlay. The pseudo-data column is a Poisson throw of the neutrino MC only, so it should be compared with Signal +  $\nu$  bkg; EXT and dirt are listed for scale and enter the comparison only with real beam-on data.

| Beam | Sideband     | Pseudo-data | Signal | $\nu$ bkg | EXT    | Dirt | Sig. frac. |
|------|--------------|-------------|--------|-----------|--------|------|------------|
| FHC  | CC0 $\pi$    | 11 887      | 1 122  | 10 736    | 8 939  | 54   | 9.5%       |
| FHC  | multi- $\pi$ | 127         | 30     | 94        | 42     | 0.1  | 24.0%      |
| FHC  | $\pi^0$      | 6 555       | 661    | 5 931     | 5 148  | 12   | 10.0%      |
| FHC  | cosmic       | 69          | 6      | 68        | 308    | 2.2  | 8.1%       |
| RHC  | CC0 $\pi$    | 12 261      | 1 204  | 11 034    | 9 281  | 42   | 9.8%       |
| RHC  | multi- $\pi$ | 176         | 35     | 124       | 43     | 0.1  | 22.0%      |
| RHC  | $\pi^0$      | 8 139       | 778    | 7 234     | 5 346  | 9    | 9.7%       |
| RHC  | cosmic       | 100         | 6      | 74        | 320    | 1.7  | 8.0%       |
| Comb | CC0 $\pi$    | 24 148      | 2 327  | 21 770    | 18 219 | 54   | 9.7%       |
| Comb | multi- $\pi$ | 303         | 65     | 217       | 85     | 0.1  | 22.9%      |
| Comb | $\pi^0$      | 14 694      | 1 439  | 13 165    | 10 494 | 12   | 9.9%       |
| Comb | cosmic       | 169         | 12     | 142       | 629    | 2.2  | 8.0%       |



## 9.4 Control regions for the proton-tagged subsample

The proton-tagged selection inherits its cut logic from `CC1mu1piXp`, and with it the four sideband definitions of §9.3: the reprocessed proton-tagged files already carry `CC1mu1pi1p_sb_cc0pi`, `_sb_multi`, `_sb_pi0` and `_sb_cosmic`. No new definitions or reprocessing are required to exercise the background model of the W/TKI measurement; the same four regions can be evaluated either as inherited (the inclusive phase space) or with a reconstructed tagged proton additionally required, which places them in the same phase space as the measurement itself. Table 36 gives both.

Three things follow from the numbers. First, all four regions are *more* signal-depleted for the proton-tagged signal than they were for the inclusive one: 6.0–7.3% signal contamination in  $CC0\pi$  and  $\pi^0$ , against 9.5–10.0% for the same regions in the inclusive phase space and a 48% purity in the signal region, because requiring a proton does not make a pion-less or  $\pi^0$  event look like signal. Second, the proton requirement costs about a third of the statistics ( $CC0\pi$ : 11 887  $\rightarrow$  8 268 for FHC) but suppresses the beam-off contribution by roughly a factor six (8 760  $\rightarrow$  1 418): cosmic-induced events rarely contain a reconstructed proton above threshold, so the proton tag is itself a strong cosmic rejector. This is worth noting for the systematic budget; the EXT normalisation matters far less in the proton-tagged measurement than in the inclusive one. Third, the pseudo-data to neutrino-MC ratios remain consistent with unity in every region and configuration (0.956–1.158 with the tag), so the per-run normalisation machinery is validated in the proton-tagged phase space as well.

**What these regions cannot do.** All four *apply* the proton tag rather than invert it, so none of them constrains the proton *mis-tag* background; the  $\sim 13\%$  of tagged protons that are really a pion or another track, which smears  $W_{\pi p}$  and the TKI variables directly. Testing that requires a region defined by inverting the proton PID (selecting candidates that pass the momentum and containment requirements but fail the LLR cut), which does need a new branch in `CC1mu1pi1p` and a reprocess. Slicing a measured W/TKI observable is *not* an alternative: because the signal definition is inclusive in those variables, cutting on them splits the signal region rather than depleting it. Explicitly, on the Run-1 FHC overlay the signal region has 53.7% purity and the slices  $W_{\pi p} > 1.5$  GeV,  $W_{\pi p} < 1.15$  GeV,  $p_n > 0.8$  GeV/ $c$  and  $\delta p_T > 0.7$  GeV/ $c$  give 50.0%, 48.9%, 47.3% and 52.0% respectively; no depletion at all. Even the kinematically forbidden region  $W_{\text{had}} < m_\pi + m_p$  retains 53.4% signal, because the calorimetric mass is systematically under-reconstructed and true signal populates it through resolution.



Table 36: Sideband yields for the proton-tagged subsample at full exposure, evaluated with the inherited definitions both as-is and with a tagged proton additionally required. “Sig.” is the proton-tagged signal contamination of the neutrino MC.  $N_{\text{data}}/N_{\nu\text{MC}}$  compares the Poisson-thrown pseudo-data with the CV-weighted neutrino MC; EXT and dirt are listed for scale.

| Beam                                                           | Region       | $N_{\text{data}}$ | Signal | $\nu$ bkg | EXT    | Sig. frac. | $N_{\text{data}}/N_{\nu\text{MC}}$ |
|----------------------------------------------------------------|--------------|-------------------|--------|-----------|--------|------------|------------------------------------|
| <i>inherited definitions (inclusive phase space)</i>           |              |                   |        |           |        |            |                                    |
| FHC                                                            | CC0 $\pi$    | 11 887            | 1 122  | 10 736    | 8 760  | 9.5%       | 1.002                              |
| FHC                                                            | multi- $\pi$ | 127               | 30     | 94        | 41     | 24.0%      | 1.030                              |
| FHC                                                            | $\pi^0$      | 6 555             | 660    | 5 931     | 5 045  | 10.0%      | 0.994                              |
| FHC                                                            | cosmic       | 69                | 6.0    | 68        | 302    | 8.1%       | 0.937                              |
| RHC                                                            | CC0 $\pi$    | 12 261            | 1 204  | 11 034    | 9 095  | 9.8%       | 1.002                              |
| RHC                                                            | multi- $\pi$ | 176               | 35     | 124       | 42     | 22.0%      | 1.109                              |
| RHC                                                            | $\pi^0$      | 8 139             | 778    | 7 234     | 5 239  | 9.7%       | 1.016                              |
| RHC                                                            | cosmic       | 100               | 6.4    | 74        | 314    | 8.0%       | 1.239                              |
| Comb                                                           | CC0 $\pi$    | 24 148            | 2 326  | 21 770    | 17 855 | 9.7%       | 1.002                              |
| Comb                                                           | multi- $\pi$ | 303               | 64     | 217       | 83     | 22.9%      | 1.075                              |
| Comb                                                           | $\pi^0$      | 14 694            | 1 439  | 13 165    | 10 284 | 9.9%       | 1.006                              |
| Comb                                                           | cosmic       | 169               | 12     | 142       | 616    | 8.0%       | 1.095                              |
| <i>with a tagged proton required (measurement phase space)</i> |              |                   |        |           |        |            |                                    |
| FHC                                                            | CC0 $\pi$    | 8 268             | 484    | 7 636     | 1 418  | 6.0%       | 1.018                              |
| FHC                                                            | multi- $\pi$ | 54                | 9.8    | 46        | 17     | 17.7%      | 0.978                              |
| FHC                                                            | $\pi^0$      | 3 988             | 292    | 3 730     | 1 162  | 7.3%       | 0.991                              |
| FHC                                                            | cosmic       | 24                | 1.7    | 23        | 5.8    | 6.7%       | 0.956                              |
| RHC                                                            | CC0 $\pi$    | 8 287             | 499    | 7 812     | 1 473  | 6.0%       | 0.997                              |
| RHC                                                            | multi- $\pi$ | 83                | 14     | 58        | 18     | 19.8%      | 1.143                              |
| RHC                                                            | $\pi^0$      | 5 031             | 332    | 4 487     | 1 207  | 6.9%       | 1.044                              |
| RHC                                                            | cosmic       | 36                | 2.3    | 29        | 6.0    | 7.5%       | 1.158                              |
| Comb                                                           | CC0 $\pi$    | 16 555            | 983    | 15 448    | 2 891  | 6.0%       | 1.008                              |
| Comb                                                           | multi- $\pi$ | 137               | 24     | 104       | 36     | 18.9%      | 1.071                              |
| Comb                                                           | $\pi^0$      | 9 019             | 624    | 8 218     | 2 370  | 7.1%       | 1.020                              |
| Comb                                                           | cosmic       | 60                | 4.0    | 52        | 12     | 7.1%       | 1.067                              |

## 10 Proton-tagged subsample: hadronic invariant mass and transverse kinematic imbalance

Requiring an identified proton in the final state, in addition to the muon and charged pion, reconstructs the full visible hadronic system and opens two classes of observable inaccessible to the inclusive CC  $1\mu 1\pi Xp$  measurement: the *hadronic invariant mass*, which probes the  $\Delta(1232)$  and higher resonances directly, and the *transverse kinematic imbalance* (TKI) between the muon and the hadronic system, which is sensitive to Fermi motion and final-state interactions. This subsample is implemented as a dedicated selection, CC1 $\mu$ 1 $\pi$ 1p, that inherits the full CC  $1\mu 1\pi Xp$  selection (§6) and additionally requires a leading proton candidate; the most proton-like track (log-likelihood-ratio PID score below the proton cut), contained in the fiducial volume and above the 0.3 GeV/ $c$  tracking threshold. The signal is correspondingly narrowed to the inclusive signal plus at least one true proton above 0.3 GeV/ $c$ . On the Run-1 FHC sample this yields a selection efficiency of 11.5% and a purity of 52% (versus 17.5% and  $\sim 56\%$  for the inclusive selection); the proton tag trades signal statistics for a fully reconstructed hadronic final state.

### 10.1 Selection performance and cut-flow

The full-exposure cut-flow of the proton-tagged subsample is the inherited CC  $1\mu 1\pi Xp$  sequence followed by the leading-proton identification (“IdentProton”), which defines the subsample.

Table 37 gives the per-category, per-cut breakdown for the three beam configurations and Figure 29 the corresponding stacks; both use the same per-run POT scaling as the inclusive cut-flow (Table 9), and “signal” here is the proton-tagged signal (inclusive signal plus a true proton above 0.3 GeV/c). The proton-identification cut is the key discriminator: at FHC it rejects  $\sim 64\%$  of the neutrino background while retaining  $\sim 71\%$  of the signal, lifting the purity from 33% at the inclusive final cut (Nonprotons) to 48% in the proton-tagged sample.




Figure 29: Proton-tagged (CC1mu1pi1p) cut-flow yields at full exposure (log scale; MC-prediction stack, Okabe-Ito: signal blue, background orange, EXT grey, dirt green). The last stage, IdentProton, is the leading-proton identification that defines the subsample.

Table 37: Proton-tagged (CC1 $\mu$ 1 $\pi$ 1 $p$ ) cut-flow yields at full exposure for FHC, RHC and combined (MC prediction; data blind). Stages None–Nonprotons are the inherited inclusive selection with the proton-tagged signal definition; the final IdentProton stage adds the leading-proton identification. Signal/Bkg from the  $\nu_\mu$  overlays POT-scaled per run, EXT by the data/EXT trigger ratio, dirt from the GENIE NuMI dirt overlay. Pred. = Signal + Bkg + EXT + Dirt. Eff. is the cumulative signal efficiency relative to the generated signal (the None stage, where all true signal is counted); Pur. = Signal/Pred.

| Cut                                                                     | Signal     | Bkg MC     | EXT        | Dirt       | Pred.        | Eff. [%]    | Pur. [%]    |
|-------------------------------------------------------------------------|------------|------------|------------|------------|--------------|-------------|-------------|
| <i>FHC (Runs 1,2,4,5; <math>8.857 \times 10^{20}</math> POT)</i>        |            |            |            |            |              |             |             |
| None                                                                    | 3 387      | 281 372    | 24 410 022 | 36 142     | 24 730 924   | 100.0       | 0.01        |
| InFiducialVol                                                           | 2 834      | 63 944     | 1 041 637  | 1 890      | 1 110 306    | 83.7        | 0.26        |
| Topological                                                             | 2 238      | 29 303     | 35 184     | 204        | 66 929       | 66.1        | 3.3         |
| MuonCandidate                                                           | 2 031      | 20 092     | 15 516     | 137        | 37 776       | 60.0        | 5.4         |
| ContainedPion                                                           | 852        | 2 637      | 1 376      | 25         | 4 890        | 25.2        | 17.4        |
| MuonIn3Planes                                                           | 812        | 2 499      | 1 145      | 17         | 4 473        | 24.0        | 18.2        |
| PionIn3Planes                                                           | 754        | 2 255      | 872        | 7.0        | 3 888        | 22.3        | 19.4        |
| ShowerCut                                                               | 632        | 1 313      | 569        | 6.2        | 2 521        | 18.7        | 25.1        |
| OpeningAngle                                                            | 628        | 1 226      | 190        | 2.0        | 2 045        | 18.5        | 30.7        |
| Nonprotons                                                              | 568        | 1 049      | 130        | 1.3        | 1 749        | 16.8        | 32.5        |
| <b>IdentProton</b>                                                      | <b>401</b> | <b>373</b> | <b>53</b>  | <b>0.4</b> | <b>828</b>   | <b>11.8</b> | <b>48.4</b> |
| <i>RHC (Runs 1,2,3,4a,4b,4c; <math>1.108 \times 10^{21}</math> POT)</i> |            |            |            |            |              |             |             |
| None                                                                    | 3 638      | 306 840    | 25 344 643 | 28 032     | 25 683 152   | 100.0       | 0.01        |
| InFiducialVol                                                           | 3 064      | 71 162     | 1 081 520  | 1 466      | 1 157 212    | 84.2        | 0.26        |
| Topological                                                             | 2 455      | 33 134     | 36 532     | 158        | 72 278       | 67.5        | 3.4         |
| MuonCandidate                                                           | 2 234      | 22 769     | 16 110     | 106        | 41 219       | 61.4        | 5.4         |
| ContainedPion                                                           | 935        | 3 291      | 1 429      | 19         | 5 674        | 25.7        | 16.5        |
| MuonIn3Planes                                                           | 892        | 3 119      | 1 189      | 13         | 5 213        | 24.5        | 17.1        |
| PionIn3Planes                                                           | 831        | 2 835      | 905        | 5.5        | 4 577        | 22.9        | 18.2        |
| ShowerCut                                                               | 687        | 1 544      | 591        | 4.8        | 2 827        | 18.9        | 24.3        |
| OpeningAngle                                                            | 683        | 1 444      | 197        | 1.5        | 2 325        | 18.8        | 29.4        |
| Nonprotons                                                              | 611        | 1 217      | 136        | 1.0        | 1 964        | 16.8        | 31.1        |
| <b>IdentProton</b>                                                      | <b>436</b> | <b>443</b> | <b>55</b>  | <b>0.3</b> | <b>934</b>   | <b>12.0</b> | <b>46.6</b> |
| <i>Combined (<math>1.994 \times 10^{21}</math> POT)</i>                 |            |            |            |            |              |             |             |
| None                                                                    | 7 025      | 588 212    | 49 755 076 | 36 142     | 50 386 456   | 100.0       | 0.01        |
| InFiducialVol                                                           | 5 898      | 135 106    | 2 123 174  | 1 890      | 2 266 069    | 84.0        | 0.26        |
| Topological                                                             | 4 693      | 62 436     | 71 717     | 204        | 139 050      | 66.8        | 3.4         |
| MuonCandidate                                                           | 4 265      | 42 860     | 31 627     | 137        | 78 889       | 60.7        | 5.4         |
| ContainedPion                                                           | 1 787      | 5 928      | 2 805      | 25         | 10 545       | 25.4        | 16.9        |
| MuonIn3Planes                                                           | 1 704      | 5 618      | 2 333      | 17         | 9 672        | 24.3        | 17.6        |
| PionIn3Planes                                                           | 1 585      | 5 091      | 1 777      | 7.0        | 8 460        | 22.6        | 18.7        |
| ShowerCut                                                               | 1 320      | 2 857      | 1 161      | 6.2        | 5 344        | 18.8        | 24.7        |
| OpeningAngle                                                            | 1 310      | 2 670      | 387        | 2.0        | 4 369        | 18.7        | 30.0        |
| Nonprotons                                                              | 1 179      | 2 266      | 266        | 1.3        | 3 712        | 16.8        | 31.8        |
| <b>IdentProton</b>                                                      | <b>837</b> | <b>816</b> | <b>109</b> | <b>0.4</b> | <b>1 762</b> | <b>11.9</b> | <b>47.5</b> |

**Proton-tag performance.** Beyond the yields, the leading-proton identification (log-likelihood-ratio PID score below the proton cut,  $p_p \geq 0.3$  GeV/ $c$ , contained) is a reliable tag. Truth-matched on the FHC  $\nu_\mu$  overlay, 87% of the tagged events contain an energetic true proton ( $p_p > 0.3$  GeV/ $c$ ); a  $\sim 13\%$  mis-tag rate, dominated by a charged pion or other track being taken for the proton, and 52% are the full  $\mu\pi p$  signal topology. The reconstructed proton momentum, from the proton-hypothesis range kinetic energy, carries a  $-9\%$  bias (underestimate) and a 21% resolution. In cross-section terms the tag rejects  $\sim 64\%$  of the neutrino background for a  $\sim 30\%$  signal loss, improving the signal-to-background figure of merit  $S/\sqrt{B}$  by  $\sim 17\%$ ; the residual mis-tag and

momentum smearing are folded into the response matrix and corrected by the unfolding.

Eleven differential observables are measured in the proton-tagged subsample: the six hadronic-mass and transverse-kinematic-imbalance (W/TKI) variables specific to this selection, plus the five muon/pion kinematic observables of the inclusive analysis ( $p_\mu$ ,  $p_\pi$ ,  $\cos\theta_\mu$ ,  $\cos\theta_\pi$ ,  $\theta_{\mu\pi}$ ) re-measured on the proton-tagged phase space so that the two selections share a common set of kinematic distributions. The two hadronic-mass variables are the *directly measured*  $\pi p$  invariant mass,  $W_{\pi p} = [(E_\pi + E_p)^2 - (\vec{p}_\pi + \vec{p}_p)^2]^{1/2}$ , which makes no assumption on the neutrino direction, and the *calorimetric* hadronic mass  $W_{\text{had}} = [m_p^2 + 2m_p(E_{\text{cal}} - E_\mu) - Q^2]^{1/2}$  built from the reconstructed calorimetric energy. The four TKI observables; the transverse-momentum imbalance  $\delta p_T$ , its two angles  $\delta\alpha_T$  and  $\delta\phi_T$ , and the inferred struck-nucleon momentum  $p_n$ , are computed from the muon versus the summed ( $\pi + p$ ) hadronic system using the framework STV tool. Proton momenta are obtained from the proton-hypothesis range-based kinetic energy. The six W/TKI observables are each measured in six equal-population bins, matching the granularity of the inclusive analysis, now that the per-run POT normalisation (§B) removes the background-subtraction bias that had previously forced a coarser three-bin binning for the lower-statistics reverse-horn-current sample, while the five inherited kinematic observables use the inclusive-analysis binning (four to seven bins). All bin edges are set from the validated true-signal distributions (Table 38).

### Reconstruction from input quantities

All six observables are built from the three candidate tracks (muon, charged pion, proton) using only reconstructed quantities stored in the input ntuples: the track direction cosines  $\hat{n} = (\text{track\_dirx}, \text{track\_diry}, \text{track\_dirz})$ , the muon-hypothesis range and multiple-scattering momenta (`track_range_mom_mu`, `track_mcs_mom_mu`) and the proton-hypothesis calorimetric kinetic energy (`track_kinetic_energy_p`).

*Candidate three-momenta.* The muon momentum magnitude is range-based when the muon track ends inside the fiducial volume,  $p_\mu = \text{track\_range\_mom\_mu}[\mu]$ , and multiple-Coulomb-scattering based otherwise,  $p_\mu = \text{track\_mcs\_mom\_mu}[\mu]$ . The charged pion, being minimum-ionising, is assigned the muon-hypothesis range momentum  $p_\pi = \text{track\_range\_mom\_mu}[\pi]$ . The proton momentum follows from its calorimetric kinetic energy under the proton hypothesis,  $p_p = \sqrt{T_p^2 + 2m_p T_p}$  with  $T_p = \text{track\_kinetic\_energy\_p}[p]$ . Each three-momentum is  $\vec{p}_i = p_i \hat{n}_i$ , and the energies are  $E_\mu = \sqrt{p_\mu^2 + m_\mu^2}$ ,  $E_\pi = \sqrt{p_\pi^2 + m_\pi^2}$ ,  $E_p = T_p + m_p$  (masses  $m_\mu = 105.7$ ,  $m_\pi = 139.6$ ,  $m_p = 938.3$  MeV).

*Directly measured mass  $W_{\pi p}$ .* The invariant mass of the pion-proton four-vector sum, using no neutrino-direction assumption:

$$W_{\pi p} = \sqrt{(E_\pi + E_p)^2 - |\vec{p}_\pi + \vec{p}_p|^2}.$$

*Transverse kinematic imbalance.* The TKI observables treat the muon against the visible hadronic system  $\vec{p}_{\text{had}} = \vec{p}_\pi + \vec{p}_p$ ,  $E_{\text{had}} = E_\pi + E_p$ . With the components transverse to the beam ( $\hat{z}$ ) written  $\vec{p}_T^\mu$  and  $\vec{p}_T^{\text{had}}$ ,

$$\delta p_T = |\vec{p}_T^\mu + \vec{p}_T^{\text{had}}|, \quad \delta\alpha_T = \arccos \frac{-\vec{p}_T^\mu \cdot (\vec{p}_T^\mu + \vec{p}_T^{\text{had}})}{|\vec{p}_T^\mu| \delta p_T}, \quad \delta\phi_T = \arccos \frac{-\vec{p}_T^\mu \cdot \vec{p}_T^{\text{had}}}{|\vec{p}_T^\mu| |\vec{p}_T^{\text{had}}|},$$

both angles in degrees folded onto  $[0, 180]$ . For a stationary nucleon with no final-state interactions  $\delta p_T \rightarrow 0$ ; Fermi motion broadens it and FSI drive  $\delta\alpha_T \rightarrow 180^\circ$ .

*Calorimetric mass  $W_{\text{had}}$  and struck-nucleon momentum  $p_n$ .* The calorimetric available energy is  $E_{\text{cal}} = E_\mu + (E_{\text{had}} - m_p) + E_b$  with the argon removal energy  $E_b = 24.78$  MeV, giving the momentum transfer  $Q^2 = -[(0, 0, E_{\text{cal}}, E_{\text{cal}}) - p^\mu]^2$  and

$$W_{\text{had}} = \sqrt{m_p^2 + 2m_p(E_{\text{cal}} - E_\mu) - Q^2}$$

(returned as 0 where reconstruction fluctuations make the argument negative). The inferred initial-state nucleon momentum combines the transverse imbalance with a longitudinal component  $p_L$  from energy-momentum balance,  $p_n = \sqrt{\delta p_T^2 + p_L^2}$ . The transverse-imbalance and calorimetric quantities are evaluated with the framework STV tool (`STVTools::CalculateSTVs`, option 1); it was formulated for a single-proton hadronic system and is applied here to the summed ( $\pi p$ ) system with the proton rest mass, an approximation adequate for the leading-order kinematics.

Table 38: Binning of the six W/TKI proton-tagged observables (six equal-population bins each), chosen from the validated true-signal distributions. The first and last bins are open so the full signal phase space is partitioned.  $W_{\pi p}$  brackets the  $\Delta(1232)$  peak; the  $\delta\alpha_T$  and  $\delta\phi_T$  binnings are finer toward  $180^\circ$ , where final-state interactions pile up. The five inherited kinematic observables ( $p_\mu$ ,  $p_\pi$ ,  $\cos\theta_\mu$ ,  $\cos\theta_\pi$ ,  $\theta_{\mu\pi}$ ) use the inclusive-analysis binning of §D unchanged.

| Observable       | Definition                                 | Bin edges                                           |
|------------------|--------------------------------------------|-----------------------------------------------------|
| $W_{\pi p}$      | $M(\pi p)$ , direct                        | 1.08, 1.19, 1.23, 1.27, 1.34, 1.47, 2.00 GeV/ $c^2$ |
| $W_{\text{had}}$ | calorimetric                               | 0, 0.78, 1.08, 1.21, 1.31, 1.44, 1.90 GeV/ $c^2$    |
| $\delta p_T$     | $ \vec{p}_T^\mu + \vec{p}_T^{\text{had}} $ | 0, 0.43, 0.59, 0.72, 0.88, 1.17, 1.60 GeV/ $c$      |
| $\delta\alpha_T$ | TKI angle                                  | 0, 80, 120, 142, 158, 170, $180^\circ$              |
| $\delta\phi_T$   | TKI angle                                  | 0, 37, 63, 84, 107, 139, $180^\circ$                |
| $p_n$            | inferred nucleon momentum                  | 0, 0.52, 0.67, 0.81, 0.98, 1.25, 1.70 GeV/ $c$      |

The pipeline is validated end-to-end on FHC with a fake-data closure: the Poisson-thrown central-value pseudo-data are unfolded with the same Wiener-SVD procedure (§8) and compared to the  $A_C$ -smeared truth. Table 39 shows the result. The bin-by-bin  $\chi^2/\text{ndf}$  is well below unity for every observable, confirming the unfolded pseudo-data are statistically consistent with truth and that the response, background subtraction and normalisation are self-consistent. The integrated-cross-section ratios (0.94–1.19) reflect the same non-norm-preserving Wiener-SVD additional smearing  $A_C$  seen in the inclusive measurement (§8.14), largest for  $\delta\alpha_T$  and  $\theta_{\mu\pi}$ . The systematic treatment here includes the flux, cross-section and reinteraction terms from the MC weight universes; the detector-variation term is deferred pending a dedicated reprocessing of the detector-variation samples through this selection.

The reverse-horn-current measurement is complete at the same finer binning (Table 40). All eleven RHC observables close positive, with  $\sigma_{\text{int}}$  clustered in  $0.41\text{--}0.58 \times 10^{-38}$  cm<sup>2</sup>/Ar, just below the FHC band (0.50–0.67), as expected for the  $\bar{\nu}_\mu$ -enriched RHC flux; and  $\chi^2/\text{ndf} \lesssim 1$  for almost every observable.

Reaching this point required resolving a normalisation subtlety that is worth recording, because it produced spurious *negative* cross sections for this subsample at an intermediate stage. The processed ntuples use two different POT conventions: the files behind the inclusive selection store the *total* run MC POT duplicated in every file of a multi-file run, whereas the proton-tagged files store each file’s *own* POT. Normalising by the individual file POT is therefore correct for the first set but scales every split of a multi-file run to the full run exposure in the second, inflating the simulated background by the split multiplicity ( $\sim 3\times$  for RHC) and driving the background-subtracted spectrum negative. Normalising by the summed per-file POT has the mirror problem. The analysis now normalises by the sum of the *distinct* per-file POT values, which reduces to the correct run total under either convention; FHC, with one MC file per run, is unaffected by the choice.

The combined-sample measurement is in production and will be added here.

Table 39: FHC fake-data closure for the eleven proton-tagged observables at the finer binning.  $\sigma_{\text{int}}$  is the integrated (unfolded) cross section over the measured phase space ( $10^{-38} \text{ cm}^2/\text{Ar}$ ); “unf./truth” is the ratio of integrals;  $\chi^2$  is the bin-by-bin comparison of unfolded pseudo-data to  $A_C$ -smeared truth (ndf = number of bins). The upper block is the six W/TKI observables (six bins each); the lower block is the five kinematic observables inherited from the inclusive analysis. The small  $\chi^2/\text{ndf}$  validates the chain; the integral ratios are the  $A_C$  smearing effect, not a bias.

| Observable        | $\sigma_{\text{int}}$ | unf./truth | $\chi^2/\text{ndf}$ |
|-------------------|-----------------------|------------|---------------------|
| $W_{\pi p}$       | 0.550                 | 1.12       | 0.19/6              |
| $W_{\text{had}}$  | 0.500                 | 1.13       | 0.86/6              |
| $\delta p_T$      | 0.482                 | 1.15       | 1.87/6              |
| $\delta \alpha_T$ | 0.551                 | 1.20       | 0.67/6              |
| $\delta \phi_T$   | 0.592                 | 1.17       | 4.02/6              |
| $p_n$             | 0.432                 | 1.00       | 0.19/6              |
| $p_\mu$           | 0.555                 | 1.16       | 0.54/7              |
| $p_\pi$           | 0.596                 | 1.15       | 0.15/2              |
| $\cos \theta_\mu$ | 0.555                 | 1.11       | 0.18/5              |
| $\cos \theta_\pi$ | 0.479                 | 1.13       | 0.18/4              |
| $\theta_{\mu\pi}$ | 0.515                 | 1.14       | 0.66/5              |

Figure 30 shows the six FHC W/TKI differential cross sections at the finer six-bin binning, overlaid with all four generator predictions. The generator W/TKI predictions were regenerated at this binning (GENIE and GiBUU natively, NuWro and NEUT from their event vectors inside the SL7 container); the rebinning conserves each generator’s integral to the previous three-bin value (GENIE 0.501, NuWro 0.646, GiBUU 0.735, NEUT  $0.783 \times 10^{-38} \text{ cm}^2/\text{Ar}$ ). All curves, data, tune and generators, are  $A_C$ -smeared, so the comparison is made consistently in the measurement space (§10.2). The predictions are built at truth level from each generator’s event-level final state, GENIE `gst` events, GiBUU `FinalEvents`, and the NuWro and NEUT event vectors reprocessed in their respective SL7-container environments, by applying the inclusive signal plus a leading true proton above 0.3 GeV/ $c$  and evaluating the six observables with the same formulas as the selection (§10.1), then combining the  $\nu_\mu$  and  $\bar{\nu}_\mu$  samples over the FHC flux to a per-argon cross section. The four generators span a factor of  $\sim 1.6$  in the integrated proton-tagged cross section, GENIE lowest (0.50), then NuWro (0.65), GiBUU (0.74) and NEUT highest ( $0.78 \times 10^{-38} \text{ cm}^2/\text{Ar}$ ), bracketing the data ( $\sim 0.58$ ), and they differ most in the low- $\delta p_T$  / high- $p_n$  region sensitive to the proton kinematics and final-state interactions. As throughout, the unfolded fake data sit above the  $A_C$ -smeared truth by the Wiener-SVD normalisation factor.

Table 40: RHC fake-data closure for the eleven proton-tagged observables at the finer binning, columns as in Table 39. All observables close positive with the convention-agnostic MC POT normalisation described above;  $\sigma_{\text{int}}$  sits just below the FHC band, consistent with the  $\bar{\nu}_\mu$ -enriched RHC flux.

| Observable        | $\sigma_{\text{int}}$ | unf./truth | $\chi^2/\text{ndf}$ |
|-------------------|-----------------------|------------|---------------------|
| $W_{\pi p}$       | 0.515                 | 1.22       | 3.42/6              |
| $W_{\text{had}}$  | 0.540                 | 1.34       | 3.02/6              |
| $\delta p_T$      | 0.452                 | 1.21       | 1.16/6              |
| $\delta \alpha_T$ | 0.472                 | 1.22       | 5.12/6              |
| $\delta \phi_T$   | 0.596                 | 1.21       | 1.55/6              |
| $p_n$             | 0.464                 | 1.07       | 3.18/6              |
| $p_\mu$           | 0.499                 | 1.23       | 1.21/7              |
| $p_\pi$           | 0.497                 | 1.10       | 2.36/2              |
| $\cos \theta_\mu$ | 0.559                 | 1.28       | 0.78/5              |
| $\cos \theta_\pi$ | 0.474                 | 1.24       | 4.04/4              |
| $\theta_{\mu\pi}$ | 0.465                 | 1.16       | 0.48/5              |

Table 41: Combined FHC+RHC fake-data closure for the eleven proton-tagged observables at the finer binning, columns as in Table 39. All eleven close positive and every  $\chi^2$  is comfortably below its number of bins. This completes the proton-tagged measurement in all three beam configurations. Note that the combined result is an independent extraction, not a fluence-weighted average of the FHC and RHC ones, so it is not required to lie between them; see the discussion below.

| Observable        | $\sigma_{\text{int}}$ | unf./truth | $\chi^2/\text{ndf}$ |
|-------------------|-----------------------|------------|---------------------|
| $W_{\pi p}$       | 0.531                 | 1.17       | 1.34/6              |
| $W_{\text{had}}$  | 0.460                 | 1.14       | 2.18/6              |
| $\delta p_T$      | 0.463                 | 1.12       | 0.74/6              |
| $\delta \alpha_T$ | 0.516                 | 1.22       | 3.44/6              |
| $\delta \phi_T$   | 0.576                 | 1.14       | 1.45/6              |
| $p_n$             | 0.472                 | 1.13       | 0.90/6              |
| $p_\mu$           | 0.554                 | 1.22       | 0.75/7              |
| $p_\pi$           | 0.539                 | 1.13       | 1.01/2              |
| $\cos \theta_\mu$ | 0.580                 | 1.17       | 0.68/5              |
| $\cos \theta_\pi$ | 0.462                 | 1.16       | 2.41/4              |
| $\theta_{\mu\pi}$ | 0.484                 | 1.15       | 0.19/5              |

With all three configurations in hand, the combined  $\sigma_{\text{int}}$  lies between the FHC and RHC values for seven of the eleven observables. The four exceptions are small and worth stating explicitly rather than glossing:  $p_n$  sits 6.9% above the FHC value,  $\cos \theta_\mu$  1.7% above it, and  $W_{\pi p}$  and  $p_\mu$  fall 2.4% and 1.5% below the RHC value. The combined measurement is *not* required to bracket the two single-mode ones: it is an independent extraction with its own response matrix, its own regularisation and its own  $A_C$ , not a fluence-weighted average of the FHC and RHC results. Excursions at the few-per-cent level are therefore expected, and all four are far inside the  $\sim 30\%$  systematic uncertainty.

The spread across observables within a configuration, 0.45 to 0.67 for the combined set, is likewise not a physics spread but the residual of the additional smearing discussed in the next section: each observable's  $A_C$  rescales the integral by a different factor, so the eleven numbers

are not eleven independent measurements of the same quantity.

## 10.2 Additional smearing in the proton-tagged observables

Because the Wiener–SVD additional-smearing matrix  $A_C$  is not norm-preserving, each observable’s predictions are rescaled by a different factor when they are brought into the measurement space. Table 42 gives that factor, measured as the ratio of the  $A_C$ -smeared to the truth-level generator integral, for the six proton-tagged observables. It is a property of the response and the binning, not of the model: GENIE and GiBUU agree to a few per cent throughout.

The two hadronic-mass variables are the *least* affected in the whole analysis (0.77–0.88). This is worth stating explicitly because  $W_{\pi p}$  is the sharpest distribution measured here; the ratio of its largest to smallest bin is  $\sim 36$ , against  $\sim 19$  for  $\cos\theta_\mu$ , whose forward peak is suppressed by a factor  $\sim 0.33$  (§9.2). Sharpness alone therefore does not drive the suppression:  $A_C$  removes structure that the *response* cannot resolve, and  $W_{\pi p}$  is built from directly measured pion and proton momenta with good resolution and equal-population bins that track the  $\Delta(1232)$  peak, so the peak survives. No reparameterisation of  $W$  is required.

Two observables do carry a caveat.  $\delta p_T$  is strongly suppressed (0.43–0.45), closer to the  $\cos\theta_\mu$  regime than to  $W$ , and  $p_n$  is *enhanced* by a factor  $\sim 1.7$ . Both effects are applied identically to data, tune and generators, so the closure is unaffected, but any model comparison read off the  $\delta p_T$  and especially the  $p_n$  panels is distorted by these factors and should not be interpreted as a direct statement about the truth-level predictions until the binning and response for those two observables have been revisited.

Table 42: Additional-smearing factor  $A_C$  for the proton-tagged observables (FHC), measured as the ratio of the  $A_C$ -smeared to the truth-level generator integral. The inclusive observables are given for scale.

| Observable                                                                                                                                                                                                                                                                                    | GENIE | GiBUU | Comment            |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------|-------|--------------------|
| $W_{\text{had}}$                                                                                                                                                                                                                                                                              | 0.69  | —     |                    |
| $W_{\pi p}$                                                                                                                                                                                                                                                                                   | 0.77  | —     | least smeared      |
| $\delta\alpha_T$                                                                                                                                                                                                                                                                              | 0.71  | —     |                    |
| $\delta\phi_T$                                                                                                                                                                                                                                                                                | 0.80  | —     |                    |
| $\delta p_T$                                                                                                                                                                                                                                                                                  | 0.67  | —     |                    |
| $p_n$                                                                                                                                                                                                                                                                                         | 0.69  | —     | no longer enhanced |
| <i>Post-fix values, mean <math>A_C</math> row sum. The pre-fix set (0.88, 0.78, 0.63, 0.61, 0.43, 1.71) spanned a factor of four and amplified <math>p_n</math>; it is now uniform to within 0.13, which is why the <math>\chi^2</math> ordering can no longer be attributed to smearing.</i> |       |       |                    |






Figure 30: Proton-tagged (CC1mu1pi1p) differential cross sections for FHC: the  $\pi p$  invariant mass  $W_{\pi p}$ , the calorimetric hadronic mass  $W_{\text{had}}$ , and the four transverse-kinematic-imbalance observables  $\delta p_T$ ,  $\delta\alpha_T$ ,  $\delta\phi_T$  and  $p_n$ . Unfolded fake data (black points) are compared to the  $A_C$ -smeared fake-data truth (grey dashed), the MicroBooNE tune (black), and the GENIE (blue), GiBUU (green), NEUT (purple) and NuWro (orange) predictions built for the same proton-tagged phase space.  $W_{\pi p}$  falls from the  $\Delta(1232)$  threshold region;  $\delta\alpha_T$  rises toward  $180^\circ$ , the signature of final-state interactions.

## 11 Systematic Uncertainties

Full systematic uncertainty propagation has not yet been performed. Table 43 lists the sources and their estimated impacts.

Table 43: Systematic uncertainty sources. “Not quantified” means the branch is readable but propagation code has not been written.

| Source                        | Branch / method           | Impact                              |
|-------------------------------|---------------------------|-------------------------------------|
| Flux (PPFX multisims)         | <code>weightsFlux</code>  | $\sim 10\text{--}15\%$ norm.        |
| GENIE cross sections          | <code>weightsGenie</code> | $\sim 20\text{--}30\%$ (model)      |
| Hadron re-interaction         | <code>weightsReint</code> | $\sim 5\text{--}10\%$               |
| Detector response             | Dedicated samples         | $\sim 5\text{--}15\%$               |
| EXT normalisation             | Trigger-count ratio       | $\sim 3\text{--}5\%$                |
| POT counting                  | Beam toroids              | $\sim 2\%$                          |
| Flux integral (absolute)      | PPFX integral             | Blocks absolute $\sigma$            |
| Statistical (427 data events) | n/a                       | $\sim 5\%$ on $\sigma_{\text{int}}$ |

## 12 Known Issues and Required Follow-up

Table 44: Outstanding issues and their impact on the final result.

| Issue                                                              | Status                                  | Impact                                                                                                                                            |
|--------------------------------------------------------------------|-----------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------|
| FLUX_PER_POT confirmed                                             | <b>Resolved</b>                         | $\Phi = 6.81159 \times 10^{-10} \text{ cm}^{-2} \text{ POT}^{-1}$<br>from NuMI flux histograms                                                    |
| Unfolding covariance (stat-only $\rightarrow$ full)                | <b>Resolved</b>                         | Framework <code>PredTotal</code> +data<br>Poisson imported; removed<br>$\sim 26\%$ bias                                                           |
| EXT subtraction on NuWro fake data                                 | <b>Resolved</b>                         | Cancelled via <code>fake_data_ext_cancel</code> .<br><i>Follow-up:</i> re-enable before unblinding                                                |
| $\theta_{\mu\pi}$ binning (10 vs 9, C4)                            | <b>Resolved</b>                         | Custom aligned to framework 9-bin scheme                                                                                                          |
| Multisim systematics not propagated (custom)                       | Not started                             | Imported from framework for the covariance; full custom propagation pending                                                                       |
| Negative tail bins                                                 | <b>Resolved</b>                         | All bins now positive after EXT cancellation; no longer present                                                                                   |
| Runs 2, 4, 5 beam-on files not available                           | Data files needed                       | $\times 5.4$ exposure increase for the real-data measurement                                                                                      |
| Generator comparisons (GENIE, NuWro)                               | Blocked by missing AVX on compute nodes | NEUT comparison runnable locally                                                                                                                  |
| CRT veto branches ( <code>crthitpe</code> , <code>crtveto</code> ) | Dead (all-zero in NuMI ntuples)         | Replaced by <code>bdt_cosmic</code> (disabled at default 0.0)                                                                                     |
| Official XSecAnalyzer NuWro closure                                | <b>Done</b> (all 5 obs)                 | Cross-validates the unfolding; NuWro $\approx 2\times$ GENIE (Sec. C)                                                                             |
| Native RHC detector variations                                     | <b>Resolved</b>                         | Run-4 RHC detVar (8 knobs) now used; replaces the Run-1 FHC stand-in (Sec. 9)                                                                     |
| Run-5 FHC detVar Recomb2/SCE                                       | <b>Resolved</b>                         | Were intrinsic- $\nu_e$ overlays (0% $\nu_\mu$ ); Run-4 FHC used as the stand-in for those two knobs                                              |
| Combined FHC+RHC joint flux systematic                             | <b>Resolved</b>                         | Per-universe summing over the shared 600 PPFx universes is correct; the block was a detVar duplicate-file guard, now fixed to accumulate (Sec. 9) |
| Combined per-mode normalisation                                    | <b>Resolved</b>                         | Per-mode <code>summed_pot</code> so each horn scales to its own data exposure, not MC POT (Sec. 9)                                                |

## 13 Analysis Pipeline

The primary measurement is produced by a single driver, `run_analysis.sh`, which chains the selection, the per-observable cross-section extraction, and the comparison/diagnostic stages:

```

selection/ccpi_selection.C  ->  histograms.root + unfolding_inputs.root
                             | (5 observables x 8 methods)
xsec/ccpi_xsec.C(obs,method) -> xsec_{obs}_{method}.root  (40 files)
xsec/compare_unfolding.C    -> compare_plots/

```

```
xsec/compare_generators.C    -> gen_compare/
selection/event_display.C    -> event_displays/
```

Each output file is validated for freshness and non-zero content after the step that produces it. The official-framework cross-check of Section C is driven separately (`xsec_analyzer/run_nuwro_observables.sh`).

## 14 Binning Optimisation, Systematic Breakdown, and Threshold Validation

The reconstructed spectra, response matrices, efficiencies, and differential cross sections shown throughout this note are the XSECANALYZER framework outputs on the *optimised* binning defined here; the four-generator comparisons of Fig. 32 onward likewise use it.

### 14.1 Momentum-threshold validation

The two signal-definition momentum thresholds sit at the selection efficiency turn-on, measured against a threshold-free true signal (the CC  $\nu_\mu$   $1\mu 1\pi^\pm$  topology with the momentum/angle cuts removed):

| Threshold             | below                  | above                  | turn-on                   |
|-----------------------|------------------------|------------------------|---------------------------|
| $p_\mu > 0.15$ GeV/c  | $\sim 0\text{--}1.5\%$ | $\sim 9\text{--}12\%$  | 0.145 $\rightarrow$ 0.155 |
| $p_\pi > 0.175$ GeV/c | $\sim 0\text{--}3\%$   | $\sim 12\text{--}16\%$ | 0.165 $\rightarrow$ 0.185 |

Both are correctly placed, lowering them would add near-zero-efficiency phase space, raising them would discard measurable signal. The reconstructed pion momentum (muon-hypothesis range) is biased low and increasingly so with momentum (mean reco $-\text{true} \approx 0$  at threshold,  $-0.15$  GeV/c at 0.40) from the  $\mu$ -vs- $\pi$  mass hypothesis plus pion re-interaction; a pion-hypothesis mass correction is applied to the  $p_\pi$  reco bins.

### 14.2 Binning optimisation

Bins were designed to be at least the local RMS detector resolution wide (so the response stays well conditioned and the Wiener-SVD does not amplify the correlated flux/cross-section uncertainty) and to hold  $\geq 45$  selected-signal events (statistical floor). This lowers the total fractional uncertainty in every observable, chiefly through the statistical and unfolding-amplified terms; the flux/beamline term is correlated across bins and sets a floor:

| Observable        | bins               | Total before | Total after  |
|-------------------|--------------------|--------------|--------------|
| $p_\mu$           | 22 $\rightarrow$ 7 | 33.2%        | <b>25.0%</b> |
| $p_\pi$           | 5 $\rightarrow$ 5  | 27.7%        | 27.0%        |
| $\cos \theta_\mu$ | 12 $\rightarrow$ 5 | 31.3%        | <b>26.7%</b> |
| $\cos \theta_\pi$ | 12 $\rightarrow$ 5 | 48.5%        | <b>33.9%</b> |
| $\theta_{\mu\pi}$ | 9 $\rightarrow$ 7  | 28.4%        | <b>23.5%</b> |

### 14.3 Systematic breakdown by source

The framework builds a covariance matrix for every systematic; the source-group decomposition method is illustrated below on a single configuration (the definitive per-configuration breakdowns for the full FHC, RHC, and combined exposures are Tables 31–33). The bin-averaged fractional uncertainty from each source group (on the optimised binning) is:

Table 45: Bin-averaged fractional systematic uncertainty (%) by source group on the optimised binning.

| Source                | $p_\mu$ | $p_\pi$ | $\cos \theta_\mu$ | $\cos \theta_\pi$ | $\theta_{\mu\pi}$ |
|-----------------------|---------|---------|-------------------|-------------------|-------------------|
| <b>Total</b>          | 27.2    | 28.3    | 28.8              | 33.8              | 25.1              |
| Cross section (GENIE) | 8.6     | 8.5     | 9.1               | 9.5               | 8.1               |
| Flux, PPFX            | 21.8    | 23.6    | 20.7              | 25.4              | 18.7              |
| Flux, beamline geom.  | 2.3     | 2.2     | 2.3               | 3.3               | 1.6               |
| Detector              | 9.6     | 10.2    | 14.0              | 14.6              | 10.5              |
| Reinteraction         | 3.2     | 2.6     | 3.4               | 3.4               | 2.5               |
| MC + data stats       | 8.6     | 6.5     | 9.4               | 11.9              | 9.3               |
| POT + targets         | 2.7     | 2.9     | 2.7               | 3.0               | 2.4               |

The NuMI flux uncertainty has two parts, listed separately above: the PPFX hadron-production term (the dominant systematic, 19–26%) and the beamline-geometry term. The latter is built from the 12 geometry-variation weights (`Horn_2kA`, `Horn1/2.x/y_3mm`, `Beam_spot_*`, `Beam_shift_*`, `Horns*_water`, `Target_z_7mm`) injected into the numuMC overlay via `AddBeamlineGeometryWeights`, summed in quadrature as two-sided unisim variations. It is sub-dominant, 1.6–3.3% across the observables. (Building it required a framework fix: the covariance code dereferenced a null histogram when a systematic, here the beamline weights, present in the numuMC but not the dirt overlay, was absent from some MC sample; it now falls back to that sample’s central value, i.e. unit weight.) The per-bin breakdowns are shown in Fig. 31.




Figure 31: Bin-averaged fractional systematic uncertainty by source vs. observable, for FHC (left), RHC (centre) and combined (right); the figure form of Tables 31–33. Colours use the Okabe-Ito colourblind-safe palette with distinct markers per source: total (black), flux (blue), detector (green), cross section (vermillion), reinteraction (purple), data stat (orange).

#### 14.4 Anatomy of the flux term: genuine flux vs. unfolding amplification

The flux (PPFX) term dominates the budget, and at 17–26% it is larger than the  $\sim 17\%$  normalisation uncertainty usually quoted for the NuMI flux. It is therefore worth asking how much of it is the flux model itself and how much is statistical amplification introduced by the unfolding. To separate the two we evaluate the flux covariance at two stages, on the reconstructed spectrum (before unfolding) and on the unfolded result, and, at each stage, both integrated (the fully correlated, normalisation-like number, from the sum of all covariance elements) and bin-averaged (the mean of the diagonal fractional errors, i.e. the number quoted in Table 45 and Fig. 31).

Table 46: Flux (PPFX) fractional uncertainty before and after unfolding. The reconstructed-level value is  $\sim 17\%$  and flat across observables; the genuine flux-normalisation uncertainty. The unfolded value is that same  $\sim 17\%$  amplified by the Wiener–SVD propagation through the finite-statistics response; the amplification tracks the data+MC statistical quality of each observable.

|                                | $p_\mu$ | $p_\pi$ | $\cos \theta_\mu$ | $\cos \theta_\pi$ | $\theta_{\mu\pi}$ |
|--------------------------------|---------|---------|-------------------|-------------------|-------------------|
| Flux, reco (integrated / norm) | 16.9    | 17.0    | 16.7              | 16.7              | 16.7              |
| Flux, reco (bin-averaged)      | 17.7    | 16.4    | 16.6              | 17.0              | 17.0              |
| Flux, unfolded (bin-averaged)  | 19.8    | 22.4    | 19.8              | 25.5              | 17.5              |
| Amplification factor           | 1.12    | 1.37    | 1.20              | 1.49              | 1.03              |
| Excess from unfolding (abs.)   | +2.1    | +6.0    | +3.2              | +8.5              | +0.5              |
| Data+MC stats (unfolded)       | 7.9     | 6.3     | 8.4               | 12.2              | 8.8               |

Three points follow. First, the *genuine* flux uncertainty is  $\sim 17\%$  and identical for all five observables, as expected for a normalisation term. Second, it is not itself contaminated by Monte Carlo statistics: at reco level the integrated (correlated) and bin-averaged (diagonal) numbers agree to better than a percent, whereas statistical noise leaking into the covariance would inflate the diagonal well above the correlated norm. This is expected; the  $\sim 600$  PPFX universes reweight the *same* events, so there is no independent per-universe statistical scatter. Third, the elevated unfolded values are the  $\sim 17\%$  flux *amplified* by the Wiener–SVD propagation  $A_C M V_{\text{flux}} M^\top A_C^\top$  through the finite-statistics response. It is at this step, not in the flux multisim, that the response statistics transfer into the flux term: the amplification is largest exactly where the statistics are thinnest ( $\cos \theta_\pi$ : stat. 12.2%, amplification 1.49 $\times$ , +8.5%) and negligible where the response is well conditioned ( $\theta_{\mu\pi}$ : 1.03 $\times$ , +0.5%). The lever for reducing the flux term is therefore the response conditioning (more overlay statistics, or the observable’s binning and regularisation strength), not the flux model;  $\cos \theta_\pi$ ; the noisiest observable across the entire budget, is where it would matter most.

## 14.5 Multi-run overlay and a resonance-model closure

Two facts frame what more statistics can and cannot do for the flux term. First, the flux *systematic* is a normalisation-and-shape model uncertainty and is therefore POT-independent: adding data or overlay does not change it. Second, Section 14.4 showed that the elevated *unfolded* flux is that  $\sim 17\%$  model uncertainty amplified by the Wiener–SVD propagation through a statistically-limited response, and that the limitation is the *data* statistics, not the overlay. Increasing the overlay alone therefore sharpens the response but leaves the data-limited amplification in place.

To study a genuine multi-run configuration we combine the FHC overlays for Runs 1, 2 and 4 (with Run4=Run4c+Run4d counted once; the file first distributed as “Run 5 FHC” was set aside on inspection, being in fact an intrinsic- $\nu_e$  overlay, 86%  $\nu_e$ , 14%  $\bar{\nu}_e$ , no  $\nu_\mu$  events, hence the large  $\nu_e$ -enriched POT-equivalent and no  $\nu_\mu$  signal; the genuine Run 5  $\nu_\mu$  overlay, a *reweightedPPFX* sample, was subsequently located and is used for the extended set below),  $2.32 \times 10^6$  unique events, normalised to the FHC data POT  $6.626 \times 10^{20}$  (Section 14.6). Because  $\text{CC}\pi^\pm$  is resonance-dominated, robustness to the resonant axial mass is tested with Poisson-thrown fake data reweighted to  $M_A^{\text{RES}} = 1.12 \pm 0.22$  GeV ( $\pm 1\sigma$ ; axial dipole ratio in the true interaction  $Q^2$ , a +8.1% / – 7.2% rate distortion). The pseudo-data are thrown at the true FHC event count so their statistical covariance is realistic; the closure condition is that the unfolded result recovers the  $A_C$ -smeared shifted truth within uncertainties. The Wiener–SVD first- versus second-derivative regularisation is compared on the same response.

The closure succeeds for every observable, both signs of the shift, and both regularisations: the unfolded result recovers the  $A_C$ -smeared shifted truth with  $\chi^2/\text{ndf} < 1$  throughout (Table 48), all  $p$ -values above 0.63 and most above 0.9. The Wiener–SVD machinery is therefore unbiased under a  $\pm 1\sigma$   $M_A^{\text{RES}}$  distortion, and the first- and second-derivative regularisations perform equivalently in the closure (the first-derivative variant carries a slightly larger statistical term). Restoring a realistic (Poisson) data covariance also restores the flux amplification that the degenerate near-Asimov pseudo-data had suppressed.

To separate the mean amplification from single-realisation noise, we throw 15 independent Poisson pseudo-datasets (central value only, no  $M_A^{\text{RES}}$  shift, at the same  $6.626 \times 10^{20}$  FHC exposure) and unfold each (Table 47). Averaged over throws, the unfolded flux stabilises at 24.7–27.1%, a mean amplification of  $1.44\text{--}1.63\times$  over the  $\sim 17\%$  floor, with a data-statistical term of 8–13%. This confirms that the amplification is driven by the data statistics, not the overlay, and is consistent with the mechanism of Section 14.4. The throw-to-throw scatter is small for the momenta ( $p_\mu, p_\pi$ :  $\pm 0.06\text{--}0.08\times$ , flux  $\pm 1.1\text{--}1.4\%$ ) but grows for the angular observables, and is largest for  $\theta_{\mu\pi}$ , whose amplification spans  $1.27\text{--}2.02\times$  (flux 21.4–34.2%) across the 15 throws. At these statistics the amplification is therefore well defined in the mean but individually unstable, most so for  $\theta_{\mu\pi}$  and  $\cos\theta_\pi$ ; a direct consequence of the sparse, data-limited response.

Table 47: Flux amplification averaged over 15 independent Poisson-thrown pseudo-datasets at the FHC data POT: mean  $\pm$  standard deviation and min–max range over throws, for the unfolded flux (bin-averaged) and the amplification relative to the  $\sim 17\%$  reco-level floor.

| Observable        | Amplification (mean $\pm$ sd) | Amp. range   | Unfolded flux    | Flux range   |
|-------------------|-------------------------------|--------------|------------------|--------------|
| $p_\mu$           | $1.44 \pm 0.08\times$         | [1.27, 1.60] | $24.7 \pm 1.4\%$ | [21.9, 27.5] |
| $p_\pi$           | $1.52 \pm 0.06\times$         | [1.40, 1.65] | $26.2 \pm 1.1\%$ | [24.0, 28.4] |
| $\cos\theta_\mu$  | $1.60 \pm 0.14\times$         | [1.41, 1.85] | $26.8 \pm 2.3\%$ | [23.7, 31.1] |
| $\cos\theta_\pi$  | $1.63 \pm 0.11\times$         | [1.45, 1.89] | $27.1 \pm 1.9\%$ | [24.1, 31.6] |
| $\theta_{\mu\pi}$ | $1.58 \pm 0.21\times$         | [1.27, 2.02] | $26.7 \pm 3.6\%$ | [21.4, 34.2] |

Table 48: Fake-data closure:  $\chi^2/\text{ndf}$  of the unfolded result against the  $A_C$ -smeared  $M_A^{\text{RES}}$ -shifted truth, for the two shift signs and the two Wiener–SVD regularisations (number of bins in parentheses). All values are below unity per degree of freedom ( $p > 0.63$ ).

| Observable            | +1 $\sigma$ (2nd) | +1 $\sigma$ (1st) | −1 $\sigma$ (2nd) | −1 $\sigma$ (1st) |
|-----------------------|-------------------|-------------------|-------------------|-------------------|
| $p_\mu$ (7)           | 1.26              | 2.15              | 0.95              | 2.14              |
| $p_\pi$ (5)           | 1.24              | 3.44              | 0.25              | 0.39              |
| $\cos\theta_\mu$ (5)  | 0.47              | 0.63              | 1.90              | 1.24              |
| $\cos\theta_\pi$ (5)  | 2.99              | 1.07              | 0.24              | 0.33              |
| $\theta_{\mu\pi}$ (7) | 1.71              | 3.32              | 1.96              | 4.33              |

Adding the genuine Run 5  $\nu_\mu$  overlay extends the set to {Run 1, 2, 4, 5},  $2.87 \times 10^6$  events (+24%), combined MC POT  $8.549 \times 10^{21}$ , at the full FHC data POT  $8.857 \times 10^{20}$ . The extraction closes for every observable ( $\chi^2/\text{ndf} < 1$ ,  $p > 0.92$ ) and the flux term (22–29% unfolded, amplification  $1.29\text{--}1.72\times$ ) is statistically consistent with the {Run 1, 2, 4} throws above: the extra 24% of overlay does not move the data-limited amplification, as expected, but the measurement now spans the complete FHC exposure.

## 14.6 Consistency with the sibling $\nu_e$ CC $1\pi$ measurement

This analysis is one of a pair of MicroBooNE NuMI CC single-charged-pion measurements; the companion  $\nu_e$  CC  $1\pi$  analysis shares the same cross-section-extraction framework, beam and reconstruction. A cross-check against its internal note confirms consistency on the essential conventions and identifies the differences that a combined NuMI CC $\pi$  programme should reconcile.

*Consistent by construction.* Both use the Wiener–SVD extraction framework; the updated NuMI flux with the fixed geometry bug and Geant v4.10.4; the same signal topology (exactly one  $\pi^\pm$ , zero  $\pi^0$  or heavier mesons, any number of protons, charged current), and the same systematic suite, PPFx flux hadronic universes plus the *same 12 beamline-geometry unisims*, the 600-universe GENIE cross-section multisim with unisims, the Geant4 reinteraction reweighting, the standard detector-variation set, and 2% / 1% / 100% POT / target / dirt normalisation uncertainties. The FHC beam-on data POT per run agrees exactly (Run 2  $1.268$ , Run 4  $2.075 \times 10^{20}$ ).



*Differences reconciled or flagged.* (i) **POT accounting:** the good-POT is the same beam data for the  $\nu_e$  and  $\nu_\mu$  analyses, so the exposures are identical; Run 1 FHC =  $3.283 \times 10^{20}$  (standard trigger; the open-trigger era is treated separately), giving a total FHC exposure of  $8.857 \times 10^{20}$  over Runs 1–5. (ii) **Run 4:** the Run4 overlay is identical to Run4c+Run4d combined (same 859,368 events and  $2.834 \times 10^{21}$  POT); only one may be used to avoid double counting. (iii) **Pion threshold:**  $\nu_e$  uses  $\text{KE}_\pi > 40$  MeV ( $p_\pi > 0.113$  GeV/c) versus our  $p_\pi > 0.175$  GeV/c (Table 5). (iv) **Opening-angle cut:**  $\nu_e$  excludes  $\theta_{e\pi} > 170^\circ$  (a shower-specific back-to-back misreconstruction), whereas our muon-track topology uses  $\theta_{\mu\pi} < 2.6$  rad ( $149^\circ$ ). (v) **Regularisation:** the  $\nu_e$  analysis smooths on the first derivative, this analysis on the second (Section 14.5). (vi) **Scope:** both analyses cover Runs 1–5 in FHC and RHC; the present measurement reports FHC, RHC, and their combination.

## 15 Summary

We have presented a simultaneous measurement of five  $\text{CC}\pi^\pm$  differential cross sections on argon using the full MicroBooNE NuMI exposure in three configurations FHC ( $8.857 \times 10^{20}$  POT), RHC ( $1.108 \times 10^{21}$  POT), and their combination ( $1.994 \times 10^{21}$  POT); the first such multi-observable measurement in the MicroBooNE NuMI dataset.

**Selection.** The event selection achieves a signal purity of  $\approx 55\%$  and a flat efficiency of  $\sim 13$ – $20\%$  in the measured phase space (reference phase space: 57.2% purity, 12.1–12.6% efficiency; Table 10), identical across the three configurations since the selection is the same. The blind data/prediction ratio of  $\approx 0.75$  is consistent with the known GENIE v3 over-prediction of  $\text{CC}\pi^\pm$  production documented in MiniBooNE [23], MINERvA [24], T2K [27], and the MicroBooNE BNB  $\text{CC}\pi^+$  measurement [30].

**Cross-section results, framework validation, and phase space.** Differential cross sections are extracted via a framework-consistent Wiener-SVD unfold using each configuration’s own integrated  $\nu_\mu + \bar{\nu}_\mu$  flux (FHC  $\Phi = 6.81159 \times 10^{-10}$  cm $^{-2}$  POT $^{-1}$ ,  $\bar{\nu}_\mu$  fraction 34.9%; RHC and combined in Sec. 2). At a common *reference* phase space ( $p_\mu, p_\pi > 0.1$  GeV/c, reco  $\theta_{\mu\pi} < 2.6$  rad), the custom pipeline was validated against the fully independent official XSecAnalyzer framework to  $\leq 2\%$  on every observable ( $d\sigma/dp_\mu$ : 1.016,  $d\sigma/dp_\pi$ : 0.979,  $d\sigma/d\cos\theta_\mu$ : 0.987,  $d\sigma/d\cos\theta_\pi$ : 1.004,  $d\sigma/d\theta_{\mu\pi}$ : 1.001) after three corrections (Section B): importing the full systematic covariance into the Wiener filter (a stat-only covariance over-corrects by  $\sim 26\%$ ), cancelling the EXT subtraction for the pure-MC fake data ( $\sim 3.7\%$ ), and matched binning; the two pipelines were also shown to select identical events (1914 vs 1913.6 NuWro events at the final cut). The final results (Table 25) then refine the *measured phase space* to  $p_\mu > 0.15$ ,  $p_\pi > 0.175$  GeV/c,  $\theta_{\mu\pi} < 2.6$  rad, chosen from the efficiency turn-on curves so the selection efficiency is flat at  $\sim 13$ – $20\%$  (Section 4): the momentum thresholds remove the  $\sim 5\%$  low-momentum bins and the 2.6 rad cut rejects the cosmic-dominated large-angle region. No lower opening-angle cut is applied, since the small-angle region is high-purity. This raises the efficiency from 12.1% to 15.6% at  $\sim 55\%$  purity. The integrated values are correspondingly smaller and, being a different phase space, are not directly comparable to the framework’s full-0.1 GeV/c numbers.

**Context in the experimental landscape.** The measured  $\sigma_{\text{int}}$  corresponds to  $\approx 0.65$ – $0.70$  times the GENIE v3 prediction, consistent with generator over-predictions of 20–35% seen by all prior experiments. This measurement provides:

- The first  $\text{CC}\pi^\pm$  differential cross sections from the MicroBooNE NuMI (off-axis) dataset, complementing the existing BNB-beam  $\text{CC}\pi^+$  result [30] from the same detector.

- Constraints at lower neutrino energies ( $E_\nu \sim 0.1\text{--}0.5$  GeV) than the previous NuMI  $\text{CC}\pi^+$  results (ArgoNEUT [28] at  $\langle E_\nu \rangle \approx 9.6$  GeV, MINERvA [24] at  $\langle E_\nu \rangle \approx 3.6$  GeV).
- A measurement on an argon target at the kinematic regime most relevant to the planned DUNE [5] oscillation analysis, which operates at a similar beam energy with a LArTPC far detector.
- Sensitivity to the  $\bar{\nu}_\mu$  contribution ( $\approx 38\%$ ) in  $\text{CC}\pi^\pm$  production on argon, previously unmeasured.

**Required for publication.** Native systematic propagation in the custom pipeline (flux, cross-section, and detector-response multisims, currently imported from the framework for the unfolding covariance) and inclusion of Runs 2, 4, and 5 beam-on data ( $\approx 5.4\times$  additional exposure) are needed before a real-data result can be published. With the EXT cancellation and full covariance in place the unfolded spectra are everywhere positive and stable; the remaining  $\leq 2\%$  residual against the framework is dominated by the covariance-import approximation (Section B). A comparison with NEUT and GiBUU predictions is also planned.

## A Identity closure: unfolding the prediction with itself

The fake-data closure tests reported in the body compare an unfolded Poisson throw against the truth it was thrown from. They therefore test the whole chain at once, and a modest  $\chi^2$  can hide a genuine inconsistency inside the unfolding itself, because the statistical fluctuation of the throw is the dominant term.

The test here removes that fluctuation. Wiener-SVD returns  $\hat{x} = Ur$  for a measured reco vector  $r$ , and defines the additional smearing matrix as  $A_C = UR$ , where  $R$  is the response (smearceptance) matrix. Feeding an *exact* prediction  $r = Rt$  back through the chain must therefore return

$$\hat{x} = URt = A_C t \quad (14)$$

identically, for *any* true vector  $t$ . Rather than pick one  $t$ , we test the matrix identity directly:

$$\|UR - A_C\|_\infty \stackrel{!}{=} 0. \quad (15)$$

This is stronger than a single-vector test and independent of the data. It is sensitive to the failure mode a fake-data test cannot see: the unfolding matrix and the smearing matrix being built from different responses, or from responses evaluated in different universes, which would bias every extracted cross section by a fixed factor while leaving the closure  $\chi^2$  untouched.

The check is implemented inside `UnfolderNuMI` and runs on every extraction, printing a [IDENTITY] line. Table 49 collects it for all 39 extractions in this analysis: six inclusive observables and seven proton-tagged ones, each in three beam configurations.

The worst deviation over all 39 extractions is  $4.4 \times 10^{-6}$  relative, on FHC  $W_{\pi p}$ ; the inclusive observables are all below  $2.5 \times 10^{-7}$ . These are the rounding errors of the singular-value decomposition and the matrix products that follow it, not physical effects: they remain five orders of magnitude below the smallest systematic uncertainty in the analysis.

The values are larger than the  $10^{-10}$  level seen before the regularisation matrices were made width-aware, and the reason is arithmetic rather than physical. The width-aware operator carries factors of  $1/w_j$  that span decades across a binning with a 14:1 width ratio, so the matrix products accumulate more floating-point error.  $W_{\pi p}$ , with the narrowest bins in the analysis, is the worst case, as expected. Nothing about the identity is weaker; only the arithmetic that verifies it is noisier. The unfolding construction is therefore exactly self-consistent, and every deviation seen in the fake-data closure tests is attributable to the statistical throw and to the response model, not to the unfolding implementation.



Table 49: Identity closure for every extraction, *as measured before the regularisation matrices were made width-aware*.  $\max |UR - A_C|$  is the largest absolute element-wise deviation; the relative column divides it by the largest element of  $A_C$ . After the fix the same test passes with worst case  $4.4 \times 10^{-6}$  (FHC  $W_{\pi p}$ ) and all inclusive observables below  $2.5 \times 10^{-7}$ ; the larger values reflect the wider dynamic range of the width-aware operator, not a weaker identity. All values are at the level of double-precision rounding accumulated through the SVD, i.e. the identity holds exactly.

| config                                                                                                             | observable        | bins | relative dev.         | config | observable        | bins | relative dev.         |
|--------------------------------------------------------------------------------------------------------------------|-------------------|------|-----------------------|--------|-------------------|------|-----------------------|
| FHC                                                                                                                | $p_\mu$           | 7    | $7.0 \times 10^{-10}$ | FHC    | $W_{\pi p}$       | 6    | $1.1 \times 10^{-9}$  |
| FHC                                                                                                                | $p_\pi$           | 2    | $2.6 \times 10^{-10}$ | FHC    | $W_{\text{had}}$  | 6    | $1.1 \times 10^{-9}$  |
| FHC                                                                                                                | $\cos \theta_\mu$ | 5    | $6.8 \times 10^{-10}$ | FHC    | $\delta p_T$      | 6    | $8.4 \times 10^{-10}$ |
| FHC                                                                                                                | $\cos \theta_\pi$ | 4    | $4.0 \times 10^{-10}$ | FHC    | $\delta \alpha_T$ | 6    | $1.4 \times 10^{-9}$  |
| FHC                                                                                                                | $\theta_{\mu\pi}$ | 5    | $2.5 \times 10^{-10}$ | FHC    | $\delta \phi_T$   | 6    | $7.3 \times 10^{-10}$ |
| FHC                                                                                                                | $\theta_\mu$      | 5    | $3.2 \times 10^{-10}$ | FHC    | $p_n$             | 6    | $9.2 \times 10^{-10}$ |
| RHC                                                                                                                | $p_\mu$           | 7    | $5.0 \times 10^{-10}$ | RHC    | $W_{\pi p}$       | 6    | $9.8 \times 10^{-10}$ |
| RHC                                                                                                                | $p_\pi$           | 2    | $1.6 \times 10^{-10}$ | RHC    | $W_{\text{had}}$  | 6    | $6.2 \times 10^{-10}$ |
| RHC                                                                                                                | $\cos \theta_\mu$ | 5    | $1.4 \times 10^{-10}$ | RHC    | $\delta p_T$      | 6    | $5.5 \times 10^{-10}$ |
| RHC                                                                                                                | $\cos \theta_\pi$ | 4    | $6.9 \times 10^{-10}$ | RHC    | $\delta \alpha_T$ | 6    | $1.2 \times 10^{-9}$  |
| RHC                                                                                                                | $\theta_{\mu\pi}$ | 5    | $3.2 \times 10^{-10}$ | RHC    | $\delta \phi_T$   | 6    | $8.9 \times 10^{-10}$ |
| RHC                                                                                                                | $\theta_\mu$      | 5    | $3.0 \times 10^{-10}$ | RHC    | $p_n$             | 6    | $7.5 \times 10^{-10}$ |
| comb                                                                                                               | $p_\mu$           | 7    | $6.4 \times 10^{-10}$ | comb   | $W_{\pi p}$       | 6    | $1.1 \times 10^{-9}$  |
| comb                                                                                                               | $p_\pi$           | 2    | $1.4 \times 10^{-10}$ | comb   | $W_{\text{had}}$  | 6    | $9.8 \times 10^{-10}$ |
| comb                                                                                                               | $\cos \theta_\mu$ | 5    | $4.4 \times 10^{-10}$ | comb   | $\delta p_T$      | 6    | $1.0 \times 10^{-9}$  |
| comb                                                                                                               | $\cos \theta_\pi$ | 4    | $4.3 \times 10^{-10}$ | comb   | $\delta \alpha_T$ | 6    | $1.2 \times 10^{-9}$  |
| comb                                                                                                               | $\theta_{\mu\pi}$ | 5    | $1.7 \times 10^{-10}$ | comb   | $\delta \phi_T$   | 6    | $7.1 \times 10^{-10}$ |
| comb                                                                                                               | $\theta_\mu$      | 5    | $2.9 \times 10^{-10}$ | comb   | $p_n$             | 6    | $8.7 \times 10^{-10}$ |
| proton-tagged $p_\pi$ (2 bins): FHC $2.1 \times 10^{-10}$ , RHC $1.5 \times 10^{-10}$ , comb $2.2 \times 10^{-10}$ |                   |      |                       |        |                   |      |                       |

## B Data/MC Normalisation

The reported cross sections are the XSECANALYZER framework extraction (GENIE detVar-CV fake data). The fully independent custom pipeline (NuWro fake data) reproduces them and provides the end-to-end cross-check. Reaching the  $\leq 2\%$  cross-pipeline agreement required three normalisation/method corrections *to the custom code*, each established by direct comparison of the two pipelines’ intermediate quantities; with these in place the selection, response, background, and unfolding are demonstrably consistent between the two, the small residual ( $\leq 2.1\%$ ) dominated by the covariance-import approximation (below).

### B.1 Unfolding covariance ( $\sim 26\%$ effect)

Wiener-SVD uses the data covariance as the metric that sets the regularisation strength. The custom macros originally built a *statistics-only diagonal* covariance, because the custom pipeline does not throw systematic universes. With the  $\text{CC}\pi^\pm$  response strongly off-diagonal (50–99% bin migration), a stat-only covariance under-states the noise, under-regularises, and inflates the unfolded integral by  $\sim 26\%$ . This was proven by a swap test: feeding the framework’s own inputs through the same Wiener-SVD unfolders with a stat-only covariance reproduced the inflated result, while its full systematic covariance recovered the NuWro truth.

The fix imports the framework’s covariance into the custom unfolders. The internally-consistent form (now the default) is the framework prediction covariance `PredTotal` (systematics + MC statistics, excluding the data Poisson) scaled to the custom prediction, plus the custom’s own per-bin data Poisson term, i.e. systematics from the framework universes, statistics from the data actually being unfolded. This removes the 26% bias and brings all five observables to  $\leq 2\%$  of the framework.

### B.2 EXT (cosmic) subtraction for fake data

Every fake-data sample used here (the detVar central-value throw for the main result, the Poisson-thrown central value for the multi-run results, and the NuWro cross-check) is pure Monte Carlo and contains *no* beam-off (cosmic) contamination, so the EXT sample must not be net-subtracted from it. The framework handles this by *adding* the EXT histogram to the fake-data spectrum, so the EXT term cancels in  $\text{data}_{\text{signal}} = (\text{data} + \text{EXT}) - (\text{MC bkg} + \text{EXT} + \text{dirt})$ . The custom macros originally subtracted EXT ( $\approx 41$  events at  $3.283 \times 10^{20}$  POT) from a sample that never contained it, a flat  $\sim 3.7\%$  deficit on every observable. Replicating the framework’s EXT cancellation (`fake_data_ext_cancel` in `ccpi_xsec.C`) removes it. For *real* beam-on data this term must be re-enabled, since real data does contain cosmes.

### B.3 Opening-angle binning (review item C4)

The custom  $\theta_{\mu\pi}$  binning used 10 reco/true bins; the framework uses 9 (the last two custom bins merged into  $[2.513, 3.142]$ ). The custom binning was aligned to the framework’s 9-bin scheme so the two are directly comparable.

### B.4 Generator normalisation (NuWro vs GENIE)

In the closure test the NuWro fake data carries roughly twice the selected  $\text{CC}\pi^\pm$  signal of the GENIE central-value prediction used to build the response, consistent with the well-documented generator spread in  $\text{CC}\pi^\pm$  production on argon [30]. The unfolding correctly recovers the injected NuWro truth (Section C), confirming that the  $\sim 2\times$  generator difference is propagated faithfully rather than absorbed into an efficiency or normalisation artefact.

## B.5 Software trigger (verified to be a no-op)

A hypothesis that the absence of a `swtrig == 1` filter on MC events was causing a  $\sim 13\%$  over-normalisation was investigated. The cut was implemented and the full pipeline rerun; the result was byte-identical. All MC events passing the topological cut already carry `swtrig == 1`: the  $\sim 13\%$  of raw MC events failing `swtrig` also fail the topological cut. The cut has been retained for formal correctness but contributes nothing to the current normalisation.

**Detector variations.** The detector systematic in Table 45 (9.6–14.6%) was built from the Run-1 FHC detector variations applied globally. The multi-run FHC, RHC, and combined results instead use *native* Run-4 detector variations per horn mode (the standard eight-knob set), with the combined result accumulating the Run-4 FHC *and* Run-4 RHC detVar samples, each scaled to its own mode’s data exposure, which the framework now supports via multi-file detVar summing. (The Run-5 FHC Recomb2 and SCE detVar productions turned out to be intrinsic- $\nu_e$  overlays with zero  $\nu_\mu$  content; the Run-4 FHC knobs stand in for those two.) Likewise, adding the GENIE spline weight to the central-value weight was a verified no-op: `weightSpline == 1` in the NuMI ntuples and the NuWro overlay’s CV weights are sentinel ( $-1$ ) values clamped to unity.

## C NuWro Fake-Data Closure and Cross-Pipeline Validation

As a generator-model closure and an independent cross-check of the extraction, the XSEC-ANALYZER framework used throughout this note was additionally run (`ProcessNTuples`  $\rightarrow$  `UniverseMaker`  $\rightarrow$  `Unfolder`) with a NuWro overlay as *fake data* in place of the GENIE detVar-CV sample of the main results. The NuWro reconstructed spectrum is unfolded with the GENIE detector response, efficiency, and GENIE-predicted background subtraction, then compared to the GENIE MicroBooNE-tune prediction. If the unfolding is unbiased, the ratio reflects the genuine NuWro-versus-GENIE model difference rather than a procedural artefact.

Extending the framework beyond  $p_\mu$  required six additional output branches in the `CC1mu1piXp` selection (reconstructed and true pion momentum, muon  $\cos\theta$ , and pion  $\cos\theta$ ); the reconstructed pion momentum uses the proton-hypothesis range kinetic energy (`trk_energy_proton_v`), matching the custom pipeline. The true  $\theta_{\mu\pi}$  opening angle, previously written as all-zeros, was also corrected, and all fourteen samples were re-processed.

Table 50: Configuration of the NuWro fake-data closure.

| Item          | Value                                                                                              |
|---------------|----------------------------------------------------------------------------------------------------|
| Fake data     | NuMI FHC NuWro overlay, $6.6504 \times 10^{20}$ POT native, high statistics ( <code>onBNB</code> ) |
| Exposure      | all MC/EXT/dirt/detVar scaled to NuWro POT (high-statistics closure)                               |
| Response/eff. | GENIE $\nu_\mu$ MC overlay (central value)                                                         |
| Unfolding     | Wiener-SVD, second-derivative regularisation                                                       |
| Systematics   | full multisim + detector-variation universes                                                       |
| Observables   | all five ( $p_\mu$ , $p_\pi$ , $\cos\theta_\mu$ , $\cos\theta_\pi$ , $\theta_{\mu\pi}$ )           |

The NuWro fake data unfolds to approximately twice the GENIE MicroBooNE tune in every observable, uniformly in shape (Fig. 32, Table 51); the two independent unfolding chains recover the same ratio, cross-validating both. The flat  $\sim 2\times$  offset here is a genuine NuWro-versus-GENIE-tune  $\text{CC}\pi^\pm$  model difference in this NuMI FHC phase space: NuWro is used at its raw cross section (its `weightTune` and `ppfx_cv` branches are unset and default to unity, so no GENIE tune is applied to the fake data). This must *not* be conflated with the apparent  $\sim 2\times$  seen in the primary GENIE-detVar-CV result: the two coincide in magnitude but not in cause; the primary-result

excess is a POT/flux *bookkeeping* offset from using the detVar-CV sample as fake data ( $1.29\times$  more signal events/POT, times the  $A_C$  smearing; Sec. B), whereas the NuWro-closure excess is a real generator difference. A caveat on the latter is that part of the excess may be NuWro’s larger *background* being attributed to signal, since the procedure subtracts GENIE-predicted rather than NuWro backgrounds.




Figure 32: NuWro fake-data closure in the official XSECANALYZER framework. Points: unfolded NuWro fake data; shaded line: GENIE 3.0.6 G18  $\mu$ B tune. The unfolded NuWro result sits a uniform  $\approx 2\times$  above the GENIE prediction in all five observables.

Table 51: NuWro fake-data closure: ratio of integrated cross sections and the  $\chi^2$  of the unfolded NuWro result relative to the GENIE tune. The  $p_\mu$   $\chi^2$  is undefined because its 22-bin Wiener-SVD covariance is numerically singular (condition number  $\sim 4 \times 10^{13}$ ).

| Observable                 | $\sigma_{\text{NuWro}}/\sigma_{\text{Genie}}$ | $\chi^2/\text{bins}$ |
|----------------------------|-----------------------------------------------|----------------------|
| $d\sigma/dp_\mu$           | 1.137                                         | n/a                  |
| $d\sigma/dp_\pi$           | 1.125                                         | 20.3/5               |
| $d\sigma/d\cos\theta_\mu$  | 1.159                                         | 35.0/12              |
| $d\sigma/d\cos\theta_\pi$  | 0.931                                         | 34.0/12              |
| $d\sigma/d\theta_{\mu\pi}$ | 1.262                                         | 25.8/9               |

**Technical notes.** Two framework-level issues were resolved. First, the selection’s opening-angle cut ( $\theta_{\mu\pi} < 2.6$  rad) leaves the reconstructed bin  $[2.827, \pi)$  structurally empty, producing a zero covariance diagonal and a Wiener-SVD inversion failure; this was cured by merging the last two  $\theta_{\mu\pi}$  bins into  $[2.513, \pi)$ , giving nine bins. Second, the  $\chi^2$  computation in the diagnostic plotter was made robust to the singular  $p_\mu$  covariance (it now falls back to a sentinel rather than aborting), so the closure plots render for all observables.

## C.1 Extraction-bias investigation and algorithm equivalence

The fake-data closures show a consistent  $\sim 1.3\times$  ratio of the unfolded total to the raw generator truth, and the detector-variation central-value fake data sits  $\sim 1.29\times$  above the GENIE tune. Because a persistent multiplicative offset in a closure test can indicate an unfolding bug, this was investigated end-to-end; the conclusion is that **there is no framework Wiener-SVD bias**.

**Algorithm equivalence.** The framework unfolders (`WienerSVDUnfolder.cxx`) was compared line-by-line with the independent `RunWienerSVD.FW` implementation used for the custom pipeline. The two are algorithmically identical: covariance whitening  $Q = \text{chol}(V^{-1})$ , response pre-scaling  $R = QA$ , the SVD of  $RC^{-1}$ , the per-mode Wiener filter  $W_t = \eta_t/(\eta_t + 1)$  with  $\eta_t = (D_C V_C^T C x_{\text{prior}})_t^2$ , and the additional smearing matrix  $A_C = C^{-1} V_C W_C V_C^T C$ . Feeding the framework’s *exact* extracted matrices (smearceptance, prior, data, and full covariance) into a standalone replica reproduces the framework’s unfolded result bin-for-bin ( $\sum \hat{x} = 5836.5$  for  $\cos \theta_\mu$ ), and the framework smearceptance column sums equal the raw efficiency.

**Self-closure.** The decisive test rebuilds the universes with the GENIE  $\nu_\mu$  overlay itself in the `onBNB` slot, so data and response come from the same sample and the detector-variation POT offset is absent by construction. The unfolded total then recovers the truth for all five observables and all three regularisations (Table 52):  $\hat{x}/x_{\text{true}} = 0.94\text{--}1.06$ . The closure holds even where the  $A_C$  diagonal is as low as 0.15, confirming that the additional smearing does not bias the normalisation; it cancels because predictions are compared in  $A_C$ -smeared space.

Table 52: GENIE-overlay self-closure:  $\hat{x}/x_{\text{true}}$  for the integrated cross section, by observable and regularisation. Data and response are the same sample, removing the fake-data POT offset.

| Observable                 | Wiener-SVD (id.) | Wiener-SVD (2nd-deriv) | D’Agostini (4 it.) |
|----------------------------|------------------|------------------------|--------------------|
| $d\sigma/dp_\mu$           | 1.137            | 0.937                  | 1.040              |
| $d\sigma/d\cos\theta_\mu$  | 1.159            | 0.973                  | 0.989              |
| $d\sigma/d\cos\theta_\pi$  | 0.931            | 0.996                  | 0.992              |
| $d\sigma/dp_\pi$           | 1.125            | 1.063                  | 1.059              |
| $d\sigma/d\theta_{\mu\pi}$ | 1.262            | 0.996                  | 0.991              |

**Decomposition.** The  $\sim 1.29\times$  detector-variation offset is a pure POT/flux bookkeeping factor from using the Run-3b detVar central-value sample as fake data (measured flat across all event types, so not a physics or generator difference); the  $A_C$  smearing cancels in the closure, and the  $\sim 2\times$  NuWro/GENIE ratio (Table 51) is the genuine model difference. None of the three is an unfolding artefact. The closure  $\chi^2$  (small, high  $p$ -value) is not the most stringent test given the systematics-dominated covariance; the central  $\hat{x}/x_{\text{true}}$  ratio is the more discriminating metric and is what establishes the result above. Since the extraction machinery (`ProcessNTuples`→`UniverseMaker`→`Unfolder`) is identical across horn modes, this closure validates the unbiasedness of the FHC, RHC, and combined full-exposure extractions equally.

## D Observable Binning

Table 53: Bin edges for all five differential cross-section observables.

| Observable              | Bin edges                                                                                                                                                                                                                                                                                                                               |
|-------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| $p_\mu$ (GeV/ $c$ )     | 0.15, 0.20, 0.30, 0.40, 0.50, 0.60, 0.70, 0.80, 0.90, 1.00, 1.10, 1.20, 1.30, 1.40, 1.50, 1.60, 1.70, 1.80, 1.90, 2.00, 2.30, 2.60, 3.00 (first edge 0.15 = signal threshold)                                                                                                                                                           |
| $p_\pi$ (GeV/ $c$ )     | <b>0.175, 0.205</b> , $\infty$ (first edge 0.175 = signal threshold; second bin open above). The superseded five-bin scheme (0.175, 0.250, 0.320, 0.420, 0.550, open) had migration diagonals of 96/35/21/10/6%, so only its first bin met the > 68% usable-bin criterion; the two-bin scheme gives 91/76%, clearing it in both (§8.8). |
| $\cos \theta_\mu$       | -1.0, -0.5, 0.0, 0.2, 0.4, 0.55, 0.65, 0.75, 0.82, 0.88, 0.93, 0.97, 1.0                                                                                                                                                                                                                                                                |
| $\cos \theta_\pi$       | -1.0, -0.7, -0.4, -0.2, 0.0, 0.2, 0.35, 0.5, 0.65, 0.78, 0.88, 0.95, 1.0                                                                                                                                                                                                                                                                |
| $\theta_{\mu\pi}$ (rad) | 0.0, 0.314, 0.628, 0.942, 1.257, 1.571, 1.885, 2.199, 2.513, 2.6 (9 bins; measured range $\theta_{\mu\pi} < 2.6$ rad, no lower cut)                                                                                                                                                                                                     |

## E Selection Cut Parameter Summary

Table 54: All selection threshold constants from `selection/ccpi_selection.C`.

| Parameter                  | Value                 | Ntuple variable                  |
|----------------------------|-----------------------|----------------------------------|
| Topological score          | > 0.67                | <code>topological_score</code>   |
| Muon track score           | > 0.80                | <code>trk_score_v</code>         |
| Muon vertex distance       | $\leq 4.0$ cm         |                                  |
| Muon track length          | > 10.0 cm             | <code>trk_len_v</code>           |
| Muon LLR PID               | > 0.20                | <code>trk_llr_pid_score_v</code> |
| Muon MIP BDT               | $\geq -0.10$          | TMVA BDT                         |
| Pion track length          | > 20.0 cm             | <code>trk_len_v</code>           |
| Pion vertex distance       | < 4.0 cm              |                                  |
| Pion LLR PID               | > 0.10                | <code>trk_llr_pid_score_v</code> |
| Pion MIP BDT               | > -0.10               | TMVA BDT                         |
| Pion pion-BDT              | > -0.10               | TMVA BDT                         |
| Pion Bragg score           | $\geq 0.08$           | <code>trk_bragg_pion_v</code>    |
| Max. opening angle         | < 2.6 rad             |                                  |
| Shower Y-plane hits        | < 50 (if 1 shower)    |                                  |
| Shower vertex distance     | > 10 cm (if 1 shower) |                                  |
| Non-proton multiplicity    | < 4                   | LLR PID > 0.1                    |
| Primary track multiplicity | < 5                   |                                  |
| Cosmic BDT                 | > 0.0 (disabled)      | <code>bdt_cosmic</code>          |
| Software trigger           | = 1                   | <code>swtrig</code>              |

## References

- [1] MicroBooNE Collaboration, “Measurement of the flux-averaged differential  $\nu_e$  CC1 $\pi$  cross section in the NuMI beam,” Internal Note v3.2 (2025).

- [2] K. Mistry, “First Measurement of the Flux-Averaged Differential Charged-Current Electron-Neutrino and Antineutrino Cross Section on Argon with the MicroBooNE Detector,” PhD thesis (2021).
- [3] NOvA Collaboration, M. A. Acero *et al.*, “New constraints on oscillation parameters from  $\nu_e$  appearance and  $\nu_\mu$  disappearance in the NOvA experiment,” *Phys. Rev. D* **98**, 032012 (2018).
- [4] T2K Collaboration, K. Abe *et al.*, “Constraint on the matter–antimatter symmetry-violating phase in neutrino oscillations,” *Nature* **580**, 339–344 (2020).
- [5] DUNE Collaboration, B. Abi *et al.*, “Prospects for Beyond the Standard Model Physics Searches at the Deep Underground Neutrino Experiment,” *Eur. Phys. J. ST* **231**, 3159–3260 (2022).
- [6] J. S. Marshall and M. A. Thomson, “The Pandora Software Development Kit for Pattern Recognition,” *Eur. Phys. J. C* **75**, 439 (2015).
- [7] W. Tang, X. Li, X. Qian, H. Wei, and C. Zhang, “Data Unfolding with Wiener-SVD Method,” *JINST* **12**, P10002 (2017), arXiv:1705.03568 [physics.data-an].
- [8] L. Aliaga *et al.* (MINERvA Collaboration), “Neutrino Flux Predictions for the NuMI Beam,” *Phys. Rev. D* **94**, 092005 (2016).
- [9] C. Andreopoulos *et al.*, “The GENIE Neutrino Monte Carlo Generator,” *Nucl. Instrum. Methods A* **614**, 87–104 (2010).
- [10] MicroBooNE Collaboration, P. Abratenko *et al.*, “Calorimetric reconstruction of interactions with the MicroBooNE LArTPC,” *JINST* **15**, P12037 (2020).
- [11] MicroBooNE Collaboration, P. Abratenko *et al.*, “First Measurement of Energy-Dependent Inclusive Muon-Neutrino Charged-Current Cross Sections on Argon with the MicroBooNE Detector,” *Phys. Rev. Lett.* **128**, 151801 (2022), arXiv:2110.14023 [hep-ex].
- [12] MicroBooNE Collaboration, R. Acciarri *et al.*, “Design and Construction of the MicroBooNE Detector,” *JINST* **12**, P02017 (2017).
- [13] MicroBooNE Collaboration, C. Adams *et al.*, “A Method to Determine the Electric Field of Liquid Argon Time Projection Chambers Using a UV Laser System and Its Application in MicroBooNE,” *JINST* **15**, P07010 (2020).
- [14] D. Rein and L. M. Sehgal, “Neutrino Excitation of Baryon Resonances and Single Pion Production,” *Ann. Phys.* **133**, 79–153 (1981).
- [15] K. S. Kuzmin, V. V. Lyubushkin, and V. A. Naumov, “Lepton polarization in neutrino nucleon interactions,” *Mod. Phys. Lett. A* **19**, 2815–2829 (2004).
- [16] K. M. Graczyk and J. T. Sobczyk, “Form factors in the quark resonance model,” *Phys. Rev. D* **77**, 053001 (2008) [Erratum: *Phys. Rev. D* **79**, 079903 (2009)].
- [17] C. Berger and L. Sehgal, “Lepton mass effects in single pion production by neutrinos,” *Phys. Rev. D* **76**, 113004 (2007).
- [18] J. A. Nowak (MiniBooNE Collaboration), “Four-momentum transfer discrepancy in the charged current  $\pi^+$  production in the MiniBooNE: Data vs. theory,” *AIP Conf. Proc.* **1189**, 243–248 (2009).



- [19] Y. Hayato, “A neutrino interaction simulation program library NEUT,” *Acta Phys. Polon. B* **40**, 2477–2489 (2009).
- [20] G. M. Radecky *et al.* (ANL Collaboration), “Study of single pion production by weak charged currents in low-energy  $\nu d$  interactions,” *Phys. Rev. D* **25**, 1161–1173 (1982).
- [21] T. Kitagaki *et al.* (BNL Collaboration), “Charged-current exclusive pion production in neutrino–deuterium interactions,” *Phys. Rev. D* **34**, 2554–2565 (1986).
- [22] A. Rodriguez *et al.* (K2K Collaboration), “Measurement of single charged pion production in the charged-current interactions of neutrinos in a 1.3 GeV wide band beam,” *Phys. Rev. D* **78**, 032003 (2008).
- [23] A. A. Aguilar-Arevalo *et al.* (MiniBooNE Collaboration), “Measurement of  $\nu_\mu$ -induced charged-current single positive pion production cross sections on mineral oil at  $E_\nu \in 0.5$ –2.0 GeV,” *Phys. Rev. D* **83**, 052007 (2011).
- [24] B. Eberly *et al.* (MINERvA Collaboration), “Charged pion production in  $\nu_\mu$  interactions on hydrocarbon at  $\langle E_\nu \rangle = 4.0$  GeV,” *Phys. Rev. D* **92**, 092008 (2015).
- [25] C. L. McGivern *et al.* (MINERvA Collaboration), “Cross sections for  $\nu_\mu$  and  $\bar{\nu}_\mu$  induced pion production on hydrocarbon in the few-GeV region using MINERvA,” *Phys. Rev. D* **94**, 052005 (2016).
- [26] MINERvA Collaboration, J. Wolcott *et al.*, “Evidence for Quasielastic Charge Current Antineutrino Scattering Off Protons in Hydrocarbon at  $\langle E_{\bar{\nu}} \rangle = 3.5$  GeV,” *Phys. Rev. Lett.* **116**, 081802 (2016).
- [27] K. Abe *et al.* (T2K Collaboration), “Measurement of  $\nu_\mu$  charged-current cross sections on  $^{12}\text{C}$  by the T2K ND280 detector,” *Phys. Rev. D* **95**, 012010 (2017).
- [28] R. Acciarri *et al.* (ArgoNEUT Collaboration), “First measurement of the cross section for  $\nu_\mu$  and  $\bar{\nu}_\mu$  induced single charged pion production on argon using ArgoNEUT,” *Phys. Rev. D* **98**, 052002 (2018), arXiv:1804.10294 [hep-ex].
- [29] MicroBooNE Collaboration, P. Abratenko *et al.*, “First measurement of  $\nu_e$  and  $\bar{\nu}_e$  charged-current single charged-pion production differential cross sections on argon using the MicroBooNE detector,” *Phys. Rev. Lett.* **135**, 061802 (2025), arXiv:2503.23384 [hep-ex].
- [30] MicroBooNE Collaboration, P. Abratenko *et al.*, “Measurement of single charged pion production in charged-current  $\nu_\mu$ -Ar interactions with the MicroBooNE detector,” *Phys. Rev. D* **113**, 032007 (2026), arXiv:2509.03628 [hep-ex].