

NOTES ON A NONCONFORMING FEM

JESSE CHAN

1. A hybrid nonconforming method. Consider a coercive variational formulation

$$a(u, v) = l(v), \quad \forall v \in V.$$

We assume further that $a(u, v)$ is coercive when restricted to a single element $u, v \in V(K)$. Standard methods approximate u using a conforming subspace of V . If we choose instead a broken test space

$$V_h := \{v \in L^2(\Omega), v|_K \in V(K)\},$$

we can enforce continuity between elements by penalizing the jumps of e using a Lagrange multiplier method

$$\begin{aligned} a_h(u, v) + \langle \lambda, \llbracket v \rrbracket \rangle_{\Gamma_h} &= l(v) \\ \langle \delta \lambda, \llbracket u \rrbracket \rangle_{\Gamma_h} &= 0. \end{aligned}$$

where $a_h(u, v) = \sum a_K(u, v)$, and λ is a function with support only on Γ_h , the mesh skeleton. The advantage of this formulation is that the degrees of freedom for e can be condensed out and eliminated, leaving a positive definite system for λ .

2. Stability analysis. We define norms on our variables u and λ :

$$\begin{aligned} \|u\|_{\Omega_h}^2 &:= \sum_{K \in \Omega_h} a_K(u, u) \\ \|\lambda\|_{\Gamma_h} &:= \min_{q \in Q, \gamma_Q(q)|_{\Gamma_h} = \lambda} \|q\|_Q. \end{aligned}$$

where λ is discretized as the trace of some space Q with trace $\gamma_Q(q)$, such that $\gamma_Q(q)$ is dual to $\gamma_V(u)$ s, the trace of u , on Γ_h . We also define an auxiliary norm on the jumps of functions in V_h

$$\|\llbracket v \rrbracket\|_{\Gamma_h} := \sup_{\lambda} \frac{\langle \lambda, \llbracket v \rrbracket \rangle_{\Gamma_h}}{\|\lambda\|_{\Gamma_h}}$$

Since the condensed system is equivalent to the mixed system, we can use Brezzi mixed theory to analyze the stability of this method. We require two conditions

- Inf-sup relating the two spaces

$$\inf \sup \frac{\langle \lambda, \llbracket u \rrbracket \rangle_{\Gamma_h}}{\|\lambda\|_{\Gamma_h} \|u\|_{\Omega_h}} \geq \gamma_0 > 0,$$

- Inf-sup in the kernel: for $u_0 \in U_0 := \{u \in V : \langle \lambda, \llbracket u \rrbracket \rangle_{\Gamma_h} = 0\}$

$$\inf \sup \frac{a_h(u_0, v)}{\|u_0\|_{\Omega_h} \|v\|_{\Omega_h}} \geq \gamma_0 > 0,$$

The second condition holds trivially due to coercivity of $a_h(u, v)$ on K , while the first condition reduces to

$$\|\llbracket u \rrbracket\|_{\Gamma_h} \geq \gamma_0 \|u\|_{\Omega_h}.$$

In other words, the jumps of u should be bounded from below by the nonconforming broken norm.




FIG. 2.1. L^2 errors for conforming vs nonconforming discretizations.

2.1. Example: Poisson's equation with first order term. We consider the equation with $\alpha > 0$

$$-\Delta u + \alpha u = f$$

which gives the following conforming variational formulation

$$a(u, v) := (\nabla u, \nabla v)_{\Omega} + \alpha (u, v)_{\Omega} = (f, v)_{\Omega}.$$

The nonconforming formulation is given by

$$a_h(u, \lambda, v) := \sum_{K \in \Omega_h} [a_K(u, v) + \langle \lambda, v \rangle_{\partial K}] = \sum_{K \in \Omega_h} a_K(u, v) + \langle \lambda, \llbracket v \rrbracket \rangle_{\Gamma_h}$$

In this specific case, λ is the trace of Raviart-Thomas elements, with corresponding norm

$$\|\lambda\|_{\Gamma_h} := \min_{q \in H(\text{div}), q \cdot n|_{\Gamma_h} = \lambda} \|q\|_{H(\text{div})}.$$

The first condition then requires only a mesh-independent Poincare inequality for broken H^1 functions, given in Lemma 4.2 of [1].

$$\|v\|_{\Omega_h} \leq C (\|\nabla v\|_{\Omega_h} + \|\llbracket v \rrbracket\|_{\Gamma_h})$$

Both the primal and original C_0 mixed DPG methods for convection-diffusion display optimal rates, and display nearly identical L^2 errors on the same mesh. Below are comparisons of L^2 -errors under uniform refinement. The flux variable λ is taken to be the trace of Raviart-Thomas elements of equal order to the field variables u (which gives a polynomial degree of $p-1$ on the edge). If we decrease our polynomial degree to be Raviart-Thomas elements of order $p-1$ instead of p , the rate of convergence falls to h^p for L^2 errors.¹

REFERENCES

- [1] L. Demkowicz and J. Gopalakrishnan. Analysis of the DPG method for the Poisson equation. *SIAM J. Numer. Anal.*, 49(5):1788–1809, September 2011.

¹In general, the optimal rate of convergence appears to be limited to $p_f + 1$.

Order	Conforming	Non-conforming
$p = 1$	1.98898	2.00114
$p = 2$	3.00635	3.01681
$p = 3$	4.01242	4.00886

(a) $p_f = p$

Order	Non-conforming
$p = 1$	N/A
$p = 2$	2.00317
$p = 3$	2.98382

(b) $p_f = p - 1$

FIG. 2.2. Rates of convergence for conforming vs nonconforming discretizations.