

Homework 1 Problems

(1) Define the following terms:

- (a) Population
- (b) Sample
- (c) Parameter
- (d) Statistic

(2) Explain the difference between a **qualitative** (categorical) and a **quantitative** variable

(3) Explain the difference between a **discrete** and a **continuous** variable and give an example of each

(4) Explain the difference between a **nominal** and an **ordinal** variable and give an example of each

(5) **At What age did women marry?** A historian wants to estimate the average age at marriage of women in New England in the early 19th century. Within her state archives, she finds marriage records for the years 1800 – 1820, which she treats as a sample of all marriage records from the early 19th century. The average age of the women in the records is 24.1 years of age. Using the appropriate statistical method, she estimates that the average age of brides in the early 19th- century New England was between 23.5 and 24.7 years of age.

- (a) Which part of this example gives a descriptive summary of the data?
- (b) Which part of this example draws an inference about a population?
- (c) What population is the historian studying?
- (d) The average age the historian computed from the historical records was 24.1 years of age. Is 24.1 years of age a statistic or a parameter? Why?

(6) Consider the following data obtained from flipping a coin 15 times. A value of H denotes the coin was heads whereas a value of T denotes the coin flip was tails.

##	Flip	Result
## 1	1	H
## 2	2	H
## 3	3	H
## 4	4	T
## 5	5	H
## 6	6	T
## 7	7	T
## 8	8	T
## 9	9	H
## 10	10	H
## 11	11	T
## 12	12	T
## 13	13	T
## 14	14	H
## 15	15	T

coin.flip.result.	H	H	H	T	H	T	T	T	H	H	T	T	T	H	T
-------------------	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---

Compute the frequency table for the variable “coin flip result” and answer the following questions:

Result	Frequency	Relative Frequency
H		
T		

- (a) What type of variable is “Coin Flip Result”?
- (b) What proportion of the coin flips were heads?
- (c) Why do we not compute the cumulative relative frequency for this variable?
- (d) What is the best graphical display for this variable and why?

(7) A survey about color preferences reported the age distribution of the people who responded. Below are the results

##	Age Group (Years)	Counts
## 1	1-18	10
## 2	19-24	97
## 3	25-35	70
## 4	36-50	36
## 5	51-69	14
## 6	70 and over	5

1	Age Group (Years)	1-18	19-24	25-35	36-50	51-69	70 and over
2	Counts	10	97	70	36	14	5

Use this table to answer parts a - d

- (a) Compute the relative frequency for each age group
- (b) Make a bar graph where the heights of the bars are relative frequencies
- (c) Describe the distribution
- (d) Explain why your bar graph is not a histogram

(8) Email spam is the curse of the internet. The table below gives a compilation of the most common types of spam

Type of Spam	Percentage
Adult	14.5
Financial	16.2
Health	7.3
Leisure	7.8
Products	21.0
Scams	14.2

Use the table to answer the questions a and b

- (a) Report the modal spam category
- (b) Construct a **Pareto chart** using the table above

(9) A farmer in Idaho is interested in the number of rainy days in a given year so he records the number of rainy R , cloudy C and sunny S days over two weeks in May. His observations are

$$\{R, R, C, C, C, S, S, C, R, R, C, R, S, S\}$$

Use the farmer’s observations to answer the following questions.

- (a) Based on the farmer's observations, what proportion of days were rainy?
- (b) Is the proportion in part (a) a statistic or a parameter? why?
- (c) What population is the farmer studying? is it a finite or infinite population?

(10) Consider the following data from a survey conducted on college students in the state of Florida. As part of their research, surveyors recorded the high school performance (measured in grade point average - GPA) of 35 college students from across the state. For your convenience the data have been sorted from least to greatest:

2.0, 2.1, 2.3, 2.8, 3.0, 3.0, 3.0, 3.0, 3.0, 3.0,
 3.0, 3.0, 3.0, 3.1, 3.2, 3.3, 3.3, 3.4, 3.4, 3.4,
 3.4, 3.5, 3.5, 3.5, 3.5, 3.6, 3.6, 3.7, 3.7, 3.8,
 3.8, 3.8, 3.8, 4.0, 4.0


The following frequency table gives the distribution of the variable high school GPA (with values rounded to nearest 0.5). Fill in the table and answer the following questions:

##	Student	High School GPA	GPA Rounded to Nearest 0.5
## 1	1	2.0	2.0
## 2	2	2.1	2.0
## 3	3	2.3	2.5
## 4	4	2.8	3.0
## 5	5	3.0	3.0
## 6	6	3.0	3.0
## 7	7	3.0	3.0
## 8	8	3.0	3.0
## 9	9	3.0	3.0
## 10	10	3.0	3.0
## 11	11	3.0	3.0
## 12	12	3.0	3.0
## 13	13	3.0	3.0
## 14	14	3.1	3.0
## 15	15	3.2	3.0
## 16	16	3.3	3.5
## 17	17	3.3	3.5
## 18	18	3.4	3.5
## 19	19	3.4	3.5
## 20	20	3.4	3.5
## 21	21	3.4	3.5
## 22	22	3.5	3.5
## 23	23	3.5	3.5
## 24	24	3.5	3.5
## 25	25	3.5	3.5
## 26	26	3.6	3.5
## 27	27	3.6	3.5
## 28	28	3.7	3.5
## 29	29	3.7	3.5
## 30	30	3.8	4.0
## 31	31	3.8	4.0
## 32	32	3.8	4.0
## 33	33	3.8	4.0
## 34	34	4.0	4.0
## 35	35	4.0	4.0


GPA	Frequency	Relative frequency	Cumulative RF
2.0	2	0.06	
2.5	1		
3.0			0.43
3.5	14	0.4	
4.0	6	0.17	

- (a) What type of variable is GPA?
 - (b) What proportion of college students had a high school GPA > 3.0 ? (use the rounded frequency table)
 - (c) What proportion of college students had a high school GPA < 3.0 ? (use the rounded frequency table)
 - (d) What is the **mean** GPA in this sample?
 - (e) What is the **median** GPA in this sample?
 - (f) What is the **mode** of GPA in this sample?
 - (g) Construct a dot plot for the variable GPA (hint use the frequency table)
 - (h) Construct a histogram for the variable GPA (hint use the frequency table)
 - (i) Construct a stem and leaf plot for the variable GPA
- (11) Which statistic is more resistant to outliers, the mean or median? Why?
- (12) Describe the shape of the following distributions and for each distribution identify if the mean will be larger, smaller or the same as the median.


(a)



(b)



(c)



(13) Consider the following four sets of observations of a quantitative variable x . For your convenience the observations have been sorted in increasing order. Match datasets 1 – 4 with the correct histogram (labeled A – D)


Dataset 1 = {0.1, 1.1, 2.6, 2.7, 3.4, 3.4, 4.1, 4.4, 8.8, 9.6}

Dataset 2 = {0.1, 0.3, 1.2, 2.4, 4.4, 4.5, 8.0, 8.9, 9.3, 9.3}


Dataset 3 = {1.1, 3.8, 5.3, 6.0, 6.2, 6.9, 7.9, 7.9, 8.1, 8.7}

Dataset 4 = {3.4, 4.5, 5.4, 5.6, 7.0, 8.5, 8.9, 9.2, 9.7, 9.7}


Distribution A




Distribution B



Distribution C



Distribution D



(14) Consider the following set of 10 observations of a variable X sorted from least to greatest:

3.3, 3.8, 4.0, 4.8, 4.8, 5.1, 5.2, 5.6, 5.7, 6.9

Use the data to answer parts a-b

- (a) Given the data, compute the 40th percentile of X
- (b) Compute Interquartile Range (IQR) range of X

(15) Consider the following $n = 20$ observations of several popular cereal brands and answer questions a - g about the variable “Sugar (g)”:

Brand	Sodium (mg)	Sugar (g)	Type
Frosted Mini Wheats	0	11	A
Raisin Bran	340	18	A
All Bran	70	5	A
Apple Jacks	140	14	C
Captain Crunch	200	12	C
Cheerios	180	1	C
Cinnamon Toast Crunch	210	10	C
Crackling Oat Bran	150	16	A
Fiber One	100	0	A
Frosted Flakes	130	12	C
Froot Loops	140	14	C
Honey Bunches of Oats	180	7	A
Honey Nut Cheerios	190	9	C
Life	160	6	C
Rice Krispies	290	3	C
Honey Smacks	50	15	A
Special K	220	4	A
Wheaties	180	4	A
Corn Flakes	200	3	A
Honeycomb	210	11	C

- (a) Construct a histogram of the data using the following bins: $\leq 2(g)$, $2-4(g)$, $4-6(g)$, $6-8(g)$, $8-10(g)$, $10-12(g)$, $12-14(g)$, $14-16(g)$, $> 16(g)$ (hint start by creating a frequency table)
- (b) Compute the mean, median, and mode of the variable “Sugar (g)”
- (c) Compute the variance and standard deviation of the variable “Sugar (g)”
- (d) Compute the quartiles (Q1, Q2, Q3) and interquartile range of the variable “Sugar (g)”
- (e) Create a box and whisker plot for the variable “Sugar (g)” and mark the Q1, Q2, and Q3 quartiles, the median and any potential outliers
- (f) Plot the Cumulative Distribution of the variable “Sugar (g)”
- (g) What proportion of cereals have a sugar content less than or equal to 10 grams?