Weeks 4-5 Notes

STAT 251 SECTION 01

Last week we wrapped up **descriptive statistics**. We can now describe the shape, center, and spread of a distribution, as well as the position of any individual value within it. All of that material is fair game for Exam 1.

This week we take a step back and ask a question that really should come first: how were the data collected in the first place? How a set of data is collected determines what we are allowed to conclude from it. In particular, there are two questions we want to be able to answer about any study:

- **Can we generalize the results from the sample to the whole population?** This depends on the **sampling design**, which is the topic of the first half of this lecture.
- **Can we conclude that one variable causes a change in another?** This depends on the **experimental design**, which we will cover in the second half.

Sampling designs

Starting with a question

Suppose we wanted to know how many hours a week a typical University of Idaho student spends exercising. We obviously can't ask every student, so how would we go about getting the data? Who would we ask, and how would we choose them?

- Take a minute and think about how you would do it. We will come back to your answer throughout this section.

Some terminology


Before we go any further, let's start with some terminology:

- The **population** is the entire group of individuals we want to learn about. In our example, the population is every UI student.
- A **parameter** is a number that describes the population, such as the population mean μ or the population proportion p. Here, the parameter is the mean weekly exercise hours of *all* UI students.

- The **sample** is the part of the population that we actually observe, i.e., the students we end up asking.
- A **statistic** is a number computed from the sample, such as $\bar{x}$ or $\hat{p}$. Here, that would be the mean hours for the students we asked.
- A **census** collects data on every member of the population. A census is usually slow and expensive, and often impossible.
- The **sampling frame** is the list of units we actually draw the sample from, for example the registrar's list of enrolled students. The sampling frame is rarely a perfect copy of the population.
- The **sampling design** is the method we use to choose the sample from the sampling frame.
- A **representative sample** is a sample that has the same characteristics as the population it came from. Getting a representative sample is the goal of every sampling design.
- **Sampling error** is the difference between a statistic and the parameter it estimates that comes purely from which units happened to be chosen. If we took a different sample, we would get a slightly different answer.

Example: census, frame and sampling error

Consider a small school with 100 students, and suppose we ask them "how many hours did you exercise last week?" If we ask all 100 students, that is a **census**, and their mean of 4.6 hours is the **parameter**. Now suppose the registrar's list, which is our **sampling frame**, only has 90 students on it. The 10 students who enrolled late (shown in grey) have no chance of being chosen. A simple random sample of 10 students from the list (shown in red) has a mean of 4.4 hours, which is our **statistic**. The **sampling error** is then 4.4 - 4.6 = -0.2 hours. If we took another random sample, we would get a different sampling error.



Practice: find the sampling frame

Practice: For each of the following, what is the sampling frame? Does it match the population well?

1. Student government wants to know how students feel about raising the parking fee. They email a survey to 500 students picked at random from the university email directory. *Frame: the email directory. It lists nearly every enrolled student, so it matches the population pretty well.*
2. A Moscow coffee shop wants its customers' opinions on a new menu. It surveys 100 people picked at random from its loyalty-app members. *Frame: the loyalty-app member list. Customers who never signed up for the app can't be chosen.*
3. A city council wants residents' views on a new bike lane. It calls numbers picked at random from a printed phone directory. *Frame: the phone directory. It misses everyone without a listed landline, and those people tend to be younger.*

Thinking critically about studies

Whenever we read about a study, there are some questions we should ask ourselves before we trust the results. Section 1.2 of the textbook gives a nice list of things to look out for (the *Critical Evaluation* list):


- **Problems with samples** - is the sample representative of the population? A biased sample will give results that are inaccurate and not valid.
- **Self-selected samples** - responses only from people who chose to respond, like call-in or online polls, are often unreliable.
- **Sample size** - samples that are too small may be unreliable. Larger samples are better when possible, but sometimes a small sample can't be avoided (e.g., crash tests or rare conditions).
- **Undue influence** - were the data collected, or the questions asked, in a way that pushes people toward a certain answer?
- **Nonresponse** - if a lot of people refused to participate, the responses we did get may not represent the population. People with strong opinions tend to be the most likely to respond.
- **Causality** - just because two variables are related does not mean one causes the other.
- **Self-funded or self-interest studies** - a study paid for by someone who wants a particular answer should be read carefully. That said, we should judge a study on its merits and not just on who paid for it.
- **Misleading use of data** - watch out for poorly displayed graphs, incomplete data, or numbers reported without any context.

- **Confounding** - when the effects of several factors on the response can't be separated, it is hard (or impossible) to draw conclusions about any one of them.

We will spend most of this lecture on these issues. We start with bias and self-selected samples, then nonresponse and question wording, and finally causality and confounding in the second half.

Examples: misleading graphs

OxyContin. Purdue promoted OxyContin, a 12-hour opioid painkiller, using a chart of blood levels over time with the caption "Fewer 'peaks and valleys' than with immediate-release oxycodone." The catch is that the chart was drawn on a logarithmic y-axis (CBS News, 2011 (https://www.cbsnews.com/news/how-purdue-used-misleading-charts-to-hide-oxycontins-addictive-power/)). On a log axis, equal distances represent equal *ratios* rather than equal differences, so a steep drop can look pretty gentle. The figure below is only an illustration and is not Purdue's actual data. It shows a single curve matched to the values in the 2008 FDA label for a 10 mg dose (a peak of 10.6 ng/mL at about 2.7 hours and a half-life of 4.5 hours), drawn on both types of axis.



On the linear axis, the blood level drops from 10.6 to 3.2 ng/mL over the 12 hours, which is a drop of about 70%. In 2007, The Purdue Frederick Company pleaded guilty to misbranding OxyContin "with the intent to defraud and mislead." The government's statement of facts says sales staff used "graphs that exaggerated the differences between blood plasma levels" of OxyContin and other pain medications, and the company paid about $600 million (U.S. Attorney, Western District of Virginia, 2007 (https://web.archive.org/web/2017/https://media.defense.gov/2007/May/10/2001711223/-1/-1/1/purduefrederick1.pdf)). To be clear, there is nothing wrong with a log axis by itself, and scientists use them all the time. The problem is using one to back up a claim that the reader has no way to check.

ExxonMobil. We should read a company's own report just as carefully as anyone else's, especially when the company has a financial stake in the answer. ExxonMobil's *2024 Advancing Climate Solutions* report shows its plan for cutting its *emissions intensity* (emissions per unit produced) as a bar chart with no numbers on the y-axis. Its abatement curve has no axis numbers either. So a reader can't tell how big the planned cut actually is, or how much of it depends on "CCS, hydrogen, and/or future advancements" (Union of Concerned Scientists, 2024 (https://blog.ucs.org/carly-phillips/greenwashing-in-graphs-an-exxonmobil-story/)).

Keep in mind that intensity can go down while total emissions go up, if the company produces more. In this case, the report's own table shows both going down from 2016 to 2023. Intensity fell from 26.4 to 23.3 tonnes per 100 tonnes of throughput or production, and emissions from its operations (Scope 1 and 2) fell from 117 to 98 million tonnes. What neither number includes is the emissions from burning the oil and gas the company produces. That was about 540 million tonnes in 2023, roughly 85% of its footprint, and the company sets no target for it (ExxonMobil, 2024 (https://corporate.exxonmobil.com/-/media/global/files/advancing-climate-solutions/2024/2024-advancing-climate-solutions-report.pdf)).

Convenience and voluntary response samples

There are two ways of getting a sample that are tempting because they are quick, cheap, and easy:

- A **convenience sample** is made up of whoever is easiest to reach. For example, we could stand outside the Student Recreation Center and ask the first 50 people who walk by.
- A **voluntary response sample** is one where we put out a call and let people decide whether or not to respond. For example, we could post a poll on a UI student Instagram account and count the replies. The textbook calls these *self-selected samples*.
- What could go wrong with each of these?

Consider the Rec Center survey first. Its average will be **too high**, since the people at the Rec Center are the ones who exercise, and students who never go there have no chance of being chosen. The Instagram poll only hears from followers who happen to see the post *and* decide to reply, which will often be the proudest gym-goers or people who feel strongly one way or the other. Also notice that asking 2,000 people instead of 50 does not fix either problem. More of the same kind of people just gives us the same wrong answer with more confidence.

What these two designs have in common is that **a person decides who ends up in the sample**. Either the researcher picks whoever is handy, or the respondents pick themselves. A design like this is called **biased**, meaning it systematically favors certain outcomes over others.

Practice: convenience or voluntary response?

Practice: For each scenario below, identify the type of sample and what could go wrong.

1. A nutrition club wants to know how many servings of vegetables UI students eat a day. Members ask the first 60 students who walk out of the campus dining hall at lunch. *Convenience sample. Only students who eat lunch at the dining hall can be chosen, so off-campus students and students who cook for themselves are left out.*
2. The campus paper wants to know whether students favor moving 8 a.m. classes later. It posts a "Vote now!" link on social media; 1,200 students vote and 85% say yes. *Voluntary response sample. Students who dread 8 a.m. classes are the most eager to vote, so 85% probably overstates support. Having 1,200 votes doesn't fix that.*
3. A professor wants to know how many hours a week the 300 students in her lecture study. She asks the 25 students who come to her office hours that week. *Convenience sample, since it's just whoever showed up. Students who come to office hours probably study more than the class as a whole, so the estimate is likely too high.*

Example: the Twitter poll

On December 18, 2022, Elon Musk posted a poll on Twitter asking *"Should I step down as head of Twitter? I will abide by the results of this poll."* In 12 hours the poll got more than 17.5 million votes, and 57.5% of them said yes (NBC News (https://www.nbcnews.com/tech/social-media/twitter-users-vote-elon-musk-quit-ceo-poll-rcna62332); Associated Press (https://www.pbs.org/newshour/nation/twitter-poll-closes-majority-of-users-vote-for-musk-to-step-down-as-ceo)). But the only people who could vote were those who saw the post and chose to vote, which was mostly Musk's own followers, and there was nothing stopping fake accounts from voting. This is a voluntary response sample. The 17.5 million votes only tell us about the people who chose to vote, and the huge sample size doesn't change that.


Example: online opt-in polls

Many online polls use what are called *opt-in panels*. Anyone can sign up to take surveys, and they get paid for each survey they complete. In opt-in samples, 12% of adults under 30 (and 24% of Hispanic respondents) said they were licensed to operate a nuclear submarine. In December 2023, an opt-in poll found that 20% of adults under 30 agreed that "the Holocaust is a myth." When Pew Research Center asked the same question to a probability-based (i.e., random) sample, the figure

was 3% (Mercer, Kennedy and Keeter, 2024 (https://www.pewresearch.org/short-reads/2024/03/05/online-opt-in-polls-can-produce-misleading-results-especially-for-young-people-and-hispanic-adults/)). Most of this difference came from bogus and careless respondents who were answering for the money rather than giving their real opinions. So when you see a shocking number about some subgroup, keep in mind that it may say more about who signed up for the poll than about what people actually think.

Bias vs. standard error

A helpful way to think about bias is to picture a target. The parameter is the bullseye, and each estimate is one shot at the target. Every time we take a sample and compute the statistic, we take a shot and see where it lands. If we take a new sample, we take a new shot. There are two different things that can go wrong:




- **Bias** is a problem with *accuracy*. The shots are centered away from the bullseye and miss in the same direction every time. Bias comes from the design (i.e., who gets into the sample), and it does not average out.
- **Standard error** is a problem with *precision*. It measures how spread out the shots are from one sample to the next. Recall from the Week 3 notes that $SE(\bar{x}) = s/\sqrt{n}$, which gets smaller as the sample size gets bigger.

Convenience and voluntary response samples are biased. No matter how carefully we carry them out, they are aimed at the wrong spot.

A bigger sample does not fix bias

To see this, consider the simulation below. Suppose that 40% of UI students exercise at least three times a week. One survey asks $n = 2000$ people at the Student Recreation Center, where 75% of the people exercise that often. The other survey is a simple random sample of only $n = 200$ students. We repeat each survey 5000 times.



Notice that the Rec Center survey is much more *precise*, since its estimates barely change from one repeat to the next. But every single one of its 5000 estimates misses the true value by about 35 percentage points. Simply put, collecting more data gets us more precision, but only a better design gets us accuracy.


Random sampling

Convenience and voluntary response samples go wrong because a person gets to decide who is in the sample, so the solution is to take the person out of the decision. **Random sampling** uses chance to choose the sample, so neither the researcher nor the respondents decide who gets in. Since nobody is doing the choosing, the selection process can't systematically favor anyone. Of course, chance will still cause our estimate to change from sample to sample. However, it changes according to rules we understand, which means we can measure how much it changes. This is exactly what the standard error does.

Simple random sampling

A **simple random sample (SRS)** of size n is chosen so that every possible group of n units has the same chance of being the sample. As a result, every unit in a population of size N has the same chance, n/N, of being included in the sample.

All of the grid pictures in these notes use the same population: $N = 100$ students on a 10×10 grid, where each row is a class year (40 first-years, 30 sophomores, 20 juniors, and 10 seniors). Every design draws a sample of $n = 20$ students, so each student in the SRS below had a $20/100$, or 1 in 5, chance of being selected.



Take a look at the class years of the students in red. This SRS happened to pick 4 seniors but only 2 juniors, even though there are twice as many juniors as seniors. This isn't a problem with the method. An SRS is fair *on average* across all of the samples it could have produced, but that doesn't mean every individual sample will be balanced.

How to draw an SRS

In practice, drawing an SRS comes down to three steps:


1. Get a sampling frame and number its units from 1 to N.
2. Use a random number generator to choose n different numbers.
3. The units with those numbers make up the sample.

In R, the `sample()` function takes care of step 2. For example, using the `heights.csv` data from the course Data page ($N = 379$):

```
heights <- read.csv('Data/heights.csv')
set.seed(251)                                  # makes the draw repeatable
srs_rows <- sample(1:nrow(heights), 30)        # 30 of the 379 row numbers
head(heights[srs_rows, ])
```

```
##     HEIGHT GENDER
## 348     73   Male
## 291     69   Male
## 48      63 Female
## 339     72   Male
## 238     69 Female
## 59      63 Female
```

By default, `sample()` draws **without replacement**, meaning once a unit has been picked it can't be picked again. This is the usual choice for surveys. When we sample **with replacement**, every unit is put back after it is drawn, so the same unit could show up in the sample more than once (`sample(..., replace = TRUE)`). When N is much larger than n, the two methods are practically the same, because the chance of drawing the same unit twice is very small.



In the figure above, when we sample without replacement each ball that is drawn is removed from the bag, so the sample 3, 1, 5 can't have any repeats. When we sample with replacement, every ball goes back in the bag before the next draw, so 3 can be drawn twice.

Examples: two recent sampling designs


A national poll. In *State of the Facts 2026*, the AP-NORC Center for Public Affairs Research and USAFacts asked U.S. adults how much they trust government information and AI. The poll interviewed 1,036 adults from July 29 to August 10, 2026: 1,017 online and 19 by phone, in English or Spanish. The margin of sampling error is ± 4.1 percentage points (AP-NORC and USAFacts, 2026 (htt

ps://apnorc.org/projects/state-of-the-facts-2026/)). The respondents came from AmeriSpeak, NORC's *probability-based* panel. NORC randomly selects U.S. addresses and then recruits those households by mail, by phone and in person. Nobody can sign themselves up, so chance decides who is invited (NORC, About AmeriSpeak (https://www.norc.org/services-solutions/amerispeak/about.html)). That is the opposite of the opt-in panels above. Random selection doesn't solve everything, though. Only 10.3% of the invited panel members completed this survey (AAPOR response rate 3), so nonresponse is still a concern. We come back to it under nonsampling error below. The methodology is on page 42 of the full report (https://apnorc.org/wp-content/uploads/2026/09/State_of_the_facts_2026.pdf).

A longitudinal study. The UK Biobank is a cohort of about 500,000 UK adults aged 40–69 who agreed to join between 2006 and 2010. Researchers have followed them through their health records ever since. From 2013 to 2015, 103,615 of them wore a wrist sensor with a light meter for 7 days. A 2026 study then removed people with unreliable sensor data, 2,770 night-shift workers, anyone who already had heart disease, and anyone without a later heart MRI. That left 11,071 people, who had an MRI of their heart a median of 3.1 years after wearing the sensor. Hospital and death records were followed for 8 to 10 years (Dai et al., 2026 (https://doi.org/10.1093/eurheartj/ehag563); ESC, 2026 (https://www.escardio.org/news/press/press-releases/too-much-light-at-night-changes-the-shape-and-working-of-the-heart/)). Because the same people are measured at more than one point in time, this is a **longitudinal** study: night light was measured first and the heart later. It is not a random sample of any population, however. The authors point out that the UK Biobank cohort is mostly White and healthier than the general population, so the results may not generalize to everyone. We return to this study under association vs. causation.

Systematic sampling

A **systematic sample** selects every k^{th} unit from an ordered list, starting at a *randomly chosen* point among the first k units. With $N = 100$ and $n = 20$, we have $k = N/n = 5$. In the figure below, the 100 students are numbered in list order. The random starting point was 2, so the sample is made up of positions 2, 7, 12, 17, …, 97.



Systematic sampling is easy to carry out in the field, where there often isn't a list at all until the units show up. As long as the order of the list has nothing to do with what we are measuring, a systematic sample behaves like an SRS. The danger is when the list has a repeating pattern that lines up with k. For example, on our seating grid with rows of 10, counting every 5th seat would select the *same two columns* every time.

For a project dataset with $N = 379$ rows and $n = 30$, we have $379/30 = 12.6$, so we would take $k = 12$ and pick a random starting point between 1 and 12:


```
set.seed(251)
k <- 12
start <- sample(1:k, 1)
sys_rows <- seq(start, by = k, length.out = 30)
sys_rows
```

```
##  [1]   7  19  31  43  55  67  79  91 103 115 127 139 151 163 175 187
199 211 223
## [20] 235 247 259 271 283 295 307 319 331 343 355
```

Stratified random sampling

Strata are groups of units that are similar in some way, and they are formed *before* we take the sample. Examples include class year, region, age group, or sex. A **stratified random sample** takes a separate SRS within each stratum and then combines them.

Under **proportional allocation**, each stratum gets its share of the total sample size n. Our population is 40% first-years, 30% sophomores, 20% juniors, and 10% seniors, so a sample of 20 students would take 8, 6, 4, and 2 students from each class year, respectively.




Stratifying guarantees that every group is represented in the sample, so we can't end up with only 2 juniors by bad luck. It also gives more precise estimates than an SRS of the same size, as long as units are similar within strata and different between strata. This is because the variation from stratum to stratum is removed from the sampling error. If the strata are no more alike inside than the population as a whole, stratifying doesn't buy us anything.

Example: proportional allocation with the heights data

The `heights.csv` data has $N = 379$ people: 262 female and 117 male. For a stratified sample of $n = 30$ we would take

$$\text{female: } 30 \times \frac{262}{379} = 20.74 \approx 21, \qquad \text{male: } 30 \times \frac{117}{379} = 9.26 \approx 9$$



We would then draw an SRS of 21 of the women and a separate SRS of 9 of the men:


```
set.seed(251)
women <- which(heights$GENDER == 'Female')
men   <- which(heights$GENDER == 'Male')
strat_rows <- c(sample(women, 21), sample(men, 9))
table(heights$GENDER[strat_rows])
```

```
##
## Female   Male
##     21      9
```

Large national surveys use stratification. For example, the Current Population Survey, which is used to produce the U.S. unemployment rate, groups areas into strata that are "as homogeneous as possible with respect to labor force and other social and economic characteristics" and then samples within each stratum (Census Bureau and BLS, Technical Paper 77 (https://www2.census.gov/programs-surveys/cps/methodology/CPS-Tech-Paper-77.pdf)).

Cluster sampling


A **cluster** is a group of units that occurs naturally, such as a dorm, a classroom, a city block, or a village. A **cluster sample** takes an SRS of clusters and then measures *every* unit in the clusters that were chosen. In the figure below, each column is a cluster, and two columns were chosen at random.



The main appeal of cluster sampling is practical. We only need a list of the clusters rather than a list of every individual, and visiting a few clusters is much cheaper than traveling to units spread out across a whole region. Cluster sampling works best when each cluster is like a *mini-population*, meaning it is mixed inside, the same way every column above contains all four class years. When units in the same cluster tend to be similar to each other (e.g., students in the same dorm or households on the same street), a cluster sample carries less information than an SRS of the same size, and its estimates are less precise.

Two-stage cluster sampling


A **two-stage cluster sample** first takes an SRS of clusters (stage 1) and then takes an SRS of units within each chosen cluster (stage 2). In the figure below, 4 of the 10 columns were chosen, and then 5 students were chosen from each of those columns.



Election exit polls are a good example of combining several of these designs. For the 2024 election, the National Election Pool drew a stratified probability sample of 279 polling places. Within each polling place, an interviewer approached every n^{th} voter leaving, which is a systematic sample within each cluster (National Election Pool methods statement, 2024 (https://s.abcnews.com/assets/dtci/elections/NEPExitPollMethodologyStatement.pdf)). Similarly, NHANES, the U.S. health and nutrition survey, uses "a complex, multistage, probability sampling design," which samples counties, then segments within counties, then households, and finally individuals (CDC, NHANES sample design (https://wwwn.cdc.gov/nchs/nhanes/tutorials/sampledesign.aspx)).

Strata vs. clusters

Strata and clusters are easy to mix up, since both involve dividing the population into groups. The difference is in what the groups look like inside, and as a result, how many of the groups we sample:



	Strata	Clusters
Units inside a group are…	alike	mixed (a mini-population)
Groups differ from each other?	yes	ideally not
Which groups are sampled?	**every** stratum	**some** clusters
Within a chosen group…	an SRS	every unit (or an SRS, in two stages)
Main benefit	precision, and every group represented	cost and convenience

Sampling and nonsampling error

No sample will match its population exactly. It helps to separate the reasons why into two types of error:

- **Sampling error** is the natural difference between a sample and the population that comes from which units happened to be chosen. Every sample has some sampling error, and a larger sample makes it smaller.
- **Nonsampling error** is error that does not come from chance. This includes biased sampling methods, poorly worded questions, data entry mistakes, or a faulty measuring instrument. A larger sample does not make nonsampling error any smaller.
- **Sampling bias** occurs when some members of the population are less likely to be chosen than others, for example in convenience and voluntary response samples, or when the sampling

frame leaves people out.

Common sources of nonsampling error

Even if we use a perfectly random design, we can still end up with a biased sample. Three common culprits are:

- **Undercoverage** - the sampling frame leaves out part of the population. For example, students who aren't on the list you sampled from, or people who don't have a phone or a fixed address.
- **Nonresponse** - people who are selected for the sample can't be reached or refuse to answer.
- **Response bias** - the answers themselves are wrong. This can be caused by leading or confusing questions, sensitive topics, or respondents wanting to please the interviewer.

Just like before, a larger sample doesn't fix any of these.

Example: nonresponse

Pew Research Center's telephone surveys reached 36% of the people they called in 1997, but only 6% in 2018 (Kennedy and Hartig, 2019 (https://www.pewresearch.org/short-reads/2019/02/27/response-rates-in-telephone-surveys-have-resumed-their-decline/)). A low response rate by itself doesn't make a survey biased. It only causes bias when the people who respond are different from the people who don't. Unfortunately, this is usually impossible to check, because we have no data on the people who didn't respond.

Nonresponse also came up after the 2020 presidential election. National polls in the final two weeks overstated Joe Biden's margin by 3.9 points on average, which was the largest polling error for the national popular vote in 40 years. The American Association for Public Opinion Research's task force ruled out several explanations, like late-deciding voters, and concluded that "at least some of the polling error in 2020 was caused by unit nonresponse." However, without knowing how the people who didn't respond compare with the people who did, they could not pin down the primary source of the error (AAPOR Task Force on 2020 Pre-Election Polling (https://aapor.org/wp-content/uploads/2022/11/Task-Force-on-2020-Pre-Election-Polling_Executive-Summary.pdf)).

Example: question wording

In September 2013, CNBC asked half of the people in its poll about "Obamacare" and the other half about "the Affordable Care Act." These are the same law! Opposition was 46% when it was called "Obamacare" and 37% when it was called "the Affordable Care Act." Interestingly, support was *also* higher for "Obamacare" (29% vs. 22%). This is because far fewer people chose "don't know" for

"Obamacare" (12%) than for the Affordable Care Act (30%) (NBC News, 2013 (https://www.nbcnews.com/news/amp/wbna53122899); Harvard Political Review (https://harvardpolitics.com/aca-vs-obamacare-american-people/)). So the words we use in a question can change the answers we get.

The wording of a question can also change what the question is actually measuring. In January 2003, 68% of Americans favored "taking military action in Iraq to end Saddam Hussein's rule." But when the question added "even if it meant that U.S. forces might suffer thousands of casualties," support dropped to 43% (Pew Research Center, Writing Survey Questions (https://www.pewresearch.org/our-methods/u-s-surveys/writing-survey-questions/)).

Sampling for your course project

Part 3 of the course project (project.html) asks you to draw a sample of at least 20-30 observations from your dataset. When you write this part up, you should:

- choose a sampling design (simple random, systematic, stratified, or cluster sampling)
- explain exactly how you drew the sample (the method, n, and the tool you used, such as the R code above) so that someone else could repeat it
- justify why you chose that design for *your* data. Is there a natural group worth stratifying on? Are there natural clusters?
- describe any bias your method might introduce

You can start on Steps 1 through 4 of the project once we finish this lecture.

If you want some practice before starting the project, try the textbook's **Sampling Experiment** lab (Section 1.6). It lists restaurants by city and by entree cost, and asks you to draw a simple random sample, a systematic sample, two stratified samples, and a cluster sample from the same list. The examples in Section 1.2 (Examples 1.11 to 1.13) are also good practice for identifying sampling designs.

Experiments and causation

Observational studies and experiments

Most studies are interested in the relationship between two or more variables. Again, let's start with some terminology:

- The **response variable** is the outcome we measure.


- The **explanatory variable** is the variable we think explains or influences the response.
- An **observational study** observes individuals and measures variables of interest without trying to influence anything.
- An **experiment** deliberately assigns treatments to the units and then measures the response.

The key question for telling these apart is: **were the groups created by the study, or did they already exist?** In an experiment the study imposes the treatments, and it settles who gets which **by chance** rather than by anyone's judgement - that is exactly what random assignment means, and it is the whole reason an experiment can support a claim about cause. In an observational study nothing is imposed: the units are already in their groups, through their own choices or through circumstance, and the researcher only records what is there.

Note that none of this has to do with how the units were *sampled*. An experiment on volunteers is still an experiment, and an observational study on a random sample is still an observational study.

Association vs. causation

- A **statistical association** between two variables is a numerical measure of their relatedness.
- A **causal relationship** between two variables means that changing one variable causes a proportional change in the other. This is also called a cause-and-effect relationship.
- A **lurking variable** is a variable unknown to the researchers and not included in the study, which is associated with both the explanatory and the response variable.
- Two variables are **confounded** when their effects on the response cannot be separated, so neither effect can be credited to one variable alone.



For example, towns that sell more ice cream also have more drownings. This association is real (it shows up in the data every year), but the causal story is wrong. Hot weather is a lurking variable that drives both. It is important to note that a lurking variable doesn't create a fake association. The

association is real, it just has the wrong explanation. An observational study by itself can't rule this out, because the groups being compared may differ in ways that nobody measured.

Example: hormone therapy

For many years, large observational studies found that women who took hormone therapy after menopause had much less heart disease. In the Nurses' Health Study, which followed 59,337 women, current users of estrogen plus progestin had a relative risk of major coronary heart disease of 0.39 compared to non-users (Grodstein et al., 1996 (https://europepmc.org/article/MED/8672166)). Millions of women were prescribed hormones, partly to protect their hearts.

Then the Women's Health Initiative *randomly assigned* 16,608 women aged 50 to 79 to take either estrogen plus progestin or a placebo. In 2002, after an average of 5.2 years, the trial was stopped early because the risks outweighed the benefits. Invasive breast cancer had crossed its stopping boundary, and the hormone group actually had *more* coronary heart disease, not less (hazard ratio 1.29) (Writing Group for the WHI Investigators, 2002 (https://doi.org/10.1001/jama.288.3.321)).

So why did the observational studies point in the wrong direction? It turns out that women who chose to take hormone therapy tended to be wealthier, better educated, and healthier to begin with. When the observational studies were pooled and adjusted for socioeconomic status, the apparent benefit went away (Humphrey, Chan and Sox, 2002 (https://europepmc.org/article/MED/12186518)). In the randomized trial, random assignment balanced these lurking variables between the two groups automatically.

It is worth being precise about what that trial did and did not test, because the result was reported far more broadly than it was designed to support. The WHI estrogen-plus-progestin trial was a **primary prevention** trial: it enrolled postmenopausal women aged 50 to 79 with an intact uterus, whose average age was about 63, more than a decade past the typical age of menopause. Every woman in the treatment group received the same fixed daily dose of one oral formulation, 0.625 mg of conjugated equine estrogens plus 2.5 mg of medroxyprogesterone acetate in a single tablet. Its primary outcome was coronary heart disease, with invasive breast cancer as the primary adverse outcome. It was not asking whether hormone therapy relieves menopausal symptoms, it did not test transdermal patches or doses tailored to the patient, and it did not enroll perimenopausal women at all. Its conclusion was about *that* regimen, started *that* late, used for *chronic disease prevention*.

The labeling has since caught up with those distinctions. In November 2025 the FDA asked manufacturers to revise the labels of menopausal hormone therapy products, noting that the average WHI participant was 63 and that the formulation tested is "no longer in common use" (FDA, 2025 (https://www.fda.gov/drugs/drug-alerts-and-statements/fda-requests-labeling-changes-related-safety-information-clarify-benefitrisk-considerations)). On February 12, 2026 the FDA approved the

first batch of those changes, removing the statements about cardiovascular disease, breast cancer, and probable dementia from the **boxed warning** of six products (FDA, 2026 (https://www.fda.gov/news-events/press-announcements/fda-approves-labeling-changes-menopausal-hormone-therapy-products)). The boxed warning about endometrial cancer remains for estrogen-alone products, and information about cardiovascular and breast cancer risk remains elsewhere in the labeling; what changed is how prominently and how broadly the risks are stated. The FDA's labeled recommendation is now to begin systemic therapy within 10 years of menopause, generally before age 60, with dose, route, and duration individualized.

Current reviews make the same point: timing, formulation, and route of administration all matter, and transdermal estradiol appears to have a safer cardiovascular profile than the older oral preparations (Blackburn and Kunadian, 2026 (https://doi.org/10.1093/ehjopen/oeag054)). A 2025 analysis of more than 3,800 postmenopausal women, presented at The Menopause Society annual meeting, found less anxiety and depression with transdermal than with oral therapy and no difference in cardiovascular disease, concluding that the route "should be individualized" (The Menopause Society, 2025 (https://menopause.org/press-releases/oral-or-transdermal-hormone-therapy-the-mental-health-risks-are-not-the-same)). That one is a conference presentation rather than a peer-reviewed paper, so treat it as preliminary.

For our purposes, the interesting part is that the lurking-variable problem from 1996 has not gone away. Observational studies again report lower cardiovascular risk among women who start therapy early: in a 2026 cohort of 83,670 matched pairs, women who began within a year of menopause had a cardiovascular disease hazard ratio of 0.70, about 30% lower than matched non-users (Lu et al., 2026 (https://doi.org/10.1093/hropen/hoag072)). But the 2025 Cochrane review of the *randomized* trials still finds little to no difference in coronary events for women starting within 10 years of menopause (relative risk 0.94, 95% CI 0.78 to 1.13), with increased stroke and venous thromboembolism (Bofill Rodriguez et al., 2025 (https://doi.org/10.1002/14651858.CD004143.pub6)).

- Two study designs, two answers. Which one can tell us whether starting therapy early *prevents* heart disease?

The randomized one. Women who choose hormone therapy are still healthier than women who do not, so the observational gap may once again be partly selection rather than effect. This is not a reason to dismiss the newer evidence, and it is not the 2002 story repeating itself: the case for early, individualized therapy in women with symptoms rests on strong evidence for **symptom relief**, on the mismatch between WHI's population and the women actually being treated, and on the timing evidence. It does not yet rest on a demonstrated randomized cardiovascular benefit. Keeping those two claims separate is exactly the skill this section is about.

Example: coffee and the risk of death

A lurking variable can even make an association point the wrong way. The NIH-AARP Diet and Health Study followed 402,260 adults (229,119 men and 173,141 women) aged 50 to 71 from 1995 to 2008, and asked them at the start how much coffee they drank. When the researchers adjusted only for age, men who drank 6 or more cups a day had a 60% higher risk of death than men who drank no coffee (hazard ratio 1.60). However, coffee drinkers were also much more likely to smoke: 34.7% of the men who drank 6 or more cups a day were current smokers, compared with 4.8% of the men who drank none. After adjusting for smoking and other potential confounders, the same comparison showed a 10% *lower* risk of death (hazard ratio 0.90), with fewer deaths from heart disease among coffee drinkers as well (Freedman et al., 2012 (https://doi.org/10.1056/NEJMoa1112010)).

- What is the lurking variable, and why did the association flip?

Smoking. Heavy coffee drinkers smoked far more, and smoking, not coffee, is what raised their risk of death. Once smoking was accounted for, the harmful-looking "coffee effect" disappeared. Keep in mind that this is still an observational study. The authors state that they cannot tell whether the lower risk after adjustment is caused by coffee or is just an association.

Two decades of better-adjusted research have kept pointing the same way. A 2026 analysis of 354,957 UK Biobank participants, followed for a median of 13 years, found that people drinking 5 or more cups a day had a lower risk of cirrhosis (hazard ratio 0.68, 95% CI 0.58 to 0.79), liver cancer (0.53, 0.34 to 0.83), and death from liver disease (0.58, 0.45 to 0.74) than non-drinkers, with less liver fat on MRI in the subgroup that was scanned (Kim et al., 2026 (https://doi.org/10.1016/j.cgh.2026.04.035)). Notice the cohort: this is the same UK Biobank used in the light-at-night study, so the same caveat applies. The authors say plainly that "residual confounding and reverse causation cannot be excluded" and that their participants were mostly of European ancestry and relatively health-conscious.

For the heart specifically, the American Heart Association reviewed this literature in a 2026 scientific statement and reported that caffeine, "studied primarily in the context of coffee, has been shown either to have no relationship or to be associated with a lower risk of coronary artery disease and heart failure", with moderate consumption also associated with lower stroke risk (Marcus et al., 2026 (https://doi.org/10.1161/CIR.0000000000001454)). The statement is careful about two things worth repeating in class. First, "the majority of human subject-based research is observational". Second, coffee is not one exposure: unfiltered coffee raises LDL cholesterol, and high-dose caffeine in energy

drinks looks harmful rather than helpful. There is even a little randomized evidence now, and it cuts both ways: in a trial among regular coffee drinkers, caffeinated coffee lowered the occurrence of atrial fibrillation but increased premature ventricular contractions.

So the whole story runs the other way from the hormone therapy example, and that contrast is the point:

- With **hormone therapy**, observational studies suggested a benefit, and the randomized trial reversed it. Healthier women were choosing the treatment. Later work on timing, dose, and formulation then reversed the broad conclusion again: the WHI harms belonged to one fixed oral regimen started a decade past menopause, and the FDA now recommends individualized therapy started within 10 years of menopause. The heart benefit for early starters, though, still rests mostly on observational studies.
- With **coffee**, observational studies suggested harm, and that harm came from a lurking variable. Once smoking was accounted for, the association reversed, and it has held up as the evidence improved.

A lurking variable does not push in a predictable direction. It pushed the estimate toward benefit in one case and toward harm in the other, which is why the fix is the study design rather than a correction applied afterwards.

Can we claim cause without an experiment?

Sometimes an experiment is impossible or unethical. For example, nobody can randomly assign people to smoke for thirty years. And yet we are confident that smoking causes lung cancer. How? The evidence was built up over decades of observational studies, and it was convincing because it had many features at once (Hill, 1965 (https://europepmc.org/article/MED/14283879)):

- a **strong** association that was found **consistently** across different populations and study designs
- a **dose-response** pattern, i.e., the more cigarettes someone smokes, the higher their risk
- the right order in **time**, i.e., smoking comes before the cancer
- a **plausible** biological explanation that is supported by laboratory evidence

In 2014, fifty years after the first federal report on smoking, the Surgeon General's report concluded that cigarette smoking and secondhand smoke cause at least 480,000 premature deaths a year in the United States, including about 127,700 deaths from lung cancer (U.S. Department of Health and Human Services, 2014 (https://www.ncbi.nlm.nih.gov/books/NBK294316/)). The point is that without an experiment, a causal claim needs this much evidence behind it. A single observational study is never enough.

Example: light at night and the heart

In September 2026, the *European Heart Journal* published a study of 11,071 adults from the UK Biobank (its sampling design is described under "Examples: two recent sampling designs" above). Each participant wore a light sensor on their wrist for 7 days between 2013 and 2015, and about 3 years later each one had an MRI of their heart. Compared with participants who had no light above 3 lux during the least active part of their night, the participants with the most light above 3 lux had 2.4% more left ventricular mass (adjusted for height), heart walls that were 1.5% thicker, and a 1.9% lower myocardial contraction fraction, which is one measure of how well the heart muscle pumps. These differences grew steadily as the amount of night light went up, i.e., a dose-response pattern (Dai et al., 2026 (https://doi.org/10.1093/eurheartj/ehag563)). The European Society of Cardiology's press release about the study was titled "Too much light at night changes the shape and working of the heart" (ESC, 2026 (https://www.escardio.org/news/press/press-releases/too-much-light-at-night-changes-the-shape-and-working-of-the-heart/)).

- Is this an observational study or an experiment? Does it show that light at night *causes* changes in the heart?

Nobody was assigned to sleep in a bright room or a dark room. The researchers simply measured the light people already had at night, so this is an **observational study**. The authors adjusted for a long list of variables, including age, smoking, physical activity, BMI, air pollution, and neighborhood deprivation, and they state themselves that the observational design does not allow them to conclude cause and effect. Keep in mind that adjusting for a variable only helps if that variable was measured. In the editorial published alongside the study, Münzel, Kuntic and Daiber point out that traffic noise was not measured and could still be confounding the results, that the participants were mostly White and healthier than average, and that "causation cannot be firmly established" (Münzel, Kuntic and Daiber, 2026 (https://doi.org/10.1093/eurheartj/ehag650)). They call for randomized trials.

Now compare this study with the smoking evidence above. The night light study has the right order in **time** (the light was measured before the MRI), a **dose-response** pattern, and a **plausible** explanation, since shorter sleep statistically accounted for part of the association. What it does not have yet is the same result found **consistently** by many studies in different populations. So a headline that says light at night "changes" the heart goes further than the study can. A fair summary is that more light at night is *associated* with these changes in the heart, and that it is worth testing with an experiment.

Experimental design terminology

- **Experimental units** are the individuals the treatments are applied to. When the experimental units are people, we call them **subjects**.
- A **factor** is an explanatory variable that the researcher controls. The specific values the factor is set to are called its **levels**.
- A **treatment** is a specific combination of factor levels.

Example: The VITAL trial was a placebo-controlled trial that randomly assigned 25,871 U.S. adults using **two factors**: vitamin D (2,000 IU a day or a placebo) and marine omega-3 fatty acids (1 g a day or a placebo). Each factor has 2 levels, so there are $2 \times 2 = 4$ treatments, and each participant received one of them. This let the researchers answer two questions with one trial. Over a median of 5.3 years, neither vitamin D nor omega-3 lowered the rate of invasive cancer or of major cardiovascular events (Manson et al., 2019a (https://doi.org/10.1056/NEJMoa1809944); Manson et al., 2019b (https://doi.org/10.1056/NEJMoa1811403)).

Principles of experimental design

A good experiment is built on three principles:

1. **Control** - compare the treatment to a **control group**, so that the treatment is the only systematic difference between the groups.
2. **Randomization** - use chance to assign units to treatments. Randomization balances out the lurking variables we know about and the ones we don't, which is something no amount of careful matching can promise.
3. **Replication** - put enough units in each group so that chance differences between the groups average out.

A **placebo** is a fake treatment that looks exactly like the real one, like a sugar pill or a saline injection. We use placebos because of the **placebo effect**, which is when people improve simply because they believe they are being treated. For example, in one study of a performance-enhancing drug, just believing they had taken the substance improved people's performance almost as much as actually taking it (McClung and Collins, 2007, quoted in Section 1.4 of the textbook). In a **blind** experiment, the subjects don't know which treatment they received. In a **double-blind** experiment, the people who measure the response don't know either.

Example: the Pfizer-BioNTech COVID-19 vaccine trial

In 2020, 43,548 people aged 16 and older were randomly assigned, in equal numbers, to receive either two doses of the Pfizer-BioNTech COVID-19 vaccine or two injections of a saline placebo. Neither the participants nor the staff who assessed them knew who had received which.

Group	COVID-19 cases from 7 days after dose 2
Vaccine	8
Placebo	162

Among the 36,523 participants with no sign of earlier infection. Source: Polack et al. (2020) (https://pmc.ncbi.nlm.nih.gov/articles/PMC7745181/).

The design is what makes this comparison meaningful. The saline injection gives the control group the same experience as the vaccine group, so the difference in cases can't be explained by people's expectations. Randomization means the people who got the vaccine weren't, for example, the more cautious people who would have chosen to get it on their own, which would be a lurking variable confounded with the vaccine. With both of these in place, we can credit the vaccine efficacy of about 95% to the vaccine itself.

Example: placebo surgery

Arthroscopic knee surgery for osteoarthritis was one of the most common operations in the United States, and patients reported that it helped. Moseley and colleagues enrolled 180 patients with knee osteoarthritis at the Houston VA Medical Center and randomly assigned each of them to one of three groups. Two groups received real arthroscopic procedures (debridement or lavage), and the third received *placebo surgery*: skin incisions and a simulated procedure, with no arthroscope inserted. Neither the patients nor the staff who later measured their pain and function knew which group a patient was in, so the trial was **double blind**, and everyone was followed for two years.

This is a **completely randomized design** with three treatments, one of them a placebo. Over the two years of follow-up, neither surgery group ever reported less pain or better function than the placebo group (Moseley et al., 2002 (https://doi.org/10.1056/NEJMoa013259)). In other words, the benefit patients felt came from believing they had been treated, and without a placebo group and blinding, nobody would have known.

Was that ethical?

It is a fair question: the patients in the placebo arm were anaesthetised and cut open with no chance of benefit. Three things are usually pointed to in answer.


- **Informed consent.** Every patient signed a consent form that stated explicitly that they might receive placebo surgery, and that it would do nothing for their arthritis.
- **Board approval.** The trial was reviewed and approved by an institutional review board before it began. We return to what that means under Ethics below.
- **Equipoise.** Nobody actually knew whether the operation worked. It was being performed very widely on the strength of what patients reported, with no randomized evidence behind it. When experts are genuinely uncertain which treatment is better, a trial is justified, and that uncertainty has a name: **equipoise**.

On balance most would say yes — real consent, board approval, genuine uncertainty, and a small risk set against an answer that changed practice for a very large number of people. Sham surgery is still debated, though, and every such trial is judged case by case.

Types of experimental designs


Completely randomized design

The simplest type of experimental design is the **completely randomized design**, where every unit is assigned to a treatment completely at random. The COVID-19 vaccine trial above was a completely randomized design with two groups.



Randomized block design


A **block** is a group of units that are similar in a way that we expect to affect the response. Blocks are formed *before* the treatments are assigned. In a **randomized block design**, the random assignment is done separately within each block, and the treatments are compared within blocks.



For example, suppose men and women respond differently to a treatment. In a completely randomized design, that difference would add to the variation within each treatment group. Blocking on sex removes this extra variation, which makes a real treatment effect easier to detect. You can think of blocking as the experimental design version of stratifying in sampling.

Matched pairs design

A **matched pairs design** is a special kind of block design where each block is a pair of very similar units, such as twins or two plots of land side by side. Within each pair, we flip a coin to decide which unit gets which treatment. Another option is to have each subject serve as their own pair by receiving both treatments in a random order.



Measurements taken before and after on the same people (for example, blood pressure before and after an exercise program) also give us paired data. However, there is no random order and no control group, so anything else that changed between the two measurements is confounded with the program. This makes before-and-after measurements much weaker evidence than a randomized matched pairs experiment.

Practice: design the study

Practice: *(Adapted from the textbook's Try It 1.21.)* Suppose you want to know whether texting slows down how quickly a driver reacts when the car ahead of them brakes.

- What are the explanatory variable, the response variable, and the treatments?
- What design would you use?


- Can the experiment be blinded?

Here is one way to think about it:

- The explanatory variable is whether or not the driver is texting, the response variable is reaction time, and the treatments are driving while texting and driving only.
- We could randomly split drivers into a texting group and a driving-only group, but reaction times vary a lot from person to person. A **matched pairs** design is better here. Each driver does both, in a random order, and is compared with themselves.
- The drivers can't be blinded since they know when they are texting, but the person measuring reaction times can be.

Random sampling vs. random assignment

Chance plays two different roles in a study, and each one lets us draw a different kind of conclusion:



- **Random sampling** allows us to **generalize** from the sample to the population it was drawn from.
- **Random assignment** allows us to conclude **cause and effect**.

Most studies only have one of the two. For example, a clinical trial of volunteers can establish that a vaccine works, but strictly speaking only for people similar to those volunteers. On the other hand, a survey based on a random sample can describe a whole population, but it can't explain why its variables are related.

Ethics

Collecting and using data may not seem like something that raises ethical concerns, but like any human endeavor it does, especially when the data are about people.

Example: the Facebook emotional contagion experiment

For one week in January 2012, Facebook ran a **completely randomized experiment** on 689,003 users. Each user was assigned at random to one of three conditions: posts with positive emotional content were less likely to show up, posts with negative content were less likely to show up, or the feed was left unchanged as a control. The study was published in 2014, and the paper stated that the research "was consistent with Facebook's Data Use Policy, to which all users agree prior to creating an account on Facebook, constituting informed consent for this research" (Kramer, Guillory and Hancock, 2014 (https://pmc.ncbi.nlm.nih.gov/articles/PMC4066473/)).

The journal had concerns. Its Editorial Expression of Concern said it was "a matter of concern that the collection of the data by Facebook may have involved practices that were not fully consistent with the principles of obtaining informed consent and allowing participants to opt out." Because Facebook is a private company, it was not bound by the federal rules for human-subjects research. Cornell University's review board decided that its researcher, who only saw the results, was not directly engaged in human research, so no review was required (PNAS Expression of Concern, via Retraction Watch (https://retractionwatch.com/2014/07/03/rapid-mood-swing-pnas-issues-expression-of-concern-for-controversial-facebook-study/)). The takeaway is that clicking "I agree" on a terms-of-service page is not informed consent, even for a well-designed experiment.

Honest data

Ethics also means being honest with data. An investigation of the social psychologist Diederik Stapel found that he had created and altered datasets, and the falsified data tainted more than 55 papers he authored and 10 Ph.D. dissertations he supervised (Section 1.4 of the textbook). But problems don't have to be that big to matter. Example 1.22 in the textbook describes a researcher surveying households in a community who:

- picks a block where they already know a lot of the neighbors. This is a convenience sample, which is biased, and it is misleading to present it as representative.
- skips four houses where nobody was home and never goes back. This is nonresponse, which may leave out groups like working families.

- fills in four forms with answers copied from the neighbors. This is fabricated data, which is never acceptable.

Why the rules exist

Many of the rules we have today came about because of past abuses. In the U.S. Public Health Service's Untreated Syphilis Study at Tuskegee (1932-1972), about 600 Black men in Macon County, Alabama were enrolled, 399 of them with syphilis. They were told they were being treated for "bad blood". They were never told their actual diagnosis, they were followed for decades without being told the true purpose of the study, and they were not offered penicillin once it became the treatment of choice in the late 1940s (CDC (https://www.cdc.gov/tuskegee/about/index.html)). The study ran for forty years and stopped only after the press exposed it in 1972.

The public response led to the National Research Act of 1974, which created the commission that wrote the 1979 **Belmont Report**. Its three principles are the foundation of the review boards that approve human-subject research today (HHS Office of Research Integrity (https://ori.hhs.gov/content/chapter-3-The-Protection-of-Human-Subjects-belmont-report-1979)):

- **Respect for persons** - people get to decide for themselves whether to take part in research.
- **Beneficence** - do no harm, maximize the possible benefits, and minimize the possible risks.
- **Justice** - the burdens and benefits of research should be shared fairly.

Research ethics today

Some of the ethical requirements that apply to studies today are:

- **Informed consent** - subjects are told the purpose, procedures, and risks of the study ahead of time, and they can refuse or withdraw at any time.
- **Confidentiality** - information about individuals is protected and is never released in a way that could identify them.
- **Institutional Review Boards (IRBs)** - these boards review studies involving human subjects before they begin. At the University of Idaho, the IRB is overseen by the Office of Research Assurances (https://www.uidaho.edu/research/our-people#res-assurance), under FSH 5200, Human Subject Research (https://www.uidaho.edu/policies/fsh/5/5200).

Keep in mind that ethics also applies to data you didn't collect yourself. If you use a public dataset for your project, make sure it doesn't contain any information that could identify individuals.

Conclusion: what you should take away

These two weeks were about where data come from, and they divide neatly in two. The first half was about **how we choose who to look at** — the sampling design. The second half was about **whether we can say one thing caused another** — the study design. The two questions are separate, and the rest of the course rests on both.

The three kinds of study

We have been using these three terms throughout, so here they are in one place.

- An **experimental study** (often just called an **experiment**) assigns the units to treatments - by chance, not by anyone's judgement - and then measures the response.
- An **observational study** measures the explanatory and response variables for each unit without trying to influence anything. The groups already exist, through the units' own choices or through circumstance, and nothing is assigned.
- A **survey** is a particular kind of observational study in which units are selected from a population and asked questions, in order to describe that population. Nothing is assigned; a survey only records what is already there.

The question that separates the first from the other two is always the same: **were the groups created by the study, or did they already exist?** If the study imposed the treatments - deciding by chance who received which - it is an experiment. If the groups were already there and we only recorded them, it is observational. And remember that none of this has to do with how the units were *sampled*: an experiment run on volunteers is still an experiment, and a survey of a perfect random sample is still observational.

The sampling designs

The five random designs, and the two non-random ones to recognize and avoid:

- **Simple random sample (SRS).** Every possible sample of size n is equally likely. The benchmark all the others are measured against.
- **Systematic sample.** Choose a random starting point, then take every kth unit from the sampling frame.
- **Stratified random sample.** Split the population into **strata** that are alike within, then draw an independent SRS from each stratum.
- **Cluster sample.** Split the population into **clusters**, each a mini-population, then randomly select whole clusters and measure everyone in them. In a **two-stage cluster sample**, sample units within the selected clusters instead of taking all of them.

- **Convenience sample** and **voluntary response sample** are *not* random. They let the easiest to reach, or the most motivated, select themselves, and both are biased in ways no amount of extra sample size will fix.

The distinction students most often lose: a **stratum** is a group whose members are alike, and we sample from *every* stratum; a **cluster** is a small copy of the whole population, and we sample *some* clusters and take everyone inside.

The experimental designs

- **Completely randomized design.** Every unit is assigned to a treatment completely at random.
- **Randomized block design.** Units are first grouped into **blocks** that are alike in some way expected to affect the response, and the random assignment is then carried out separately within each block. Blocking is to experiments what stratifying is to sampling.
- **Matched pairs design.** A block design in which each block is a pair of very similar units, or a single subject receiving both treatments in a random order.

Error, bias, and what randomness buys you

- **Sampling error** is the difference between a statistic and the parameter that comes purely from which units happened to be chosen. It shrinks as the sample grows.
- **Nonsampling error** comes from how the study was run — undercoverage, nonresponse, response bias, bad question wording, recording mistakes. It does **not** shrink as the sample grows.
- **Bias** is a systematic tendency to miss the parameter in the same direction. A bigger sample makes a biased estimate more precisely wrong, which is the single most important idea in this half of the course.
- **Random sampling** is what lets you **generalize** from the sample to the population. **Random assignment** is what lets you conclude **cause**. Most studies have one or the other, rarely both, and each buys you exactly one of those two things.

Finally, keep the causation rule in front of you: a **statistical association** between two variables is never on its own evidence that one causes the other, because a **lurking variable** associated with both can produce that association with no causal link anywhere in sight.

Sources

- AAPOR Task Force on 2020 Pre-Election Polling (2021). *Executive Summary*. American Association for Public Opinion Research.
- AP-NORC Center for Public Affairs Research and USAFacts (2026). *State of the Facts 2026*. AP-NORC project page: State of the Facts 2026 (https://apnorc.org/projects/state-of-the-facts-2026/) (report: State of the Facts 2026 full report (PDF) (https://apnorc.org/wp-content/uploads/2026/09/State_of_the_facts_2026.pdf))
- Blackburn, I. and Kunadian, V. (2026). Hormone replacement therapy and cardiovascular risk in postmenopausal women. *European Heart Journal Open*, 6(2), oeag054. Blackburn and Kunadian (2026), European Heart Journal Open article (https://doi.org/10.1093/ehjopen/oeag054)
- Bofill Rodriguez, M., Yong, L. N., Mirkov, S., Bekos, C., Lethaby, A. and Farquhar, C. (2025). Long-term hormone therapy for perimenopausal and postmenopausal women. *Cochrane Database of Systematic Reviews*, 11, CD004143. Bofill Rodriguez et al. (2025), Cochrane review (https://doi.org/10.1002/14651858.CD004143.pub6)
- Dai, J., Dai, W., Heianza, Y. and Qi, L. (2026). Nighttime light exposure and cardiac structure and function. *European Heart Journal*. Dai et al. (2026), European Heart Journal article (https://doi.org/10.1093/eurheartj/ehag563)
- European Society of Cardiology (2026, September 10). Too much light at night changes the shape and working of the heart. Press release. ESC press release: Too much light at night changes the heart (https://www.escardio.org/news/press/press-releases/too-much-light-at-night-changes-the-shape-and-working-of-the-heart/)
- Freedman, N. D., Park, Y., Abnet, C. C., Hollenbeck, A. R. and Sinha, R. (2012). Association of coffee drinking with total and cause-specific mortality. *New England Journal of Medicine*, 366(20), 1891–1904. Freedman et al. (2012), New England Journal of Medicine article (https://doi.org/10.1056/NEJMoa1112010)
- Grodstein, F. et al. (1996). Postmenopausal estrogen and progestin use and the risk of cardiovascular disease. *New England Journal of Medicine*, 335, 453–461.
- Hill, A. B. (1965). The environment and disease: association or causation? *Proceedings of the Royal Society of Medicine*, 58, 295–300.
- Humphrey, L. L., Chan, B. K. and Sox, H. C. (2002). Postmenopausal hormone replacement therapy and the primary prevention of cardiovascular disease. *Annals of Internal Medicine*, 137, 273–284.
- Illowsky, B. and Dean, S. (2023). *Introductory Statistics 2e*, Sections 1.2 and 1.4–1.6. OpenStax. OpenStax: Introductory Statistics 2e (https://openstax.org/details/books/introductory-statistics-2e)
- Kennedy, C. and Hartig, H. (2019). Response rates in telephone surveys have resumed their decline. Pew Research Center.

- Kim, H.-S. et al. (2026). Coffee consumption and improved liver outcomes: clinical, imaging, and proteomic evidence from the UK Biobank. *Clinical Gastroenterology and Hepatology*. Kim et al. (2026), Clinical Gastroenterology and Hepatology article (https://doi.org/10.1016/j.cgh.2026.04.035)
- Kramer, A. D. I., Guillory, J. E. and Hancock, J. T. (2014). Experimental evidence of massive-scale emotional contagion through social networks. *PNAS*, 111(24), 8788–8790. Kramer, Guillory and Hancock (2014), PNAS article (https://doi.org/10.1073/pnas.1320040111)
- Lu, T. F. et al. (2026). Long-term systemic outcomes of postmenopausal hormone replacement therapy by age at menopause: a real-world cohort study. *Human Reproduction Open*, 2026(3), hoag072. Lu et al. (2026), Human Reproduction Open article (https://doi.org/10.1093/hropen/hoag072)
- Manson, J. E. et al. (2019a). Vitamin D supplements and prevention of cancer and cardiovascular disease. *New England Journal of Medicine*, 380(1), 33–44. Manson et al. (2019a), New England Journal of Medicine article (https://doi.org/10.1056/NEJMoa1809944)
- Manson, J. E. et al. (2019b). Marine n−3 fatty acids and prevention of cardiovascular disease and cancer. *New England Journal of Medicine*, 380(1), 23–32. Manson et al. (2019b), New England Journal of Medicine article (https://doi.org/10.1056/NEJMoa1811403)
- Marcus, G. M., Hu, F. B., van Dam, R. M. et al. (2026). Caffeine and cardiovascular disease: a scientific statement from the American Heart Association. *Circulation*, 154, e345–e355. Marcus et al. (2026), AHA scientific statement in Circulation (https://doi.org/10.1161/CIR.0000000000001454)
- The Menopause Society (2025, October 10). *Oral or transdermal hormone therapy? The mental health risks are not the same*. Press release. The Menopause Society press release on oral vs. transdermal hormone therapy (https://menopause.org/press-releases/oral-or-transdermal-hormone-therapy-the-mental-health-risks-are-not-the-same)
- Mercer, A., Kennedy, C. and Keeter, S. (2024). Online opt-in polls can produce misleading results, especially for young people and Hispanic adults. Pew Research Center.
- Moseley, J. B. et al. (2002). A controlled trial of arthroscopic surgery for osteoarthritis of the knee. *New England Journal of Medicine*, 347(2), 81–88. Moseley et al. (2002), New England Journal of Medicine article (https://doi.org/10.1056/NEJMoa013259)
- Münzel, T., Kuntic, M. and Daiber, A. (2026). The dark night as a cardiovascular vital sign: personal light exposure, subclinical cardiac remodelling, and the urban exposome. *European Heart Journal*, ehag650. Münzel, Kuntic and Daiber (2026), European Heart Journal article (https://doi.org/10.1093/eurheartj/ehag650)
- NORC at the University of Chicago (n.d.). About AmeriSpeak. NORC: About AmeriSpeak (https://www.norc.org/services-solutions/amerispeak/about.html)

- Polack, F. P. et al. (2020). Safety and efficacy of the BNT162b2 mRNA Covid-19 vaccine. *New England Journal of Medicine*, 383(27), 2603–2615. Polack et al. (2020), New England Journal of Medicine article (https://doi.org/10.1056/NEJMoa2034577)
- U.S. Department of Health and Human Services (2014). *The Health Consequences of Smoking — 50 Years of Progress: A Report of the Surgeon General*.
- U.S. Food and Drug Administration (2025, November 10). *FDA requests labeling changes related to safety information to clarify the benefit/risk considerations for menopausal hormone therapies*. FDA statement, November 2025: labeling changes for menopausal hormone therapies (https://www.fda.gov/drugs/drug-alerts-and-statements/fda-requests-labeling-changes-related-safety-information-clarify-benefitrisk-considerations)
- U.S. Food and Drug Administration (2026, February 12). *FDA approves labeling changes to menopausal hormone therapy products*. FDA press announcement, February 2026: labeling changes approved (https://www.fda.gov/news-events/press-announcements/fda-approves-labeling-changes-menopausal-hormone-therapy-products)
- Writing Group for the Women's Health Initiative Investigators (2002). Risks and benefits of estrogen plus progestin in healthy postmenopausal women. *JAMA*, 288(3), 321–333. Women's Health Initiative (2002), JAMA article (https://doi.org/10.1001/jama.288.3.321)