

Conformal twistor KIDs

E. Gasperín ^{*,1} and J.L. Williams ^{†,2}

¹*Instituto Superior Técnico*

²*Department of Mathematical Sciences, University of Bath, Claverton Down, Bath BA2 7AY, United Kingdom.*

October 20, 2021

Abstract

Write-up of Edgar's twistor KID calculations from the new notebook.

1 Definitions

We are interested here in the twistor equation:

$$\nabla_{A'}(A\kappa_B) = 0,$$

encoded in the vanishing of the *zero quantity* $H_{A'AB} := 2\nabla_{A'}(A\kappa_B)$. It will prove useful to define the auxiliary quantity

$$\xi_{A'} := \nabla^A{}_{A'}\kappa_A.$$

We also define the zero quantity $S_{A'B'A} := \nabla_{QA'}H_{B'A}{}^Q$. This may be expressed alternatively in terms of the auxiliary field^{•1} $\xi_{A'}$, by an easy computation, as follows:

$$S_{A'B'A} = -\nabla_{AB'}\xi_{A'} + 2\Lambda\epsilon_{A'B'}\kappa_A - 2\Phi_{AQA'B'}\kappa^Q. \quad (1)$$

•1: I think we should change the sign in either the definition of S or ξ , to be more consistent.

The other (symmetrised) contraction yields the *Buchdahl constraint*:

$$0 = \nabla_{(A}{}^{A'}H_{|A'|BC)} = \Psi_{ABCD}\kappa^D,$$

though this won't feature in the calculations here.

2 Wave equations

2.1 For the twistor fields

A short computation shows that

$$\square\kappa_B = -2\Lambda\kappa_B + \frac{2}{3}\nabla^{AA'}H_{A'AB}.$$

Hence, if κ_B solves the twistor equation, then it necessarily satisfies the wave equation

$$\square\kappa_B + 2\Lambda\kappa_B = 0. \quad (2)$$

We will choose to propagate a twistor candidate according to this equation. It is maybe worth noticing that $S_{A'}{}^{A'}{}_B = \nabla^{AA'}H_{A'AB}$. Hence, if κ_A satisfies (2), then necessarily $S_{A'B'A} =$

*E-mail address: edgar.gasperin@tecnico.ulisboa.pt

†E-mail address: jlw31@bath.ac.uk

$S_{(A'B')A}$, since it is assured that $\nabla^{AA'}H_{A'AB} = 0$.

We will also need a wave equation for the auxiliary field $\xi_{A'}$. To get this, take the contracted derivative of (2) and commute derivatives, finally resulting in

$$\square \xi_{A'} + 2\Lambda \xi_{A'} - 8(\nabla_{AA'}\Lambda)\kappa^A = 0. \quad (3)$$

Again, if κ_A is a twistor, then the resulting $\xi_{A'}$ solves (3). Given a twistor candidate and its corresponding $\xi_{A'}$, we choose to propagate the latter according to this wave equation.

2.2 For the zero quantities

In order to derive wave equations for the zero quantities, we assume that twistor candidate, κ_A and its auxiliary spinor $\xi_{A'}$ satisfy the wave equations (2)–(3); at the end of the day, these will be satisfied by construction, since the candidate quantities will be propagated off the initial hypersurface using these wave equations.

To get a wave equation for $H_{A'AB}$, simply take the definition of \mathbf{S} in terms of \mathbf{H} and take a contracted derivative —i.e. consider $\nabla_A{}^{D'}S_{D'A'B}$. Ultimately, we get

$$\square H_{A'AB} = 8\Lambda H_{A'AB} - 2\Psi_{ABCD}H_{A'}{}^{CD} - 2\Phi_{ADA'D'}H^{D'D}{}_B - 2\nabla_{AD'}S^{D'}{}_{A'B}.$$

It is important to note that this is expressible (in a regular way) in terms of the rescaled Weyl spinor $\phi_{ABCD} = \Theta^{-1}\Psi_{ABCD}$:

$$\square H_{A'AB} = 8\Lambda H_{A'AB} - 2\Theta\phi_{ABCD}H_{A'}{}^{CD} - 2\Phi_{ADA'D'}H^{D'D}{}_B - 2\nabla_{AD'}S^{D'}{}_{A'B}.$$

To close the system, we need a wave equation for $S_{A'B'A}$. To get this, we will apply the D'Alembertian to (1), commute derivatives, and substitute the wave equation for $\xi_{A'}$, equation (3). Finally, we arrive at^{•2}

$$\begin{aligned} \square S_{A'B'A} = & 6\Lambda S_{A'B'A} - 4\Phi_{ABC'}(A'S_{B'}{}^{C'B} - 2\Theta\bar{\phi}_{A'B'C'D'}S^{C'D'}{}_A \\ & - \frac{2}{3}\Phi_{BCA'B'}(\nabla_{AC'}H^{C'BC} + 2\nabla^C{}_{C'}H^{C'}{}_A{}^B) \\ & + 4H_{(A'|AB|}\nabla^B{}_{B'}\Lambda - 2(\nabla_{CC'}\Phi_{ABA'B'})H^{C'BC}. \end{aligned} \quad (4)$$

•2: Is it possible to simplify this a bit more?

Note that the terms on the right-hand-side are homogeneous in \mathbf{S} , \mathbf{H} and $\nabla\mathbf{H}$.

3 What goes wrong in the higher-valence case?