


+Vincent.Knight  
@drvinceknight  
[drvinceknight.github.io/Talks](https://drvinceknight.github.io/Talks)

@IzabelaKomenda  
Professor Jeff Griffiths

$$\begin{pmatrix} (2, 2) & (5, 0) \\ (0, 5) & (4, 4) \end{pmatrix}$$



What about the controllers?


# What about the controllers?

S. Deo and I. Gurvich. **Centralized vs. Decentralized Ambulance Diversion: A Network Perspective.** *Management Science*, 57(7):13001319, May 2011.


[INCLUDE ANY PLOT FROM PAPER WITH IZA]


[INCLUDE ANY PLOT FROM PAPER WITH IZA]


**Mathematical modelling of patient flows to predict critical care capacity required following the merger of two District General Hospitals into one., *Submitted to Anaesthesia***













$$(K_{\text{NH}}, K_{\text{RG}}) = (6, 12):$$




$h = \text{NH}$




$h = \text{RG}$

$$(K_{\text{NH}}, K_{\text{RG}}) = (1, 12):$$



$h = \text{NH}$



$h = \text{RG}$

For all  $h \in \{\text{NH}, \text{RG}\}$  minimise:

$$(U_h - t)^2$$

Subject to:

$$0 \leq K_h \leq c_h$$



$$K_h \in \mathbb{Z}$$

$$A = \begin{pmatrix} (U_{\text{NH}}(1, 1) - t)^2 & \dots & (U_{\text{NH}}(1, c_{\text{RG}}) - t)^2 \\ (U_{\text{NH}}(2, 1) - t)^2 & \dots & (U_{\text{NH}}(2, c_{\text{RG}}) - t)^2 \\ \vdots & \ddots & \vdots \\ (U_{\text{NH}}(c_{\text{NH}}, 1) - t)^2 & \dots & (U_{\text{NH}}(c_{\text{NH}}, c_{\text{RG}}) - t)^2 \end{pmatrix}$$


$$B = \begin{pmatrix} (U_{\text{RG}}(1, 1) - t)^2 & \dots & (U_{\text{RG}}(1, c_{\text{RG}}) - t)^2 \\ (U_{\text{RG}}(2, 1) - t)^2 & \dots & (U_{\text{RG}}(2, c_{\text{RG}}) - t)^2 \\ \vdots & \ddots & \vdots \\ (U_{\text{RG}}(c_{\text{RG}}, 1) - t)^2 & \dots & (U_{\text{RG}}(c_{\text{RG}}, c_{\text{RG}}) - t)^2 \end{pmatrix}$$

**Theorem.**

Let  $f_h(k) : [1, c_{\bar{h}}] \rightarrow [1, c_h]$  be the best response of player  $h \in \{\text{NH}, \text{RG}\}$  to the diversion threshold of  $\bar{h} \neq h$  ( $\bar{h} \in \{\text{NH}, \text{RG}\}$ ). If  $f_h(k)$  is a non-decreasing function in  $k$  then the game has at least one Nash Equilibrium in Pure Strategies.










**Lemma.**

- ▶ If  $\lambda_{NH}^{(a)} \leq \lambda_{NH}^{(b)}$  and  $\lambda_{NH}^{(c)} \leq \lambda_{NH}^{(d)}$  then  $f_{NH}(k)$  is a non-decreasing function in  $k$ .
- ▶ If  $\lambda_{RG}^{(a)} \leq \lambda_{RG}^{(c)}$  and  $\lambda_{RG}^{(b)} \leq \lambda_{RG}^{(d)}$  then  $f_{RG}(k)$  is a non-decreasing function in  $k$ .







$(t = 0.8)$



$(t = 0.6)$

$$\text{PoA} = \frac{T^*}{\widetilde{T}}$$





# Conclusions

- ▶ Developed a strategic form game representation of CCU interaction;
- ▶ Proved structural properties of equilibrium behaviour;
- ▶ Identified a potential justified approach to obtaining policies.

# Conclusions

- ▶ Developed a strategic form game representation of CCU interaction;
- ▶ Proved structural properties of equilibrium behaviour;
- ▶ Identified a potential justified approach to obtaining policies.

**Measuring the Price of Anarchy in Critical Care Unit Interactions, *Submitted to OMEGA***



+Vincent.Knight  
@drvinceknight

[drvinceknight.github.io/Talks](https://drvinceknight.github.io/Talks)

@IzabelaKomenda  
Professor Jeff Griffiths