

Regression and Causal Inference

September 8, 2010

Prediction

- We have an input vector $X^T = (X_1, X_2, \dots, X_p)$ with dimensions of $n \times p$ and an output vector Y with dimensions $n \times 1$.

Prediction

- We have an input vector $X^T = (X_1, X_2, \dots, X_p)$ with dimensions of $n \times p$ and an output vector Y with dimensions $n \times 1$.
- The linear regression model has the form:

$$f(X) = \beta_0 + \sum_{j=1}^p X_j \beta_j$$

Prediction


- We have an input vector $X^T = (X_1, X_2, \dots, X_p)$ with dimensions of $n \times p$ and an output vector Y with dimensions $n \times 1$.
- The linear regression model has the form:

$$f(X) = \beta_0 + \sum_{j=1}^p X_j \beta_j$$

- We can pick the coefficients $\beta = (\beta_0, \beta_1, \dots, \beta_p)^T$ in a variety of ways but OLS is by far the most common, which minimizes the **residual sum of squares** (RSS):

$$\begin{aligned} RSS(\beta) &= \sum_{i=1}^N (y_i - f(x_i))^2 \\ &= \sum_{i=1}^N (y_i - \beta_0 - \sum_{j=1}^P x_{ij} \beta_j)^2 \end{aligned}$$

OLS in a Picture



Deriving the Algorithm

- Denote \mathbf{X} the $N \times (p + 1)$ matrix with each row an input vector (with a 1 in the first position) and \mathbf{y} is the output vector.

Deriving the Algorithm

- Denote \mathbf{X} the $N \times (p + 1)$ matrix with each row an input vector (with a 1 in the first position) and \mathbf{y} is the output vector.
- Write the RSS as:

$$RSS(\beta) = (\mathbf{y} - \mathbf{X}\beta)^T(\mathbf{y} - \mathbf{X}\beta)$$

Deriving the Algorithm

- Denote \mathbf{X} the $N \times (p + 1)$ matrix with each row an input vector (with a 1 in the first position) and \mathbf{y} is the output vector.
- Write the RSS as:

$$RSS(\beta) = (\mathbf{y} - \mathbf{X}\beta)^T(\mathbf{y} - \mathbf{X}\beta)$$

- Differentiate with respect to β :

$$\frac{\partial RSS}{\partial \beta} = -2\mathbf{X}^T(\mathbf{y} - \mathbf{X}\beta) \quad (1)$$

Deriving the Algorithm

- Denote \mathbf{X} the $N \times (p + 1)$ matrix with each row an input vector (with a 1 in the first position) and \mathbf{y} is the output vector.

- Write the RSS as:

$$RSS(\beta) = (\mathbf{y} - \mathbf{X}\beta)^T(\mathbf{y} - \mathbf{X}\beta)$$

- Differentiate with respect to β :

$$\frac{\partial RSS}{\partial \beta} = -2\mathbf{X}^T(\mathbf{y} - \mathbf{X}\beta) \quad (1)$$

- Assume that \mathbf{X} is full rank (no perfect collinearity among any of the independent variables) and set first derivative to 0:

$$\mathbf{X}^T(\mathbf{y} - \mathbf{X}\beta) = 0$$

Deriving the Algorithm

- Denote \mathbf{X} the $N \times (p + 1)$ matrix with each row an input vector (with a 1 in the first position) and \mathbf{y} is the output vector.

- Write the RSS as:

$$RSS(\beta) = (\mathbf{y} - \mathbf{X}\beta)^T(\mathbf{y} - \mathbf{X}\beta)$$

- Differentiate with respect to β :

$$\frac{\partial RSS}{\partial \beta} = -2\mathbf{X}^T(\mathbf{y} - \mathbf{X}\beta) \quad (1)$$

- Assume that \mathbf{X} is full rank (no perfect collinearity among any of the independent variables) and set first derivative to 0:

$$\mathbf{X}^T(\mathbf{y} - \mathbf{X}\beta) = 0$$

- Solve for β :

$$\hat{\beta} = (\mathbf{X}^T\mathbf{X})^{-1}\mathbf{X}^T\mathbf{y}$$

Making a Prediction

- The *hat matrix*, or *projection matrix*

$$\mathbf{H} = \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \text{ with } \tilde{\mathbf{H}} = \mathbf{I} - \mathbf{H}$$

Making a Prediction

- The *hat matrix*, or *projection matrix*

$$\mathbf{H} = \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \text{ with } \tilde{\mathbf{H}} = \mathbf{I} - \mathbf{H}$$

- We use the hat matrix to find the fitted values:

$$\hat{\mathbf{Y}} = \mathbf{X}\hat{\beta} = \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{Y} = \mathbf{H}\mathbf{Y}$$

Making a Prediction

- The *hat matrix*, or *projection matrix*

$$\mathbf{H} = \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \text{ with } \tilde{\mathbf{H}} = \mathbf{I} - \mathbf{H}$$

- We use the hat matrix to find the fitted values:

$$\hat{\mathbf{Y}} = \mathbf{X}\hat{\beta} = \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{Y} = \mathbf{H}\mathbf{Y}$$

- We can now write

$$\mathbf{e} = (\mathbf{I} - \mathbf{H})\mathbf{Y}$$

Making a Prediction

- The *hat matrix*, or *projection matrix*

$$\mathbf{H} = \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \text{ with } \tilde{\mathbf{H}} = \mathbf{I} - \mathbf{H}$$

- We use the hat matrix to find the fitted values:

$$\hat{\mathbf{Y}} = \mathbf{X}\hat{\boldsymbol{\beta}} = \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{Y} = \mathbf{H}\mathbf{Y}$$

- We can now write

$$\mathbf{e} = (\mathbf{I} - \mathbf{H})\mathbf{Y}$$

- If $\mathbf{H}\mathbf{Y}$ yields part of \mathbf{Y} that projects into \mathbf{X} , this means that $\tilde{\mathbf{H}}\mathbf{Y}$ is the part of \mathbf{Y} that does not project into \mathbf{X} , which is the *residual* part of \mathbf{Y} . Therefore, $\tilde{\mathbf{H}}\mathbf{Y}$ makes the residuals.

From Algorithm to Model

- 1 *Linear in Parameters*: Y is related to the independent variables and the error term as $Y = X\beta + \epsilon$

From Algorithm to Model

- ① *Linear in Parameters*: Y is related to the independent variables and the error term as $Y = X\beta + \epsilon$
- ② The X 's are fixed and take on ≥ 2 values

From Algorithm to Model

- ① *Linear in Parameters*: Y is related to the independent variables and the error term as $Y = X\beta + \epsilon$
- ② The X 's are fixed and take on ≥ 2 values
- ③ *Full Rank* (in multiple regression): There is no perfect collinearity among any of the independent variables

From Algorithm to Model

- ① *Linear in Parameters*: Y is related to the independent variables and the error term as $Y = X\beta + \epsilon$
- ② The X 's are fixed and take on ≥ 2 values
- ③ *Full Rank* (in multiple regression): There is no perfect collinearity among any of the independent variables
- ④ *Zero Conditional Mean*: $E(\epsilon|X) = 0$

From Algorithm to Model

- ① *Linear in Parameters*: Y is related to the independent variables and the error term as $Y = X\beta + \epsilon$
- ② The X 's are fixed and take on ≥ 2 values
- ③ *Full Rank* (in multiple regression): There is no perfect collinearity among any of the independent variables
- ④ *Zero Conditional Mean*: $E(\epsilon|X) = 0$
- ⑤ *Homoskedasticity*: $\text{Var}(\epsilon|X) = \sigma^2$

From Algorithm to Model

- ① *Linear in Parameters*: Y is related to the independent variables and the error term as $Y = X\beta + \epsilon$
- ② The X 's are fixed and take on ≥ 2 values
- ③ *Full Rank* (in multiple regression): There is no perfect collinearity among any of the independent variables
- ④ *Zero Conditional Mean*: $E(\epsilon|X) = 0$
- ⑤ *Homoskedasticity*: $\text{Var}(\epsilon|X) = \sigma^2$
- ⑥ *Random Sampling*: Y_i is an *iid* random sample, although this can be relaxed to
$$\text{cov}(y_i, y_j) = 0 = \text{cov}(\epsilon_i, \epsilon_j) \quad i \neq j$$

From Algorithm to Model

- ① *Linear in Parameters*: Y is related to the independent variables and the error term as $Y = X\beta + \epsilon$
- ② The X 's are fixed and take on ≥ 2 values
- ③ *Full Rank* (in multiple regression): There is no perfect collinearity among any of the independent variables
- ④ *Zero Conditional Mean*: $E(\epsilon|X) = 0$
- ⑤ *Homoskedasticity*: $\text{Var}(\epsilon|X) = \sigma^2$
- ⑥ *Random Sampling*: Y_i is an *iid* random sample, although this can be relaxed to
$$\text{cov}(y_i, y_j) = 0 = \text{cov}(\epsilon_i, \epsilon_j) \quad i \neq j$$
- ⑦ *Normal Errors* (optional): $Y \sim \mathcal{N}(X\beta, \sigma^2)$

Deriving σ^2

- Recall:

$$\begin{aligned}\hat{\beta} &= (X^T X)^{-1} X^T Y \\ &= (X^T X)^{-1} X^T (X\beta + \epsilon) \\ \Rightarrow \hat{\beta} - \beta &= (X^T X)^{-1} X^T \epsilon\end{aligned}$$

Deriving σ^2

- Recall:

$$\begin{aligned}\hat{\beta} &= (X^T X)^{-1} X^T Y \\ &= (X^T X)^{-1} X^T (X\beta + \epsilon) \\ \Rightarrow \hat{\beta} - \beta &= (X^T X)^{-1} X^T \epsilon\end{aligned}$$

- Plugging this into the covariance equation:

$$\begin{aligned}\text{cov}(\hat{\beta}|X) &= E[(\hat{\beta} - \beta)(\hat{\beta} - \beta)'|X] \\ &= E\left[\left((X^T X)^{-1} X^T \epsilon\right)\left((X^T X)^{-1} X^T \epsilon\right)'|X\right] \\ &= E\left[(X^T X)^{-1} X^T \epsilon \epsilon^T X (X^T X)^{-1}|X\right] \\ &= (X^T X)^{-1} X^T E(\epsilon \epsilon^T | X) X (X^T X)^{-1} \\ &\quad \text{where } E(\epsilon \epsilon^T | X) = \sigma^2 I_{p \times p} \\ &= (X^T X)^{-1} X^T \sigma^2 I_{p \times p} X (X^T X)^{-1} \\ &= \sigma^2 (X^T X)^{-1} X^T X (X^T X)^{-1} \\ &= \sigma^2 (X^T X)^{-1}\end{aligned}$$

Deriving σ^2

We estimate σ^2 dividing the residuals squared by the degrees of freedom because the e_i are generally smaller than the ϵ_i due to the fact that $\hat{\beta}$ was chosen to make the sum of square residuals as small as possible.

$$\hat{\sigma}^2 = \frac{1}{n - p} \sum_{i=1}^n e_i^2$$

Recall:

$$\begin{aligned}\hat{\beta} &= (X^T X)^{-1} X^T Y \\ &= (X^T X)^{-1} X^T (X\beta + \epsilon) \\ &= (X^T X)^{-1} X^T X\beta + (X^T X)^{-1} X^T \epsilon \\ &= \beta + (X^T X)^{-1} X^T \epsilon\end{aligned}$$

We know that $\hat{\beta}$ is unbiased if $E(\hat{\beta}) = \beta$

$$\begin{aligned}E(\hat{\beta}) &= E(\beta + (X^T X)^{-1} X^T \epsilon | X) \\ &= E(\beta | X) + E((X^T X)^{-1} X^T \epsilon | X) \\ &= \beta + (X^T X)^{-1} E(\epsilon | X) \\ &\quad \text{where } E(\epsilon | X) = E(\epsilon) = 0 \\ E(\hat{\beta}) &= \beta\end{aligned}$$

Regression Anatomy

- In the simple bivariate case:

$$\beta_1 = \frac{\text{Cov}(Y_i, X_i)}{\text{Var}(X_i)}$$

Regression Anatomy

- In the simple bivariate case:

$$\beta_1 = \frac{\text{Cov}(Y_i, X_i)}{\text{Var}(X_i)}$$

- In the multivariate case, β_j is:

$$\beta_j = \frac{\text{Cov}(Y_i, \tilde{X}_{ij})}{\text{Var}(\tilde{X}_{ij})}$$

where \tilde{X}_{ij} is the residual from the regression of X_{ij} on all other covariates.

Regression Anatomy

- In the simple bivariate case:

$$\beta_1 = \frac{\text{Cov}(Y_i, X_i)}{\text{Var}(X_i)}$$

- In the multivariate case, β_j is:

$$\beta_j = \frac{\text{Cov}(Y_i, \tilde{X}_{ij})}{\text{Var}(\tilde{X}_{ij})}$$

where \tilde{X}_{ij} is the residual from the regression of X_{ij} on all other covariates.

- The multiple regression coefficient $\hat{\beta}_j$ represents the additional contribution of x_j on y , after x_j has been adjusted for $x_0, x_1, \dots, x_{j-1}, x_{j+1}, \dots, x_p$

Regression Anatomy

- In the simple bivariate case:

$$\beta_1 = \frac{\text{Cov}(Y_i, X_i)}{\text{Var}(X_i)}$$

- In the multivariate case, β_j is:

$$\beta_j = \frac{\text{Cov}(Y_i, \tilde{X}_{ij})}{\text{Var}(\tilde{X}_{ij})}$$

where \tilde{X}_{ij} is the residual from the regression of X_{ij} on all other covariates.

- The multiple regression coefficient $\hat{\beta}_j$ represents the additional contribution of x_j on y , after x_j has been adjusted for $x_0, x_1, \dots, x_{j-1}, x_{j+1}, \dots, x_p$
- What happens when x_j is highly correlated with some of the other x_k 's?

Regression in Causal Analysis

- Imagine we are analyzing a *randomized* experiment with a regression using the following model:

$$Y_i = \alpha + \beta_1 \cdot T_i + \mathbf{X}_i^T \cdot \beta_2 + \epsilon_i$$

where T_i is an indicator variable for treatment status and \mathbf{X}_i is a vector of *pre-treatment characteristics*

- Under this model, what is random?

Regression in Causal Analysis

- Imagine we are analyzing a *randomized* experiment with a regression using the following model:

$$Y_i = \alpha + \beta_1 \cdot T_i + \mathbf{X}_i^T \cdot \beta_2 + \epsilon_i$$

where T_i is an indicator variable for treatment status and \mathbf{X}_i is a vector of *pre-treatment characteristics*

- Under this model, what is random?
- How do we interpret the coefficients on \mathbf{X}_i ?

Regression in Causal Analysis

- Imagine we are analyzing a *randomized* experiment with a regression using the following model:

$$Y_i = \alpha + \beta_1 \cdot T_i + \mathbf{X}_i^T \cdot \beta_2 + \epsilon_i$$

where T_i is an indicator variable for treatment status and \mathbf{X}_i is a vector of *pre-treatment characteristics*

- Under this model, what is random?
- How do we interpret the coefficients on \mathbf{X}_i ?
- How do we interpret the coefficient β_1 ?

Regression in an Observational Study

