



[Course](#) > [Module...](#) > [Pop Qui...](#) > Pop Qui...



## Pop Quiz #4

### Pop Quiz # 4

3/5 points (ungraded)

**In the following, write your BRACU ID# and section number first. After you submit the quiz, it will show that the ID# and Section numbers are wrong. Please ignore this messages. You score will based on the MCQs only.**

=====

Your Bracu ID #

19101239



Your Theory class section #:

☐ 1

Can't load web font TeX/Main/Regular

☐ 2



☐☐ 3☐ 4☐ 5☒ 6

Below are the MCQs:

=====

Q#1: In the Cauchy's interpolation formula, what would be value of

$$\left| \frac{f^{(n+1)}(\xi)}{(n+1)!} \right|$$

if  $f(x) = \tan(x) \in [-\pi/4, \pi/4]$  and  $n = 3$

☐ 0☐ 1☒  $\frac{1}{24}$ ☐ It can not be determined. Need to know the nodes.

Q#2: Which of the following is the Runge phenomenon?

- ☐ It is a phenomenon when nodes are chosen with unequal spacing.
- ☒ It is the diverging phenomena at the end points, even though at the middle it converges.
- ☐ As  $n \rightarrow \infty$   $p_n(x) \rightarrow f(x)$  always.
- ☐ It is a special phenomenon that is associated with quadratic function only.



Q#3: Why are Chebyshev nodes an optimal choice in Interpolation?

- ☒ Because it takes more nodes on the edges and less nodes in the middle.
- ☐ Because it takes nodes randomly.
- ☐ Because node selection is problem specific in this technique, ensuring minimum error in each interpolation problem.
- ☐ Because it takes equally spaced nodes.



Submit

◀ Previous

Next ▶



[About Us](#)

[BracU Home](#)

[USIS](#)

[Course Catalog](#)

Copyright - 2020

Can't load web font TeX/Main/Regular

