

ciliates of the sea urchin *Strongylocentrotus droebachiensis*. *J. Protozool.*, 1: 86–92.

4. Berger, J. & Lynn, D. H. 1984. Somatic and oral cortical ultrastructure of the plagiopylid ciliates *Lechriopyla mystax* Lynch, 1930 and *Plagiopyla minuta* Powers, 1933. *J. Protozool.*, 31: 433–443.

5. Borror, A. 1972. Tidal marsh ciliates (Protozoa): morphology, ecology, systematics. *Acta Protozool.*, 10: 29–71.

6. Corliss, J. O. 1979. *The Ciliated Protozoa*, 2nd ed. Pergamon Press, Oxford.

7. Czapik, A. 1975. Les observations sur *Trimyema compressum* Lackey (Ciliata, Trichostomata). *Acta Protozool.*, 13: 361–364.

8. Detcheva, R., Puytorac, P. de & Grolière, C. A. 1981. Some ultrastructural characteristics of the polysaprobic ciliate *Trimyema compressum*. *Trans. Am. Microsc. Soc.*, 100: 65–73.

9. Dragesco, J., Iftode, F. & Fryd-Versavel, G. 1974. Contribution à la connaissance de quelques ciliés holotriches rhabdophores; I. Prostomiens. *Protistologica*, 10: 59–75.

10. Fauré-Fremiet, E. 1962. Le genre *Trimyema* et les caractéristiques structurales des ciliés Trichostomatida. *J. Protozool.*, 9: 146–149.

11. Fernández-Galiano, D. 1976. Silver impregnation of ciliated protozoa: procedure yielding good results with the pyridinated silver carbonate method. *Trans. Am. Microsc. Soc.*, 95: 557–560.

12. Foissner, W. 1983. Taxonomische Studien über die Ciliaten des Großglocknergebietes (Hohe Tauern, Österreich). I. Familien Holophryidae, Prorodontidae, Plagiocampidae, Colepidae, Enchelyidae und Lacrymariidae nov. fam. *Ann. Naturhist. Mus. Wien.*, 84/B(1980): 49–85.

13. ———. 1984. *Infraciliatur, Silberliniensystem und Biometrie einiger neuer und wenig bekannter terrestrischer, limnischer und mariner Ciliaten (Protozoa, Ciliophora) aus den Klassen Kinetofragminophora, Colpodea und Polyhymenophora*. Stapfia, pp. 1–165.

14. Grain, J. 1966. Etude cytologique de quelques ciliés holotriches endocommensaux des ruminants et des équides (première partie). *Protistologica*, 2: 27–43.

15. Grolière, C. A., Puytorac, P. de & Grain, J. 1980. Observations de quelques espèces de ciliés endocommensaux d'échinides du Golfe du Mexique et de la mer des Antilles. *Protistologica*, 16: 233–239.

16. Jankowski, A. W. 1964. Morphology and evolution of Ciliophora III. Diagnoses and phylogenesis of 53 saproelobionts, mainly of the order Heterotrichida. *Arch. Protistenkd.*, 107: 185–294.

17. Jankowski, A. W. 1980. Conspectus of a new system of the phylum Ciliophora. *Proc. Acad. Sci. USSR*, 94: 103–121. (in Russian)

18. Kahl, A. 1926. Neue und wenig bekannte Formen der holotrichen und heterotrichen Ciliaten. *Arch. Protistenkd.*, 55: 197–438.

19. ———. 1933. Ciliata libera et ectocommensalia, in Grimpe, G. & Wagler, E., eds., *Die Tierwelt der Nord- und Ostsee*, Lief. 23 (Teil II, C3), Leipzig, pp. 29–46.

20. ———. 1935. Urtiere oder Protozoa I: Wimpertiere oder Ciliata (Infusoria), in Dahl, F., ed., *Die Tierwelt Dtl.* G. Fischer, Jena.

21. Lackey, J. B. 1925. Studies on the biology of sewage disposal. The fauna of Imhoff tanks. *New Jers. Agric. Exp. Stn. Bull.*, 417: 1–39.

22. Levine, N. D., Corliss, J. O., Cox, F. E. G., Deroux, G., Grain, J., Honigberg, B. M., Leedale, N. D., Loeblich, A. R., III, Lom, J., Lynn, D., Merinfeld, E. G., Page, F. C., Poljansky, G., Sprague, V., Vávra, J. & Wallace, F. G. 1980. A newly revised classification of the Protozoa. *J. Protozool.*, 27: 37–58.

23. Martín-González, A., Serrano, S. & Fernández-Galiano, D. 1986. *Urotricha vitrea* n. sp. (Ciliophora, Prorodontina): general morphology and cytological events during the conjugation process. *Can. J. Zool.*, 63: 1885–1891.

24. Muñoz, A., Téllez, M. C. & Fernández-Galiano, D. 1987. Morphology and infraciliature in *Urotricha nais* n. sp. and *Urotricha castalia* n. sp. (Ciliophora, Prorodontina). *Acta Protozool.*, 26: 197–204.

25. Puytorac, P. de, Batisse, A., Bohatier, J., Corliss, J. O., Deroux, G., Didier, P., Grolière, C. A., Hovasse, R., Iftode, F., Laval, M., Roque, M., Savoie, A. & Tuffrau, M. 1974. Proposition d'une classification d'un phylum Ciliophora Doflein, 1901. *C. R. Acad. Sci. Paris*, 278: 2799–2802.

26. Puytorac, P. de, Grain, J., Legendre, P. & Devaux, J. 1984. Essai d'application de l'analyse phénétique à la classification du Phylum des Ciliophora. *J. Protozool.*, 31: 496–507.

27. Rodrigues de Santa Rosa, M. 1976. Observations sur l'ultrastructure du cilié *Coleps hirtus* Nitzsch, 1817. *Protistologica*, 12: 205–216.

28. Serrano, S., Martín-González, A. & Fernández-Galiano, D. 1985. Cytological observations on the ciliate *Coleps hirtus* Nitzsch, 1817: vegetative cell, binary fission and conjugation. *Acta Protozool.*, 24: 77–84.

29. Small, E. B. & Lynn, D. H. 1985. Phylum Ciliophora, in Lee, J. J., Hutner, S. H. & Bovee, E. C., eds., *An Illustrated Guide to the Protozoa*. Society of Protozoologists, Lawrence, Kansas, pp. 393–575.

30. Tucolesco, J. 1962. Espèces nouvelles d'infusoires de la Mer Noire et des bassins salés paramarins. *Arch. Protistenkd.*, 106: 1–36.

31. Wilburt, N. & Schmall, G. 1976. Morphologie und Infraciliatur von *Coleps nolandi* Kahl, 1930. *Protistologica*, 12: 193–197.

Received 20 III 87; accepted 7 XII 87

Note to Establish as a New Species a Previously Reported Microsporidium of Philippine Prawns¹

GLORIA L. ENRIQUEZ* and VICTOR SPRAGUE**

*Natural Sciences Research Institute, University of the Philippines System, Diliman, Quezon City 3004, Philippines and

**Chesapeake Biological Laboratory, Solomons, Maryland 20688

ABSTRACT. A previously reported microsporidium in *Penaeus merguensis* de Man, 1888 (Decapoda, Penaeidae), in the Philippines has been identified as a species of *Agmasoma* Hazard & Oldacre, 1975, and named *Agmasoma aquinoae* n. sp. (Microsporida, Thelohaniidae).

A considerable amount of descriptive information regarding an unnamed "microsporidian parasite" of cultured prawns in the Philippines has already been reported (1–3). That parasite evidently belongs to the genus *Agmasoma* Hazard & Oldacre,

1975 (4), which resembles species of *Thelohania* Henneguy, 1892, but is distinguished by having an anisofilar polar tube. The purpose of this note is to summarize its taxonomic characters and to give it a name.

Agmasoma aquinoae n. sp.

Synonymy. "Microsporidian parasite" Enriquez, Baticados & Gacutan, 1980 (1). Baticados & Enriquez, 1982 (1, 3).

¹ Contribution No. 1877, Center for Environmental and Estuarine Studies of the University of Maryland.

Type host. *Penaeus merguensis* de Man, 1888 (Crustacea, Decapoda).

Site of infection. Oocytes.

Transmission. No data.

Relation to host cell. Various stages randomly dispersed in host cell cytoplasm. Meronts in direct contact with host cell cytoplasm. Sporont elaborates an envelope, isolating itself and subsequent sporulation stages from host cell cytoplasm. "... There was no apparent change in size or shape of infected oocytes. ... The parasite destroys and replaces the contents of the oocytes ..." (2).

Developmental characters. With one known sporulation sequence. Diplokarya present in meronts. Diplokaryon present in sporont, giving rise to single nuclei in sporoblasts and spores. Sporogony typically octosporoblastic (details of division process not fully clarified). "... During the sporoblast stage ... spore organelles ... start to form" (1) (see Remarks). Sporophorous vesicle present, delicate, persistent.

Spore morphology. Spores generally pyriform, many ovoid, some oblong, some teratological, most in the size range $1.5\text{--}2.2 \times 2.4\text{--}3.6 \mu\text{m}$, a few larger ones $2.2\text{--}3.6 \times 5.1\text{--}6.6 \mu\text{m}$ (fresh). Uninucleate. Exospore rather thin, endospore moderately thick. Polar tube anisofilar, being in coil with a thick proximal part forming about two turns and a thin distal part forming about five turns. Polaroplast fairly large, extending about to middle of spore; apparently with anterior laminate and posterior vesiculate parts (structural details not fully clarified). Posterior vacuole of moderate size.

Type locality. Pond of Southeast Asian Fisheries Development Center (SEAFDEC), Tigbauan, Iloilo, Philippines.

Differentiation of the taxon. The type and only other known species of *Agmasoma* is *A. penaei* (Sprague, 1950) Hazard & Oldacre, 1975 (4). This was originally *Thelohania penaei* Sprague, 1950 (5). It was found in the ovary (apparently also in male reproductive organs) of *Penaeus setiferus* (L.) collected along the Gulf coast of the United States. Its spores were originally reported to be pyriform, $2.2 \times 4.0 \mu\text{m}$ (fresh), with anisofilar polar tube about $70 \mu\text{m}$ long, the two parts being roughly equal in length. Using electron microscopy, Hazard & Oldacre (4), found that the coiled polar tube had two and one half turns in the broader proximal part and six in the narrower distal part. *Agmasoma aquinoae* is distinguished by having somewhat smaller spores, fewer turns in the coiled polar filament, a different host species, and a far removed locality.

Deposition of type specimens. Type materials of *A. aquinoae* not extant. Three slides of *Thelohania penaei* Sprague, 1950, syntype material, USNM No. 33639, have recently been deposited in the International

Protozoan Type Slide Collection, National Museum of Natural History, Smithsonian Institution, Washington, DC 20560.

Etymology. The specific name *aquinoae* is proposed in honor of Philippine President Corazon C. Aquino.

Remarks. Hazard & Oldacre (4) found in *Agmasoma penaei* a peculiar type of sporogony in which cytokinesis apparently had not yet begun after three nuclear divisions were complete and the spore walls had started to develop. That finding (much different from the finding in *A. aquinoae*) is not reliable, however, because the "... information came from electron photomicrographs of tissues taken from a shrimp packed in ice for many days, and this treatment may have produced abnormalities in the sporonts" (4). We expect that further observations, using suitable material, would show this species to have developed, like *A. aquinoae*, to the sporoblast stage before the spore organelles began to form. We also expect that adequate study would demonstrate the occurrence of meiosis in both species during sporogony.

LITERATURE CITED

1. Baticados, C. L. & Enriquez, G. L. 1982. Developmental stages of a microsporidian parasite of the white prawn, *Penaeus merguensis* de Man, 1888. *Nat. Appl. Sci. Bull.*, **34**: 255–270.
2. ——— 1982. Histopathology of microsporidian infection in white prawn, *Penaeus merguensis* de Man, 1888. *Nat. Appl. Sci. Bull.*, **34**: 299–313.
3. Enriquez, G. L., Baticados, C. L. & Gacutan, R. O. 1980. Microsporidian parasite of the white prawn *Penaeus merguensis* de Man: a preliminary report. *Nat. Appl. Sci. Bull.*, **32**: 319–325.
4. Hazard, E. I. & Oldacre, S. W. 1975. Revision of Microsporidia (Protozoa) close to *Thelohania*, with descriptions of one new family, eight new genera, and thirteen new species. *U.S. Dept. Agric. Tech. Bull.*, **1530**: i–v, 1–104.
5. Sprague, V. 1950. Notes on three microsporidian parasites of decapod Crustacea of Louisiana coastal waters. *Occ. Pap. Mar. Lab. La. St. Univ.*, **5**: 1–8.

Received 1 X 87; accepted 25 I 88

Persistence of Oral Structures in the Sexual Process of Amicronucleate *Paramecium tetraurelia*

STEPHEN F. NG

Zoology Department, University of Hong Kong, Hong Kong

ABSTRACT. In the sexual process, amicronucleate *Paramecium tetraurelia*, unlike micronucleates, fail to produce an oral apparatus, but resorb the pre-existing one. Exceptions were found in some amicronucleate cell lines in which about 1% of the cells possessed oral structures, including pieces of oral membranelles, sometimes complete with buccal cavity, after autogamy or conjugation. By following oral development in the sexual process in some detail, the present study supports the view that these oral structures are derived from the pre-existing oral apparatus and not newly developed from the oral primordium. The possible involvement of the micronucleus and the pre-existing oral apparatus in oral resorption is discussed. The possession of a functional oral apparatus after the sexual process may open up a new evolutionary avenue to the amicronucleates.

IN *Paramecium*, the oral apparatus is resorbed, and simultaneously a new one is produced during sexual reproduction (autogamy or conjugation) (Fig. 1). When conditions that lead to autogamy in micronucleate *P. tetraurelia* are applied to amicronucleates of this species, the macronucleus undergoes fragmentation and the pre-existing oral apparatus is resorbed, but a new one is not produced (Fig. 2). Stomatogenesis is arrested at the early stage of alignment of basal bodies into parallel rows

in the oral anarchic field (stage 2) (14, 16). The consequence of the sexual process for amicronucleates is thus astomy and death. The micronucleus is thus indispensable for stomatogenesis in the sexual cycle. Moreover, both of these reports (14, 16) recorded the presence in low frequency (<1%) of oral structures, including membranelles and sometimes the buccal cavity, after the sexual process in some of the amicronucleate cell lines. This generates interest as to whether such oral structures are relics