

---

---

# COGS 17 Week 5

— SPRING 2024, A03 —

---

---

# Problem Set for Today


- Link:

<https://docs.google.com/document/d/1vB2zNsDuZjfWN0OFOB9JqQthLZPJjP989ys9uTcEPo4/edit?usp=sharing>




**SCAN ME**

# Anatomy of the Ear




- Pinna -- Outer ear
- Tympanic Membrane -- eardrum, which transfers vibrations to >>
- Ossicles -- 3 smallest bones in body, which transfer vibrations to >>
- Oval window -- a membrane at the base of the upper chamber of >>
- Cochlea -- snail-shaped (coiled) tube with 3 fluid-filled chambers

# Organ of Corti




(a)



(b)


# Organ of Corti



(b)

- A structure in the cochlea of the inner ear which produces nerve impulses in response to sound vibrations
- The floor is Basilar Membrane, the ceiling is Tectorial membrane
- The auditory receptor cells, hair cells, are located between these two membranes
- Filled with a viscous, non-compressible,  $K^+$  rich fluid called endolymph


# Organ of Corti



(b)

- Vibration causes Basilar Membrane to move up and down and Tectorial Membrane to move left and right
- Cilia of Hair Cells bend, when toward longest cilium, **K<sup>+</sup> GATES OPEN**
- Na<sup>+</sup> **NOT** involved
- K<sup>+</sup> in >> Ca<sup>++</sup> in, NT out
- **Graded** response

# Place Coding




© 2001 Sinauer Associates, Inc.


- The **BASE** of the basilar membrane is narrow and stiff >> resonates to **HIGH** frequencies
- The **APEX** of the basilar membrane is wide and floppy >> resonates to **LOW** frequencies
- The more the Basilar Membrane resonates, the farther it moves >> the farther the Cilia will bend >> more NT release
- The **DISTRIBUTION** of NT release along Basilar Membrane that codes for Frequency

# Place Coding


High frequency sound = more response from Base than Apex



Low frequency sound = more response from Apex than Base





Mid frequency sound = more response from Mid-Membrane






# Temporal Coding



- The whole basilar membrane will vibrate at the rate of input
- However, Hair Cells communicate to Spiral Ganglions, which fire **ACTION POTENTIAL**, which limits the rate spiral ganglions can fire
- Maximum frequency ~1000 Hz
- Solution: Volley Principle -- No single Ganglion cell can code for a high rate, but a group of them working together can

# Temporal Coding




- Depends on Ganglions being **Phase Locked**
- The Ganglions can only fire at the same phase (e.g. peak) of the input frequency even when it is ready to fire
- These volleys occur at a rate that corresponds to the rate of the input frequency

# Localization




- Localization can use **Intensity** differences, **phase** differences, or **timing** differences to the two ears

# Localization via Intensity differences



- Sound at ear closer to source is slightly more intense than at other ear, because of the head shadow
- Works best for **higher** frequencies, since these most likely to be absorbed by head


# Localization via phase differences



Note PHASE difference at two ears


- For **lower** frequencies (longer wavelength), can detect difference in peak vs. trough of wave reaching two ears

# Localization via Timing Differences




- Onset signals race to same Superior Olive nucleus; Signal travels on axon with a branch to each Superior Olive cell in array

# Localization via Timing Differences




# Localization via Timing Differences






# Localization via Timing Differences




# Auditory Pathways -- Hair Cells




- Two types of hair cells
- Inner hair cells -- divergent connectivity, better for details
- Outer hair cells -- convergent connectivity, better for amplitude
- >> Spiral Ganglions

# Auditory Pathways -- Spiral Ganglion




- Axons of which form Auditory Nerve
- >> Cochlear Nucleus

# Auditory Pathways -- Cochlear Nucleus




- In medulla
- Monaural site (Left Cochlear Nucleus receives from left ear only, right from right only)
- Each Spiral Ganglion synapses on multiple cochlear nucleus cells
- Different types of cochlear nucleus cells: primary-like, build-up, onset, phaser
- >> Superior Olive

# Auditory Pathways -- Superior Olive




- In medulla
- Some Cochlear Nucleus axons cross-over to contralateral Superior Olive, others to ipsilateral
- First **binaural** site along pathway
- Critical for localization
- >> Inferior Colliculus

# Auditory Pathways -- Inferior Colliculus




- In tectum of midbrain
- Integrates with visual info in Superior Colliculus, help map source of sight/sound
- >> Thalamus

# Auditory Pathways -- Thalamus



- Medial Geniculate Nucleus (MGN)
- Includes Topological Maps
- Topological Maps -- Cells that respond to highest frequencies next to cells that respond to medium-high, next to medium, etc.
- >> Auditory Cortex


# Auditory Pathways -- Auditory Cortex




- A1 (Primary Auditory Cortex, along Lateral Sulcus of Temporal Cortex)
- A2 (Secondary Auditory Cortex, also in Temporal Lobe)



# A1 (Primary Auditory Cortex)




- In **MEDIAL** temporal cortex
- Responsible for Tonotopic/Amplitude Map




- Some cells respond best to simple (sine-wave) tones, others to more complex sounds (e.g. sounds that change freq)
- Other A1 areas respond per location

## A2 (Secondary Auditory Cortex)



- In **MEDIAL** temporal cortex
- Most respond best to complex sounds (familiar noises, speech sounds)
- >> Higher Auditory

# Auditory Pathways -- Higher Auditory



- Processes complex patterns; Integrates auditory input with other perceptual and cognitive activity
- Example: Wernicke's Area – Specialized for comprehending SPEECH
- Damage to Wernicke's area -- Wernicke's aphasia

# Questions?

Office Hours: Mon 5-6 pm

To get the section slides: <https://github.com/JasonC1217/COGS17-A03-Sp24>

OR:

