Sensitivity Analysis: Ranges and Reoptimization

An optimal plan answers one set of data. Sensitivity analysis tells us what survives when those data change.

> **Our route:** update a familiar dictionary → find allowable ranges → choose a reoptimization method → interpret the whole value function.


Bertsimas and Tsitsiklis, Chapter 5.

- A bakery example
- Right-hand sides
- Objective coefficients
- Solver reports
- Global value functions

One bakery, two products, three resources

Let x be bread production and y cake production. Quantities are continuous, measured in batches.

$$
\begin{aligned}
\max \quad & z = 2x + 3y \\
\text{s.t.} \quad & x + y \le 9 && \text{(hours)}, \\
& 2x + y \le 16 && \text{(flour)}, \\
& x + 2y \le 14 && \text{(sugar)}, \\
& x, y \ge 0.
\end{aligned}
$$



○ Original basis · ● Optimum
Shading: feasible plans · Dotted: profit line

The optimum is $(x, y) = (4, 5)$, with profit 23.

Hours and sugar are fully used; flour has 3 units left.

Start from an optimal dictionary we can check

Add nonnegative slacks: $x + y + s_1 = 9$, $2x + y + s_2 = 16$, $x + 2y + s_3 = 14$.

$$
\begin{aligned}
x &= 4 - 2s_1 + s_3, \\
y &= 5 + s_1 - s_3, \\
s_2 &= 3 + 3s_1 - s_3, \\
z &= 23 - s_1 - s_3.
\end{aligned}
$$

> Basic variables: x, y, s_2. Nonbasic variables: s_1, s_3. Setting the latter to zero gives $(4, 5, 3)$.

This is **maximization**: the nonbasic objective coefficients $-1, -1$ certify optimality. Increasing a nonnegative slack cannot improve profit.

After any change, check two separate things

Question	For this maximizing dictionary
Primal feasibility	Are all basic values nonnegative?
Dual feasibility	Are all nonbasic objective coefficients ≤ 0?

> If both checks pass, the current basis is still optimal.

A change in a right-hand side affects the basic values. A change in a profit coefficient affects the objective row.

We will do the arithmetic first, then summarize it in matrix form.

Change the flour limit

Maximize $2x + 3y$; $x + y \le 9,\ 2x + y \le 16 + \delta,\ x + 2y \le 14,\ x, y \ge 0$.


Original dictionary

$$x = 4 - 2s_1 + s_3,$$
$$y = 5 + s_1 - s_3,$$
$$s_2 = 3 + 3s_1 - s_3,$$
$$z = 23 - s_1 - s_3.$$

After the flour change

$$x = 4 - 2s_1 + s_3,$$
$$y = 5 + s_1 - s_3,$$
$$s_2 = 3 + \delta + 3s_1 - s_3,$$
$$z = 23 - s_1 - s_3.$$

Set $s_1 = s_3 = 0$. Feasible exactly when $3 + \delta \ge 0$: $\delta \ge -3$. Profit stays 23.



○ Original basis · ● Optimum
Shading: feasible plans · Dotted: profit line

Shown: $\delta = -3$. Flour limit: 13. Original basis: 4 bread, 5 cakes, 0 unused flour (feasible). Profit 23; optimum 23.

Change the hours limit

Maximize $2x + 3y$; $x + y \le 9 + \delta,\ 2x + y \le 16,\ x + 2y \le 14,\ x, y \ge 0$.


Original dictionary

$$x = 4 - 2s_1 + s_3,$$
$$y = 5 + s_1 - s_3,$$
$$s_2 = 3 + 3s_1 - s_3,$$
$$z = 23 - s_1 - s_3.$$

After the hours change

$$x = 4 + 2\delta - 2s_1 + s_3,$$
$$y = 5 - \delta + s_1 - s_3,$$
$$s_2 = 3 - 3\delta + 3s_1 - s_3,$$
$$z = 23 + \delta - s_1 - s_3.$$

Require $4 + 2\delta,\ 5 - \delta,\ 3 - 3\delta \ge 0$. Thus $-2 \le \delta \le 1$; profit is $23 + \delta$.



○ Original basis · ● Optimum
Shading: feasible plans · Dotted: profit line

Shown: $\delta = 1$. Hours limit: 10. Original basis: 6 bread, 4 cakes, 0 unused flour (feasible). Profit 24; optimum 24.

Practice: change the sugar limit

Maximize $2x + 3y$; $x + y \le 9,\ 2x + y \le 16,\ x + 2y \le 14 + \delta,\ x, y \ge 0.$


Original dictionary

$$x = 4 - 2s_1 + s_3,$$
$$y = 5 + s_1 - s_3,$$
$$s_2 = 3 + 3s_1 - s_3,$$
$$z = 23 - s_1 - s_3.$$

Keep $s_1 = s_3 = 0$. Find the **updated dictionary**, the allowable interval for δ, and the profit.

Use $x + y = 9 - s_1$, $x + 2y = 14 + \delta - s_3$, then $s_2 = 16 - 2x - y$.

Take two minutes. Include unused flour in the feasibility test.



Shown: $\delta = 0$. Sugar limit: 14. Find the updated dictionary and its allowable interval.

Sugar can decrease by 3 or increase by 4

Maximize $2x + 3y$; $x + y \le 9,\ 2x + y \le 16,\ x + 2y \le 14 + \delta,\ x, y \ge 0.$


Original dictionary

$$x = 4 - 2s_1 + s_3,$$
$$y = 5 + s_1 - s_3,$$
$$s_2 = 3 + 3s_1 - s_3,$$
$$z = 23 - s_1 - s_3.$$

After the sugar change

$$x = 4 - \delta - 2s_1 + s_3,$$
$$y = 5 + \delta + s_1 - s_3,$$
$$s_2 = 3 + \delta + 3s_1 - s_3,$$
$$z = 23 + \delta - s_1 - s_3.$$

Feasible exactly for $-3 \le \delta \le 4$: flour slack reaches zero at –3; bread at 4. Profit: $23 + \delta$.



○ Original basis · ● Optimum
Shading: feasible plans · Dotted: profit line

Shown: $\delta = 4$. Sugar limit: 18. Original basis: 0 bread, 9 cakes, 7 unused flour (feasible). Profit 27; optimum 27.

Recall the bakery and its optimal dictionary

x: bread batches; y: cake batches. Maximize $z = 2x + 3y$.

Original limits: $x + y \le 9$ (hours), $2x + y \le 16$ (flour), $x + 2y \le 14$ (sugar); $x, y \ge 0$.

Introduce nonnegative slacks s_1, s_2, s_3, the unused hours, flour, and sugar:

$$x + y + s_1 = 9, \qquad 2x + y + s_2 = 16, \qquad x + 2y + s_3 = 14.$$

Original optimal dictionary

$$\begin{aligned} x &= 4 - 2s_1 + s_3, \\ y &= 5 + s_1 - s_3, \\ s_2 &= 3 + 3s_1 - s_3, \\ z &= 23 - s_1 - s_3. \end{aligned}$$

Basic: x, y, s_2. Nonbasic: s_1, s_3. Set $s_1 = s_3 = 0$: $(x, y, s_2) = (4, 5, 3)$, profit 23. The objective coefficients $-1, -1$ are nonpositive, as required for maximization.

Update the hours dictionary, then set δ = 2

Maximize $2x + 3y$; $x + y \le 9 + \delta,\ 2x + y \le 16,\ x + 2y \le 14,\ x, y \ge 0.$


Old hours slack: $s_1^{\text{old}} = 9 - x - y.$

New hours slack: $s_1 = 9 + \delta - x - y.$

Substitute $s_1^{\text{old}} = s_1 - \delta$ in every original row:

Variable	With δ	$\delta = 2$
x	$4 + 2\delta - 2s_1 + s_3$	$8 - 2s_1 + s_3$
y	$5 - \delta + s_1 - s_3$	$3 + s_1 - s_3$
s_2	$3 - 3\delta + 3s_1 - s_3$	$-3 + 3s_1 - s_3$
z	$23 + \delta - s_1 - s_3$	$25 - s_1 - s_3$

At $\delta = 2$, $s_1 = s_3 = 0$ gives $s_2 = -3$. **Leave s_2; enter s_1**, the only variable with a positive coefficient in that row. Reduced costs stay $-1, -1$: dual feasible.



○ Original basis · ● Optimum
Shading: feasible plans · Dotted: profit line

Shown: $\delta = 2$. Hours limit: 11. Original basis: 8 bread, 3 cakes, -3 unused flour (infeasible). Profit 25; optimum 24.

One dual-simplex pivot restores feasibility

Maximize $2x + 3y$, with $x + y \le 11$, $2x + y \le 16$, $x + 2y \le 14$, $x, y \ge 0$.

From $s_2 = -3 + 3s_1 - s_3$, solve $s_1 = 1 + \frac{1}{3}s_2 + \frac{1}{3}s_3$ and substitute in every row.

Before pivot	After pivot
$x = 8 - 2s_1 + s_3$	$x = 6 - \frac{2}{3}s_2 + \frac{1}{3}s_3$
$y = 3 + s_1 - s_3$	$y = 4 + \frac{1}{3}s_2 - \frac{2}{3}s_3$
$s_2 = -3 + 3s_1 - s_3$	$s_1 = 1 + \frac{1}{3}s_2 + \frac{1}{3}s_3$
$z = 25 - s_1 - s_3$	$z = 24 - \frac{1}{3}s_2 - \frac{4}{3}s_3$

Set $s_2 = s_3 = 0$: $(x, y, s_1) = (6, 4, 1)$, profit **24**. All basic values are nonnegative and all nonbasic objective coefficients are nonpositive.

The old value 25 was an upper bound; it was not an attainable profit.

Two allowable changes may fail together

Bakery: maximize $z = 2x + 3y$, with $x + y \leq 9$, $2x + y \leq 16$, $x + 2y \leq 14$, $x, y \geq 0$.

Independent variation: return to basis (x, y, s_2), with $s_1 = s_3 = 0$. Change hours by δ_1 and flour by δ_2; keep sugar at 14.

From $x + y = 9 + \delta_1$, $x + 2y = 14$, and $s_2 = 16 + \delta_2 - 2x - y$:

Thus $x = 4 + 2\delta_1, \quad y = 5 - \delta_1, \quad s_2 = 3 - 3\delta_1 + \delta_2$.

Change from the original bakery	x	y	Flour slack s_2
None	4	5	3
Hours $9 \to 10$ only	6	4	0
Flour $16 \to 13$ only	4	5	0
Both changes together	6	4	-3

With 10 hours, the plan $(6, 4)$ uses $2(6) + 4 = 16$ units of flour. Only 13 are available after both changes: **this basis is infeasible**. The changed LP may still be feasible.

The 100% rule gives a sufficient check

Bakery: maximize $z = 2x + 3y$, with $x + y \le 9$, $2x + y \le 16$, $x + 2y \le 14$, $x, y \ge 0$.

Original basis: (x, y, s_2). Divide each change's magnitude by its allowance in the table; add.

Resource change	One-at-a-time allowance	Fraction used
Hours $9 \to 9.5$	Increase by 1: $9 \to 10$	$0.5/1 = 50\%$
Flour $16 \to 15$	Decrease by 3: $16 \to 13$	$1/3 \approx 33.3\%$

Sum of fractions: $R = \frac{0.5}{1} + \frac{1}{3} = \frac{5}{6} < 1$.

$R \le 1$ guarantees a feasible basis. For hours $+0.5$ and flour -1:

$$x = 5, \quad y = 4.5, \quad s_2 = 3 - 3(0.5) - 1 = 0.5.$$

The objective coefficients remain $-1, -1$, so this basis is also **optimal**; profit is 23.5.

Percentages refer to the **allowable changes**, not the original resource totals. If $R > 1$, check the basic values directly; the rule gives no conclusion.

Why the rule works · A feasible example above 100%

Change bread profit using the original dictionary

Start again from the original bakery: hours 9, flour 16, sugar 14; maximize $2x + 3y$.

Change bread profit from 2 to $2 + \theta$: $z_{\text{new}} = z_{\text{old}} + \theta x$. Substitute the original row for x.

Variable	Original dictionary	Updated dictionary
x	$4 - 2s_1 + s_3$	$4 - 2s_1 + s_3$
y	$5 + s_1 - s_3$	$5 + s_1 - s_3$
s_2	$3 + 3s_1 - s_3$	$3 + 3s_1 - s_3$
z	$23 - s_1 - s_3$	$23 + 4\theta + (-1 - 2\theta)s_1 + (-1 + \theta)s_3$

The three constraint rows are unchanged. For maximization, require $-1 - 2\theta \leq 0$ and $-1 + \theta \leq 0$: $-\frac{1}{2} \leq \theta \leq 1$, or $1.5 \leq c_x \leq 3$.

Within this interval, quantities stay $(4, 5)$ and profit becomes $23 + 4\theta$.

Selling unused hours changes one objective coefficient

Start again from the original bakery: hours 9, flour 16, sugar 14; maximize $2x + 3y$.

Suppose unused hours s_1 can be sold for κ per unit. Then $z_{\text{new}} = z_{\text{old}} + \kappa s_1$.

Variable	Original dictionary	Updated dictionary
x	$4 - 2s_1 + s_3$	$4 - 2s_1 + s_3$
y	$5 + s_1 - s_3$	$5 + s_1 - s_3$
s_2	$3 + 3s_1 - s_3$	$3 + 3s_1 - s_3$
z	$23 - s_1 - s_3$	$23 + (\kappa - 1)s_1 - s_3$

The original basis remains optimal exactly when $\kappa \leq 1$. Its profit stays 23 because $s_1 = 0$. For $\kappa > 1$, **primal simplex** can improve the still-feasible plan.

Change both profits using the same original rows

Start again from the original bakery: hours 9, flour 16, sugar 14; maximize $2x + 3y$.

Set profits to $2 + \theta_1$ and $3 + \theta_2$. Use $z_{\text{new}} = z_{\text{old}} + \theta_1 x + \theta_2 y$.

Variable	Original dictionary	Updated dictionary
x	$4 - 2s_1 + s_3$	$4 - 2s_1 + s_3$
y	$5 + s_1 - s_3$	$5 + s_1 - s_3$
s_2	$3 + 3s_1 - s_3$	$3 + 3s_1 - s_3$
z	$23 - s_1 - s_3$	$23 + 4\theta_1 + 5\theta_2 + (-1 - 2\theta_1 + \theta_2)s_1 + (-1 + \theta_1 - \theta_2)s_3$

Both objective coefficients must be nonpositive: $\theta_1 - 1 \le \theta_2 \le 1 + 2\theta_1$. If only cake profit changes, $-1 \le \theta_2 \le 1$, so $2 \le c_y \le 4$.

The bakery dictionary in matrix form

Bakery: maximize $z = 2x + 3y$, with $x + y \le 9$, $2x + y \le 16$, $x + 2y \le 14$, $x, y \ge 0$.

Add slacks $s_1, s_2, s_3 \ge 0$ for hours, flour, sugar. Keep basis (x, y, s_2); its columns form B.

$$B = \begin{pmatrix} 1 & 1 & 0 \\ 2 & 1 & 1 \\ 1 & 2 & 0 \end{pmatrix}, \qquad B^{-1} = \begin{pmatrix} 2 & 0 & -1 \\ -1 & 0 & 1 \\ -3 & 1 & 1 \end{pmatrix}.$$

Original dictionary rows	After multiplying the equalities by B^{-1}
$x = 4 - 2s_1 + s_3,$ $y = 5 + s_1 - s_3,$ $s_2 = 3 + 3s_1 - s_3,$	$\begin{pmatrix} x \\ y \\ s_2 \end{pmatrix} = \begin{pmatrix} 4 \\ 5 \\ 3 \end{pmatrix} - \begin{pmatrix} 2 & -1 \\ -1 & 1 \\ -3 & 1 \end{pmatrix} \begin{pmatrix} s_1 \\ s_3 \end{pmatrix}.$

The constant vector is $B^{-1}(9, 16, 14)^\top$. The two coefficient columns are $B^{-1}A_{s_1}$ and $B^{-1}A_{s_3}$, subtracted from it.

$$z = 2x + 3y = 23 - s_1 - s_3.$$

Dual variables and reduced costs have different roles

Bakery: maximize $z = 2x + 3y$, with $x + y \le 9$, $2x + y \le 16$, $x + 2y \le 14$, $x, y \ge 0$.

Original optimal dictionary

$$x = 4 - 2s_1 + s_3, \qquad y = 5 + s_1 - s_3,$$
$$s_2 = 3 + 3s_1 - s_3, \qquad z = 23 - s_1 - s_3.$$

Basis: (x, y, s_2), with costs $c_{\mathcal{B}} = (2, 3, 0)^\top$.

$$p^\top = c_{\mathcal{B}}^\top B^{-1} = (2, 3, 0)B^{-1} = (1, 0, 1) \quad \text{(hours, flour, sugar)}.$$

Let $\Delta b = (\delta_1, \delta_2, \delta_3)^\top$ change those limits. Write $x_{\mathcal{B}} = (x, y, s_2)^\top$.

$$x_{\mathcal{B}}^{\text{new}} = (4, 5, 3)^\top + B^{-1}\Delta b,$$
$$z_{\text{opt}}^{\text{new}} = 23 + p^\top \Delta b = 23 + \delta_1 + \delta_3.$$

Nonbasic variables are zero. This basis stays optimal if its basic values stay nonnegative.

Reduced costs are the coefficients in z: $\bar{c}_{s_1} = 0 - p_1 = -1$, $\bar{c}_{s_3} = 0 - p_3 = -1$.

Derive the dictionary for a new pastry

With pastry t: maximize $z = 2x + 3y + 4t$.

$x + y + t \le 9$ (hours), $2x + y + 2t \le 16$ (flour), $x + 2y + t \le 14$ (sugar); $x, y, t \ge 0$.

t counts pastry batches; each earns 4 and uses $(1, 2, 1)$ resources. Write the inequalities as equalities using slacks $s_1, s_2, s_3 \ge 0$. Keep basis (x, y, s_2).

Subtract hours from sugar, then substitute into hours and flour:

$$\begin{aligned}(x + 2y + t + s_3) - (x + y + t + s_1) &= 14 - 9,\\ y &= 5 + s_1 - s_3,\\ x = 9 - y - t - s_1 &= 4 - 2s_1 + s_3 - t,\\ s_2 = 16 - 2x - y - 2t &= 3 + 3s_1 - s_3.\end{aligned}$$

$$\begin{aligned}z &= 2(4 - 2s_1 + s_3 - t) + 3(5 + s_1 - s_3) + 4t\\ &= 23 - s_1 - s_3 + 2t.\end{aligned}$$

No pivot yet: s_1, s_3, t are nonbasic. Setting them to zero gives the feasible plan $(x, y, t) = (4, 5, 0)$.

The matrix calculation gives the same new column

With pastry t: maximize $z = 2x + 3y + 4t$.

$x + y + t \le 9$ (hours), $2x + y + 2t \le 16$ (flour), $x + 2y + t \le 14$ (sugar); $x, y, t \ge 0$.

Dictionary before any pastry pivot

$$x = 4 - 2s_1 + s_3 - t, \qquad y = 5 + s_1 - s_3,$$
$$s_2 = 3 + 3s_1 - s_3, \qquad z = 23 - s_1 - s_3 + 2t.$$

For basis (x, y, s_2) and pastry column $A_t = (1, 2, 1)^\top$, the new coefficients are $-B^{-1}A_t$:

$$B^{-1}A_t = \begin{pmatrix} 2 & 0 & -1 \\ -1 & 0 & 1 \\ -3 & 1 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}.$$

$$\bar{c}_t = 4 - c_{\mathcal{B}}^\top B^{-1}A_t = 4 - p^\top A_t = 4 - (1 + 0 + 1) = 2.$$

One pastry replaces one bread: $4 - 2 = 2$ more profit. Use primal simplex: two pivots

What if the pastry needs more hours?

With pastry t: maximize $z = 2x + 3y + 4t$.

$x + y + (1 + \delta)t \le 9$ (hours), $2x + y + 2t \le 16$ (flour), $x + 2y + t \le 14$ (sugar); $x, y, t \ge 0$.

Before any pivot: keep basis (x, y, s_2) and $p = (1, 0, 1)$. Only pastry hours rise, by $\delta \ge 0$.

$$\bar{c}_t^{\text{new}} = 4 - \big[1(1 + \delta) + 0(2) + 1(1)\big] = 2 - \delta.$$

Extra hours δ	Reduced cost $2 - \delta$	Effect of increasing t
0	2	Profit increases.
2	0	Profit stays constant.
3	–1	Profit decreases.

At $t = 0$, plan $(4, 5, 0)$ stays feasible. The other reduced costs stay $-1, -1$. For $\delta \ge 2$, all are nonpositive: this plan is optimal.

Derive the reduced-cost change

With pastry t: maximize $z = 2x + 3y + 4t$.

$x + y + (1+\delta)t \le 9$ (hours), $2x + y + 2t \le 16$ (flour), $x + 2y + t \le 14$ (sugar); $x, y, t \ge 0$.

Generalize the pastry calculation: a_{ij} is the coefficient of variable x_j in constraint i. Below, sum over all constraint rows k.

Change only a_{ij} to $a_{ij} + \delta$, with x_j **nonbasic**. Then $B, c_{\mathcal{B}}$ stay fixed, so $p^\top = c_{\mathcal{B}}^\top B^{-1}$ stays fixed.

$$\begin{aligned} \bar{c}_j^{\text{new}} &= c_j - \left(\textstyle\sum_k p_k a_{kj} + p_i \delta\right) \\ &= \left(c_j - \textstyle\sum_k p_k a_{kj}\right) - \delta p_i. \end{aligned}$$

$$\bar{c}_j^{\text{new}} = \bar{c}_j - \delta p_i.$$

Feasibility is unchanged: $x_j = 0$, so the changed coefficient multiplies zero. For maximization, check all nonbasic reduced costs are nonpositive.

If x_j is **basic**, B changes: recompute basic values and reduced costs. Worked basic-column example

Start again with two products: add a bread limit

Bakery: maximize $z = 2x + 3y$, with $x + y \le 9$, $2x + y \le 16$, $x + 2y \le 14$, $x, y \ge 0$.

Original optimal dictionary

$$x = 4 - 2s_1 + s_3, \qquad y = 5 + s_1 - s_3,$$
$$s_2 = 3 + 3s_1 - s_3, \qquad z = 23 - s_1 - s_3.$$

Separate variation: no pastry or recipe change. Add $x \le 3$. The original plan has $x = 4$, so it violates the new constraint.

$$x + s_4 = 3, \quad s_4 \ge 0; \qquad s_4 = 3 - (4 - 2s_1 + s_3) = -1 + 2s_1 - s_3.$$

Use basis (x, y, s_2, s_4), with $s_1 = s_3 = 0$. The new basic value $s_4 = -1$ is negative, while reduced costs remain $-1, -1$. **Use dual simplex.**

If an added constraint is satisfied by the old optimum, that solution remains optimal.

Complete repair pivot

The failed check tells us how to continue

Basic values feasible?	Reduced costs have correct signs?	Next step
Yes	Yes	Current basis remains optimal.
Yes	No	Primal simplex.
No	Yes	Dual simplex.
No	No	Repair feasibility, then optimize; Phase I is available.

> Outside a sensitivity range, reoptimize. A failed basis test alone establishes neither infeasibility nor unboundedness.

Bakery summary: ranges for the original optimal basis

Return to $\max 2x + 3y$, limits $(9, 16, 14)$, and plan $(4, 5)$. These are the **original** model's one-at-a-time ranges:

Resource	RHS	Slack	Dual variable	Absolute RHS interval
Hours	9	0	1	$[7, 10]$
Flour	16	3	0	$[13, \infty)$
Sugar	14	0	1	$[11, 18]$

Product	Quantity	Profit coefficient	Reduced cost	Absolute profit interval
Bread	4	2	0	$[1.5, 3]$
Cake	5	3	0	$[2, 4]$

Intervals give parameter values, not increments; change only one value at a time.

> **Local bakery calculations end here.** Next: verify this report in software, then study changes across several optimal bases.

Software verification: solve the original bakery

```
import gurobipy as gp
from gurobipy import GRB
m = gp.Model("bakery")
x = m.addVar(lb=0, name="bread")
y = m.addVar(lb=0, name="cake")
m.setObjective(2*x + 3*y, GRB.MAXIMIZE)
m.addConstr(x + y <= 9, name="hours")
m.addConstr(2*x + y <= 16, name="flour")
m.addConstr(x + 2*y <= 14, name="sugar")
m.Params.Method = 1  # dual simplex: obtain a basis
m.optimize()
```

The solution is continuous; these are batches, not integer counts.

Software verification: read the sensitivity report

```
if m.Status == GRB.OPTIMAL:
    for row in m.getConstrs():
        print(row.ConstrName, row.Slack, row.Pi,
              row.SARHSLow, row.SARHSUp)
    for var in m.getVars():
        print(var.VarName, var.X, var.RC,
              var.SAObjLow, var.SAObjUp)
```

`Pi`: dual variable; RC: reduced cost. SARHS… and `SAObj`…: RHS and objective-coefficient limits. Sensitivity ranges require a basic solution.

At degeneracy, another optimal basis may report different valid ranges. A zero dual variable does not imply that the constraint is inactive.

Gurobi attribute reference

From one optimal basis to the whole value function

New topic: we have finished changes that preserve one basis. Now allow the optimal basis to change, and reuse the bakery to study the full optimal-value function.

To match the textbook convention, now write the bakery as **minimization** of $f = -z = -2x - 3y$. The feasible set is unchanged.

For a standard-form minimization LP, define $F(b) = \min\{c^\top x : Ax = b,\ x \geq 0\}$, wherever the optimum is finite.

$$F(b) = \max_{A^\top p \leq c} p^\top b.$$

Each feasible dual vector supplies a linear lower bound. The best such bound is a convex, piecewise-linear value function.


The bakery's maximum profit has the opposite sign, so it is concave in resource limits.

The marginal value changes when a new resource becomes scarce

Minimize $-2x - 3y$ with $x + y \le T$, $2x + y \le 16$, $x + 2y \le 14$, $x, y \ge 0$, $T \ge 0$.

$$F(T) = \begin{cases} -3T, & 0 \le T \le 7, \\ -T - 14, & 7 \le T \le 10, \\ -24, & T \ge 10. \end{cases}$$

At $T = 9$, the slope is -1: one more hour reduces negative profit by 1, or increases profit by 1.



The dashed line –T–14 is the global lower bound supplied by the dual solution at T=9.

A dual solution gives a supporting slope even at a corner

For minimization, an optimal dual vector p^* at b^* always gives

$$F(b) \geq F(b^*) + (p^*)^\top (b - b^*).$$

A vector satisfying this inequality is called a **subgradient**: its affine prediction is a global lower bound.

At $T = 7$ in our graph, left slope is -3 and right slope is -1. Every slope between them supports the graph; there is no single two-sided derivative.

At differentiable points, this supporting vector is the gradient. At corners, do not treat one reported dual value as the unique marginal change in every direction.

Full statement and proof

Changing costs selects among fixed feasible plans

Keep the original bakery constraints and minimize $-qx - 3y$, where q is bread's profit coefficient.

$$G(q) = \min\{-21,\ -4q - 15,\ -7q - 6,\ -8q\} \quad (q \geq 0).$$

> Each expression is one feasible vertex's cost. The minimum of these affine functions is concave and piecewise linear.

As q crosses $1.5, 3, 6$, the selected production plan changes.



Each line is the cost of a fixed feasible production plan. The minimum selects the best plan.

Trace the breakpoints instead of solving unrelated cases

For minimization with costs $c + \theta d$, a fixed basis has reduced costs

$$\bar{c}_j(\theta) = c_j + \theta d_j - (c_{\mathcal{B}} + \theta d_{\mathcal{B}})^\top B^{-1} A_j.$$

> Find the interval where all required signs hold. At an endpoint, pivot to a basis valid on the next interval. Ties require the anticycling care already studied.

Cost changes leave primal feasibility intact: use primal simplex. RHS changes leave reduced costs intact: use dual simplex.

Worked breakpoint pivot in the bakery

Turn a sensitivity calculation into a purchasing decision

In the bakery, the current dual variables are $(1, 0, 1)$, with hours 9, flour 16, sugar 14.

> A supplier offers 0.5 more hours for total cost 0.3. The change is inside the hours range. Extra operating profit is $1(0.5) = 0.5$, giving net gain 0.2.

An offer of 3 extra hours needs reoptimization. The old plan's formula would predict 26, but maximum profit is only 24, so the total gain is 1.

Compare money with money, state the range, and include procurement cost separately from operating profit.

Different production policies can change the scarce resource

In computer production, configurations consume memory boards and disk drives. Demand requirements and permitted substitute boards determine the feasible plans.

Question	Sensitivity analysis helps answer
Buy more of which component?	Compare dual values, units, and allowable increases.
Allow a substitute component?	Add its activity or column and check its reduced cost.
Change a demand policy?	Rebuild the affected constraints and reoptimize.

> A resource price belongs to a particular model and basis. A price reported under one policy cannot automatically be used under another.

Practice: decide which optimality check survives

Bakery: maximize $z = 2x + 3y$, with $x + y \le 9$, $2x + y \le 16$, $x + 2y \le 14$, $x, y \ge 0$.

Change	Your task
Hours increase by 0.5	Find the quantities, slack, and new profit.
Bread profit increases to 3.2	Decide whether this basis remains optimal.
Add x ≤ 3	Choose a reoptimization method and explain why.

Work in pairs for three minutes.

$$
\begin{aligned}
x &= 4 - 2s_1 + s_3, \\
y &= 5 + s_1 - s_3, \\
s_2 &= 3 + 3s_1 - s_3, \\
z &= 23 - s_1 - s_3.
\end{aligned}
$$

Practice answers: feasibility and objective signs play different roles

Change	Result
Hours +0.5	$(x, y, s_2) = (5, 4.5, 1.5)$; profit 23.5.
Bread profit 3.2	$\theta = 1.2$; coefficient of s_3 becomes $0.2 > 0$. Use primal simplex.
Add x ≤ 3	New slack $s_4 = -1$; old reduced costs still have correct signs. Use dual simplex.

> Always identify the changed data and the surviving property before choosing an algorithm.

Study guide: a range is a claim about one basis

Be able to: update a dictionary after an RHS or cost change; check combined changes; interpret a solver report; choose primal or dual simplex; distinguish local predictions from global value functions.

Read Bertsimas and Tsitsiklis, Chapter 5. The appendix contains complete derivations, a second minimization example, and proofs of the global statements.

- Basis formulas
- 100% rule proof
- Changing A
- Value-function proofs
- Second worked example

Appendix: calculations and proofs

These pages retain the algebra behind the examples and establish the general statements.

- Derive the bakery dictionary
- Complete reoptimization pivots
- Matrix formulas and simultaneous changes
- Convexity, concavity, and supporting slopes
- A minimization example

Derive the optimal dictionary by subtraction

$$x + y + s_1 = 9, \qquad x + 2y + s_3 = 14.$$

Subtract the first equality from the second, then substitute back:

$$y = 5 + s_1 - s_3, \qquad x = 9 - s_1 - y = 4 - 2s_1 + s_3.$$

$$s_2 = 16 - 2x - y = 3 + 3s_1 - s_3,$$
$$z = 2x + 3y = 23 - s_1 - s_3.$$

All variables are nonnegative. The objective is at most 23, attained when $s_1 = s_3 = 0$.

Derive all RHS changes in one dictionary

Write limits $(9+\delta_1, 16+\delta_2, 14+\delta_3)$. Subtract the first resource equation from the third, then substitute:

$$
\begin{aligned}
x &= 4 + 2\delta_1 - \delta_3 - 2s_1 + s_3,\\
y &= 5 - \delta_1 + \delta_3 + s_1 - s_3,\\
s_2 &= 3 - 3\delta_1 + \delta_2 + \delta_3 + 3s_1 - s_3,\\
z &= 23 + \delta_1 + \delta_3 - s_1 - s_3.
\end{aligned}
$$

> The RHS changes only constants. The objective coefficients of nonbasic variables remain –1, –1.

Full dictionary after adding a product

Maximize $2x + 3y + 4t$, subject to $x + y + t \leq 9$, $2x + y + 2t \leq 16$, $x + 2y + t \leq 14$, $x, y, t \geq 0$.

$$\begin{aligned} x &= 4 - 2s_1 + s_3 - t, \\ y &= 5 + s_1 - s_3, \\ s_2 &= 3 + 3s_1 - s_3, \\ z &= 23 - s_1 - s_3 + 2t. \end{aligned}$$

Increase t. Only x decreases, so the step is 4: t enters and x leaves.

The new product replaces bread in one pivot

$$
\begin{aligned}
t &= 4 - x - 2s_1 + s_3, \\
y &= 5 + s_1 - s_3, \\
s_2 &= 3 + 3s_1 - s_3, \\
z &= 31 - 2x - 5s_1 + s_3.
\end{aligned}
$$

The point $(x, y, t) = (0, 5, 4)$ has profit 31, but the positive s_3 coefficient means it is **not yet optimal**.

Let s_3 enter. Both y and s_2 decrease; s_2 reaches zero first at step 3.

$$
(x, y, t) = (0, 2, 7), \qquad z = 34.
$$

Final dictionary and certificate

Finish the new-product optimization

From $s_2 = 3 + 3s_1 - s_3$, solve $s_3 = 3 + 3s_1 - s_2$ and substitute:

$$
\begin{aligned}
t &= 7 - x + s_1 - s_2, \\
y &= 2 - 2s_1 + s_2, \\
s_3 &= 3 + 3s_1 - s_2, \\
z &= 34 - 2x - 2s_1 - s_2.
\end{aligned}
$$

All reduced profits are nonpositive. The optimum is $(0, 2, 7)$, with profit 34. The dual solution $p = (2, 1, 0)$ gives the same bound $9(2) + 16(1) = 34$.

Complete repair after adding x ≤ 3

The new row is $s_4 = -1 + 2s_1 - s_3$. Let s_4 leave and s_1 enter:

$$
\begin{aligned}
s_1 &= \tfrac{1}{2} + \tfrac{1}{2}s_3 + \tfrac{1}{2}s_4, \\
x &= 3 - s_4, \\
y &= \tfrac{11}{2} - \tfrac{1}{2}s_3 + \tfrac{1}{2}s_4, \\
s_2 &= \tfrac{9}{2} + \tfrac{1}{2}s_3 + \tfrac{3}{2}s_4, \\
z &= \tfrac{45}{2} - \tfrac{3}{2}s_3 - \tfrac{1}{2}s_4.
\end{aligned}
$$

Thus $(x, y) = (3, 5.5)$ is optimal with profit 22.5. The new constraint reduces the best profit.

Basis formulas: equality constraints first

For $\min c^\top x$ subject to $Ax = b, x \geq 0$, partition columns into invertible B and nonbasic block N.

$$Bx_{\mathcal{B}} + Nx_{\mathcal{N}} = b \quad \Rightarrow \quad x_{\mathcal{B}} = B^{-1}b - B^{-1}Nx_{\mathcal{N}}.$$

$$c^\top x = c_{\mathcal{B}}^\top B^{-1}b + \underbrace{(c_{\mathcal{N}}^\top - c_{\mathcal{B}}^\top B^{-1}N)}_{\bar{c}_{\mathcal{N}}^\top} x_{\mathcal{N}}.$$

An optimal basis requires $B^{-1}b \geq 0$ and $\bar{c}_{\mathcal{N}} \geq 0$. For maximization, reverse only the reduced-cost sign.

A general one-at-a-time RHS interval

Change b_i by δ. Define $g = B^{-1}e_i$, where e_i has 1 in position i and zeros elsewhere.

$$x_{\mathcal{B}}(\delta) = x_{\mathcal{B}} + \delta g.$$

$$\max_{k:g_k>0} \frac{-x_{\mathcal{B},k}}{g_k} \le \delta \le \min_{k:g_k<0} \frac{-x_{\mathcal{B},k}}{g_k}.$$

An empty lower set gives $-\infty$; an empty upper set gives $+\infty$. Rows with $g_k = 0$ impose no new restriction.

The reduced costs do not change. With $p^\top = c_{\mathcal{B}}^\top B^{-1}$, the optimal value inside this interval is $F(b) + \delta p_i$.

Why the 100% rule guarantees feasibility

Let d_i be the positive finite allowable magnitude in the chosen direction, and v_i the signed change of basic values per unit. Then $x_{\mathcal{B}} + d_i v_i \geq 0$.

Set $\alpha_i = |\delta_i|/d_i$. If $R = \sum_i \alpha_i \leq 1$, then

$$x_{\mathcal{B}} + \sum_i |\delta_i| v_i = \left(1 - \sum_i \alpha_i\right) x_{\mathcal{B}} + \sum_i \alpha_i (x_{\mathcal{B}} + d_i v_i) \geq 0.$$

> This is a nonnegative combination of feasible basic-value vectors. Directions with infinite allowance have nonnegative components and contribute zero to R. A nonzero change with zero allowance is not certified.

The rule is sufficient, not necessary: changes can offset one another even when the percentage sum exceeds 100%.

Worked bakery example above 100%

General cost changes: include the changing basic costs

For minimization, change c to $c + \Delta c$. The quantities stay $B^{-1}b$, while

$$\bar{c}_{\mathcal{N}}^{\text{new}} = \bar{c}_{\mathcal{N}} + \Delta c_{\mathcal{N}} - N^{\top} B^{-\top} \Delta c_{\mathcal{B}}.$$

> Nonbasic cost c_j changes alone: require $\bar{c}_j + \delta \geq 0$. Basic cost in basis position k changes: require $\bar{c}_j - \delta (B^{-1} A_j)_k \geq 0$ for every nonbasic j.

The new value, while this basis remains optimal, is $(c_{\mathcal{B}} + \Delta c_{\mathcal{B}})^{\top} x_{\mathcal{B}}$.

Changing a nonbasic column: retain its old reduced cost

For minimization, replace nonbasic column A_j by $A_j + \delta e_i$. The basis and basic values are unchanged.

$$\begin{aligned} \bar{c}_j^{\text{new}} &= c_j - p^\top (A_j + \delta e_i) \\ &= \bar{c}_j - \delta p_i. \end{aligned}$$

Require $\bar{c}_j - \delta p_i \geq 0$. If $p_i > 0$, this is $\delta \leq \bar{c}_j / p_i$. If $p_i < 0$, reverse the inequality. If $p_i = 0$, this entry change has no effect on that reduced cost.

All other reduced costs stay unchanged.

A basic-column change is not generally linear in size

In the bakery, replace the hours constraint by $(1+\varepsilon)x + y \le 9$, keeping the other constraints and profits.

Keeping hours and sugar active gives $(1+\varepsilon)x + y = 9$ and $x + 2y = 14$, hence

$$x = \frac{4}{1+2\varepsilon}, \quad y = 7 - \frac{2}{1+2\varepsilon}, \quad s_2 = 9 - \frac{6}{1+2\varepsilon}.$$

$$z(\varepsilon) = 21 + \frac{2}{1+2\varepsilon}, \qquad z'(0) = -4 = -p_1 x^*.$$

For $\varepsilon \ge -1/6$, this basis is feasible and dual feasible. Below that limit, recheck or reoptimize.

Why the coefficient-change derivative is $-p_i x_j$

For a basic column in position k, let $B(\delta) = B + \delta e_i e_k^\top$. Suppose this basis remains invertible, feasible and dual feasible for small δ. Let $v(\delta) = c_{\mathcal{B}}^\top x_{\mathcal{B}}(\delta)$ be the optimal value.

$$B(\delta)x_{\mathcal{B}}(\delta) = b \quad \Rightarrow \quad e_i x_j^* + B x'_{\mathcal{B}}(0) = 0.$$

$$v'(0) = c_{\mathcal{B}}^\top x'_{\mathcal{B}}(0) = -c_{\mathcal{B}}^\top B^{-1} e_i x_j^* = -p_i x_j^*.$$

Strict basic values and strict nonbasic reduced-cost signs are sufficient to keep the basis optimal for sufficiently small changes. At degeneracy or a breakpoint, a unique derivative is not automatic.

RHS convexity: combine two feasible optimal plans

Fix A, c, and assume a finite optimum at feasible RHS vectors b^1, b^2. Let their optimal plans be x^1, x^2. For $0 \le \lambda \le 1$, set $\hat{x} = \lambda x^1 + (1-\lambda)x^2$.

$$A\hat{x} = \lambda b^1 + (1-\lambda)b^2, \qquad \hat{x} \ge 0.$$

$$\begin{aligned} F(\lambda b^1 + (1-\lambda)b^2) &\le c^\top \hat{x} \\ &= \lambda F(b^1) + (1-\lambda)F(b^2). \end{aligned}$$

This is exactly the convexity inequality.

Why there are finitely many linear pieces

Assume A has full row rank and the dual set $D = \{p : A^\top p \le c\}$ is nonempty. It has no lines and has finitely many vertices $p^1, \dots, p^K$.

For each feasible RHS with finite optimum, a dual vertex attains the maximum. Strong duality gives

$$F(b) = \max_{k=1,\dots,K} (p^k)^\top b.$$

> Each vertex supplies a linear function. Their maximum is convex and piecewise linear. Several may agree at a breakpoint.

Dependent equality rows can be removed first. Do not infer a unique dual vector merely because a plotted one-dimensional segment is straight.

An optimal dual vector is a subgradient

Extend $F(b)$ to $+\infty$ outside $S = \{Ax : x \geq 0\}$, and assume $D \neq \varnothing$. Let $p^* \in D$ attain the dual optimum at b^*.

$$\begin{aligned} F(b) &\geq (p^*)^\top b \\ &= F(b^*) + (p^*)^\top (b - b^*). \end{aligned}$$

> This follows directly from weak duality and equality of primal and dual values at the reference RHS.

A **subgradient** is any vector giving this supporting inequality for every b. The next page proves the converse.

Every subgradient is an optimal dual vector

Suppose q satisfies the supporting inequality at b^*. Because $F(0) = 0$ and $F(2b^*) = 2F(b^*)$, substitution of these two RHS vectors gives

$$q^\top b^* \geq F(b^*), \qquad q^\top b^* \leq F(b^*).$$

Thus $q^\top b^* = F(b^*)$. For each column A_j, the plan $x = e_j$ is feasible and costs c_j. Therefore

$$c_j \geq F(A_j) \geq F(b^*) + q^\top (A_j - b^*) = q^\top A_j.$$

So $A^\top q \leq c$ and $q^\top b^* = F(b^*)$: q is dual optimal.

Cost concavity: minimize over a fixed feasible set

Fix nonempty $P = \{x : Ax = b, x \geq 0\}$. On costs with finite optimum, let $G(c) = \min_{x \in P} c^\top x$.

$$\begin{aligned} G(\lambda c^1 + (1-\lambda)c^2) &= \min_{x \in P} \{\lambda (c^1)^\top x + (1-\lambda)(c^2)^\top x\} \\ &\geq \lambda G(c^1) + (1-\lambda) G(c^2). \end{aligned}$$

A basic feasible optimum exists, so $G(c) = \min_k c^\top x^k$ over the finite set of basic feasible solutions. It is concave and piecewise linear.

> An optimal x^* at c^* gives an upper supporting bound: $G(c) \leq G(c^*) + (x^*)^\top (c - c^*)$. Such a vector is a **supergradient**.

A cost breakpoint: bread profit rises past 3

For maximizing $qx + 3y$, the original basis has

$$z = 4q + 15 + (3 - 2q)s_1 + (q - 3)s_3.$$

At $q = 3$, the coefficient of s_3 is zero. Just above 3, let s_3 enter. The ratio test gives s_2 leaving at step 3.

$$\begin{aligned} s_3 &= 3 + 3s_1 - s_2, \\ x &= 7 + s_1 - s_2, \qquad y = 2 - 2s_1 + s_2, \\ z &= 7q + 6 + (q - 6)s_1 + (3 - q)s_2. \end{aligned}$$

The new basis is optimal for $3 \le q \le 6$. At $q = 6$, the next breakpoint is reached.

Verify all three RHS-value pieces

Use hours $T \geq 0$, flour 16, sugar 14, and maximize $2x + 3y$.

Hours	Optimal plan	Matching dual variables
$0 \leq T \leq 7$	$(0, T)$, profit $3T$	$(3, 0, 0)$
$7 \leq T \leq 10$	$(2T - 14, 14 - T)$, profit $T + 14$	$(1, 0, 1)$
$T \geq 10$	$(6, 4)$, profit 24	$(0, 1/3, 4/3)$

Every listed plan is feasible on its interval and attains its dual upper bound. Negate profits and dual variables to obtain the minimization graph.

Second example: minimization in standard form

$$\begin{aligned} \min \quad & 2x_1 + 5x_2 + 8x_3 + 6x_4 \\ \text{s.t.} \quad & x_1 + x_2 + 2x_3 = 5, \\ & x_2 + x_3 + x_4 = 3, \qquad x \geq 0. \end{aligned}$$

$$\begin{aligned} x_1 &= 2 - x_3 + x_4, \\ x_2 &= 3 - x_3 - x_4, \\ f &= 19 + x_3 + 3x_4. \end{aligned}$$

Basic variables x_1, x_2; optimum $(2, 3, 0, 0)$, cost 19. The nonnegative objective coefficients have the minimization sign.

Second example: change the second RHS

Replace 3 by $3+\delta$, with $x_3 = x_4 = 0$.

$$x_2 = 3 + \delta, \qquad x_1 = 2 - \delta.$$

$$-3 \le \delta \le 2, \qquad f^* = 2(2-\delta) + 5(3+\delta) = 19 + 3\delta.$$

Dual variables $p = (2, 3)$ are unchanged. At $\delta = -3$, $x_2 = 0$; at $\delta = 2$, $x_1 = 0$.

For simultaneous changes Δb, the basic values are $(2 + \Delta b_1 - \Delta b_2,\ 3 + \Delta b_2)$; test both together.

Second example: nonbasic and basic costs

Original objective dictionary: $f = 19 + x_3 + 3x_4$.

Change nonbasic cost $c_3 = 8 + \theta$: $f = 19 + (1 + \theta)x_3 + 3x_4$. Require $\theta \geq -1$, or $c_3 \geq 7$.

Change basic cost $c_1 = 2 + \theta$: substitute $\theta x_1 = \theta(2 - x_3 + x_4)$.

$$f = 19 + 2\theta + (1 - \theta)x_3 + (3 + \theta)x_4.$$

Require $-3 \leq \theta \leq 1$. Quantities stay $(2, 3, 0, 0)$; the dual vector becomes $(2 + \theta, 3 - \theta)$.

Second example: verify the matrix and dual formulas

$$B = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad B^{-1} = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}, \quad B^{-1}b = \begin{pmatrix} 2 \\ 3 \end{pmatrix}.$$

$$p^{\top} = (2,5)B^{-1} = (2,3).$$

$$\bar{c}_3 = 8 - (2,3)\begin{pmatrix} 2 \\ 1 \end{pmatrix} = 1, \qquad \bar{c}_4 = 6 - (2,3)\begin{pmatrix} 0 \\ 1 \end{pmatrix} = 3.$$

The same numbers appear in the objective dictionary. Matrix sensitivity and dictionary substitution express the same operations.

Degeneracy can make reported ranges basis-dependent

Consider $\max x$ subject to $x \le 1$, $x \le 1$, $x \ge 0$. The optimum is $x = 1$. Both constraints are active.

$$\min \; p_1 + p_2 \quad \text{s.t. } p_1 + p_2 \ge 1, \quad p_1, p_2 \ge 0.$$

Every $(p_1, p_2) = (t, 1 - t)$, $0 \le t \le 1$, is dual optimal. There is no unique price for one duplicated constraint.

> Increase only the first RHS: the second row still limits x to 1. Decrease the first RHS: it immediately becomes the limiting row. The value has a corner.

A reported zero allowable change or an active row with zero dual variable is therefore mathematically meaningful.

Above 100%: check the dictionary directly

Bakery: maximize $z = 2x + 3y$, with $x + y \leq 9$, $2x + y \leq 16$, $x + 2y \leq 14$, $x, y \geq 0$.

Return to the original basis (x, y, s_2). Now increase **hours by 1** and **sugar by 3**; keep flour at 16. Their one-at-a-time allowable increases are 1 and 4:

Sum of fractions: $R = \frac{1}{1} + \frac{3}{4} = \frac{7}{4} > 1$.

The changed limits are $(10, 16, 17)$. Substitution gives the full dictionary:

$$
\begin{aligned}
x &= 3 - 2s_1 + s_3, \\
y &= 7 + s_1 - s_3, \\
s_2 &= 3 + 3s_1 - s_3, \\
z &= 27 - s_1 - s_3.
\end{aligned}
$$

At $s_1 = s_3 = 0$, $(x, y, s_2) = (3, 7, 3) \geq 0$. The objective coefficients remain nonpositive: **the original basis is still optimal**, despite $R > 1$.

Return to the 100% rule