

## CORRECTION


## Correction: Palmitoylated APP Forms Dimers, Cleaved by BACE1

Raja Bhattacharyya, Rebecca H. Fenn, Cory Barren, Rudolph E. Tanzi, Dora M. Kovacs

In Fig 1, there were two instances of Fig 1B, whereas the figure legend referred to the presence of a Fig 1C. In Fig 2, the authors provide the correct HA-APP<sub>Y</sub> panel in Fig 2B. The published panel resulted from an inadvertent duplication of another gel, part of which correctly appears in lanes 2–4 of the HA-APP<sub>Y</sub> panel in Fig 3A. In Fig 6, the authors provide the correct APP-mGFP and CTF-mGFP panels in Fig 6B. The published panel resulted from an inadvertent duplication of the middle panels in Fig 6A. An error in labeling the top and middle panels of Fig 6A has been corrected. The original underlying quantitative data for this article were not included with the published article, but are now presented with this notice in S2 File.

With this correction of this article [1], the authors provide updated versions of Fig 1, Fig 2, and Fig 6; original uncropped underlying blots for the figures of concern (S1 File); and the original underlying quantitative data for this article (S2 File).

The authors apologize for the errors in the published article.




## OPEN ACCESS

**Citation:** Bhattacharyya R, Fenn RH, Barren C, Tanzi RE, Kovacs DM (2024) Correction: Palmitoylated APP Forms Dimers, Cleaved by BACE1. PLoS ONE 19(2): e0299972. <https://doi.org/10.1371/journal.pone.0299972>


**Published:** February 29, 2024

**Copyright:** © 2024 Bhattacharyya et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




**Fig 1. *pal*APP dimerizes ~4.5 times more efficiently compared *tot*APP and in *cis*-orientation.** A. Cells expressing APP<sub>V5</sub> or APP<sub>V5</sub> plus HA-APP<sub>Y</sub> were subjected to co-immunoprecipitation assays to detect APP<sub>V5</sub>/HA-APP<sub>Y</sub> interaction or APP-dimerization. APP<sub>V5</sub> was immunoprecipitated with an anti-V5 antibody. Immunoprecipitates were probed with an anti-HA antibody to detect pull-down of HA-APP<sub>Y</sub>. Subsequently the immunoprecipitates were subjected to mABE assay to detect *pal*APP<sub>V5</sub>/HA-APP<sub>Y</sub> interaction (or *pal*APP-dimerization). *Pal*APP<sub>V5</sub> pulled down both *pal*APP<sub>V5</sub> ( $M_{wt} \sim 102$  kD) and *pal*HA-APP<sub>Y</sub> ( $M_{wt} \sim 150$  kD) from cells expressing APP<sub>V5</sub> plus HA-APP<sub>Y</sub> but not from cells expressing only APP<sub>V5</sub>. B. *Tot*APP-dimers (APP<sub>V5</sub>/HA-APP<sub>Y</sub>) only form in cells expressing both APP<sub>V5</sub> and HA-APP<sub>Y</sub>. C. Quantitation of *pal*APP-dimers (*pal*APP<sub>V5</sub>/*pal*HA-APP<sub>Y</sub>) versus *tot*APP-dimers (APP<sub>V5</sub>/HA-APP<sub>Y</sub>). Error bars show the s.e.m. (\*\* $p < 0.01$ ). D. *pal*APP dimerizes in *cis*-orientation. Cells expressing HA-APP<sub>Y</sub> and cells expressing mycAPP were co-cultured in absence or presence of 1mM cell-impermeable cross-linker DTSSP. Cell extracts were subjected to a pull-down assay, using an anti-HA antibody to immunoprecipitate HA-APP<sub>Y</sub>. To test for APP-dimerization, the precipitates were probed with an anti-myc antibody (*panel b*, *co-culture*). Cells co-expressing HA-APP<sub>Y</sub> and mycAPP were also subjected to a co-IP assay using the anti-HA antibody to pull-down mycAPP with HA-APP<sub>Y</sub> (*panel b*, *co-expression*). To detect *pal*APP-dimerization, the immunoprecipitates were also subjected to mABE assay to detect co-IP of *pal*HA-APP<sub>Y</sub> with *pal*-mycAPP (*panel a*). The experiment is a representative of three independent experiments.

<https://doi.org/10.1371/journal.pone.0299972.g001>



**Fig 2. Palmitoylation-prone APP mutants exhibit increased APP dimerization compared to wtAPP.** A. Schematic representation of the Cys to Ser mutants of APP used for the following co-immunoprecipitation assays. B. Co-immunoprecipitation assay in cells co-expressing APP<sub>V5</sub> and HA-APP<sub>Y</sub> and its mutants containing indicated Cys to

<https://doi.org/10.1371/journal.pone.0299972.g002>



5 / 6

## Supporting information

**S1 File. Raw Western blot data for Figs 2, 3, and 6.**  
(PPTX)

**S2 File. Quantitative data for Figs 1, 3, 4, and 5.**  
(PPTX)

## Reference

1. Bhattacharyya R, Fenn RH, Barren C, Tanzi RE, Kovacs DM (2016) Palmitoylated APP Forms Dimers, Cleaved by BACE1. PLoS ONE 11(11): e0166400. <https://doi.org/10.1371/journal.pone.0166400> PMID: [27875558](https://pubmed.ncbi.nlm.nih.gov/27875558/)