

# Transformations between Vectors

- $T: \mathbf{x} \in \mathbb{R}^n \mapsto \mathbf{y} \in \mathbb{R}^m$ : Mapping  $n$ -dim vector to  $m$ -dim vector
- Example:

$$T: \mathbf{x} \in \mathbb{R}^3 \mapsto \mathbf{y} \in \mathbb{R}^2 \quad \mathbf{x} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \in \mathbb{R}^3 \quad \mapsto \quad \mathbf{y} = T(\mathbf{x}) = \begin{bmatrix} 4 \\ 5 \end{bmatrix} \in \mathbb{R}^2$$

# Matrix of Linear Transformation

- **Example:** Suppose  $T$  is a **linear** transformation from  $\mathbb{R}^2$  to  $\mathbb{R}^3$  such that  $T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix}$  and  $T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 0 \\ 1 \\ 2 \end{bmatrix}$ . With no additional information, find a formula for the image of an arbitrary  $\mathbf{x}$  in  $\mathbb{R}^2$

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = x_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$$\Rightarrow T(\mathbf{x}) = T\left(x_1 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = x_1 T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) + x_2 T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right)$$

$$= x_1 \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix} + x_2 \begin{bmatrix} 0 \\ 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ -1 & 1 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

# Matrix of Linear Transformation

- In general, let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a **linear** transformation. Then  $T$  is always written as a matrix-vector multiplication, i.e.,

$$T(\mathbf{x}) = A\mathbf{x} \text{ for all } \mathbf{x} \in \mathbb{R}^n$$

- In fact, the  $j$ -th column of  $A \in \mathbb{R}^{m \times n}$  is equal to the vector  $T(\mathbf{e}_j)$ , where  $\mathbf{e}_j$  is the  $j$ -th column of the identity matrix in  $\mathbb{R}^{n \times n}$ :

$$A = [T(\mathbf{e}_1) \quad \cdots \quad T(\mathbf{e}_n)]$$

- Here, the matrix  $A$  is called the **standard matrix** of the linear transformation  $T$

# Matrix of Linear Transformation

- **Example:** Find the standard matrix  $A$  of a linear transformation from  $\mathbb{R}^3$  to  $\mathbb{R}^2$  such that

$$T\left(\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 1 \\ 2 \end{bmatrix}, T\left(\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}\right) = \begin{bmatrix} 4 \\ 3 \end{bmatrix}, \text{ and } T\left(\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}\right) = \begin{bmatrix} 5 \\ 6 \end{bmatrix}.$$

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = x_1 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + x_3 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

$$\begin{aligned} \Rightarrow T(\mathbf{x}) &= T\left(x_1 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + x_2 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + x_3 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}\right) = x_1 T\left(\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}\right) + x_2 T\left(\begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}\right) + x_3 T\left(\begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}\right) \\ &= x_1 \begin{bmatrix} 1 \\ 2 \end{bmatrix} + x_2 \begin{bmatrix} 4 \\ 3 \end{bmatrix} + x_3 \begin{bmatrix} 5 \\ 6 \end{bmatrix} = \begin{bmatrix} 1 & 4 & 5 \\ 2 & 3 & 6 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = A\mathbf{x} \end{aligned}$$