

Schaum's Outline of Abstract Algebra Notes and Exercises

Jasper Ty

Contents

I	Sets and Relations	5
1	Sets	7
	Exercises	7
2	Relations and Operations	9
II	Number Systems	11
3	The Natural Numbers	13
4	The Integers	15
5	Some Properties of Integers	17
6	The Rational Numbers	19
7	The Real Numbers	21
8	The Complex Numbers	23
III	Groups, Rings, and Fields	25
9	Groups	27
	9.1 Groups	27
	9.2 Simple Properties of Groups	27
10	Further Topics on Group Theory	29
11	Rings	31
12	Integral Domains, Division Rings, and Fields	33
13	Polynomials	35
14	Vector Spaces	37

15 Matrices	39
16 Matrix Polynomials	41
17 Linear Algebras	43
18 Boolean Algebras	45

Part I

Sets and Relations

Chapter 1

Sets

I don't have much to write here as far as notes go.

Exercises

Q1.1. Exhibit in tabular form:

- (a) $A = \{a : a \in \mathbb{N}, 2 < a < 6\}$
- (b) $B = \{p : p \in \mathbb{N}, p < 10, p \text{ is odd}\}$
- (c) $C = \{x : x \in \mathbb{Z}, 2x^2 + x - 6 = 0\}$

A1.1.

- (a) $A = \{3, 4, 5\}$
- (b) $B = \{9, 7, 5, 3, 1\}$
- (c) $C = \{-2\}$

Q1.2. Let $A = \{a, b, c, d\}$, $B = \{a, c, g\}$, $C = \{c, g, m, n, p\}$. Then $A \cup B = \{a, b, c, d, g\}$, $A \cup C = \{a, b, c, d, g, m, n, p\}$, $B \cup C = \{a, c, g, m, n, p\}$;

A1.2. WHAT IS THE QUESTION?

Q1.3. Consider the subsets $K = \{2, 4, 6, 8\}$, $L = \{1, 2, 3, 4\}$, $M = \{3, 4, 5, 6, 8\}$ of $U = \{1, 2, 3, \dots, 10\}$.

- (a) Exhibit K' , L' , M' in tabular form.
- (b) Show that $(K \cup L)' = K' \cap L'$

A1.3.

$$\begin{aligned} \text{(a)} \quad K' &= \{1, 3, 5, 7, 9, 10\} \\ L' &= \{5, 6, 7, 8, 9, 10\} \\ U' &= \{1, 2, 7, 9, 10\} \end{aligned}$$

(b) $K \cup L = \{1, 2, 3, 4, 6, 8\}$, so $(K \cup L)' = \{5, 7, 9, 10\}$. Using the above, $K' \cap L' = \{5, 7, 9, 10\}$.

Q1.4.

Q1.5.

Q1.6.

Q1.7.

Q1.8. Prove $(A \cup B) \cup C = A \cup (B \cup C)$

A1.8. Any element x belongs to the left hand side if $(x \in A \vee x \in B) \vee x \in C$. It belongs to the right hand side if $x \in A \vee (x \in B \vee x \in C)$. Both expressions are logically equivalent.

Q1.9. Prove $(A \cap B) \cap C = A \cap (B \cap C)$

A1.9. Similar answer as the previous— follows from the associativity of \wedge

Q1.10. Prove $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$

A1.10. Follows from \wedge distributing over \vee

Q1.11. Prove $(A \cap B)' = A' \cap B'$

A1.11. Follows from DeMorgan's laws

Q1.12. Prove $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$

A1.12. Follows from \vee distributing over \wedge

Q1.13. Prove $A - (B \cup C) = (A - B) \cap (A - C)$

A1.13. Follows again from DeMorgan's laws

Chapter 2

Relations and Operations

Part II

Number Systems

Chapter 3

The Natural Numbers

Chapter 4

The Integers

Chapter 5

Some Properties of Integers

Chapter 6

The Rational Numbers

Chapter 7

The Real Numbers

Chapter 8

The Complex Numbers

Part III

Groups, Rings, and Fields

Chapter 9

Groups

9.1 Groups

DEFINITION 9.1: A non-empty set \mathcal{G} equipped with a binary operation \circ is a *group* if

P₁: $(a \circ b) \circ c = a \circ (b \circ c)$ (Associativity)

P₂: There exists an element $1 \in \mathcal{G}$ such that $1 \circ a = a \circ 1 = a$ for all $a \in \mathcal{G}$ (Unit)

P₃: For all $a \in \mathcal{G}$, there exists an element a^{-1} such that $a \circ a^{-1} = a^{-1} \circ a = 1$ (Inverse)

9.2 Simple Properties of Groups

THEOREM I. (Left Cancellation) Let $a, b, c \in \mathcal{G}$. Then $a \circ b = a \circ c$ implies $b = c$.

THEOREM II. (Latin Square Property)

Chapter 10

Further Topics on Group Theory

Chapter 11

Rings

Chapter 12

Integral Domains, Division Rings, and Fields

Chapter 13

Polynomials

Chapter 14

Vector Spaces

Chapter 15

Matrices

Chapter 16

Matrix Polynomials

Chapter 17

Linear Algebras

Chapter 18

Boolean Algebras