

Applied Estimation Lab 1

John Folkesson

Preparation Questions (Chap 1-3,7 in Thrun)



Question 1

- What is the difference between a 'control' \mathbf{u}_t , a 'measurement' \mathbf{z}_t and the state \mathbf{x}_t ? Give examples of each?

Question 1

- What is the difference between a 'control' \mathbf{u}_t , a 'measurement' \mathbf{z}_t and the state \mathbf{x}_t ? Give examples of each?
- The control is part of the dynamic model between \mathbf{x}_{t-1} and \mathbf{x}_t .
- \mathbf{x}_t is modeled as depending on both \mathbf{u}_t and \mathbf{x}_{t-1} by $\mathbf{g}(\mathbf{u}_t, \mathbf{x}_{t-1})$.
- Measurement is modeled as depending on the state at time t , \mathbf{x}_t .
- \mathbf{z}_t gives undirect information on the state at time t , \mathbf{x}_t thru the function $\mathbf{h}(\mathbf{x}_t)$.
- difference is that \mathbf{u}_t is used in predict while \mathbf{z}_t is used in update.

Question 2

- Can the uncertainty in the belief increase during an update?
Why (or not)?

Question 2

- Can the uncertainty in the belief increase during an update?
Why (or not)?
- No
- $K_t = \bar{\Sigma}_t C_t^T (C_t \bar{\Sigma}_t C_t^T + Q_t)^{-1}$
- $\Sigma_t = \bar{\Sigma}_t - K_t C_t \bar{\Sigma}_t$
- change is $-\bar{\Sigma}_t C_t^T (C_t \bar{\Sigma}_t C_t^T + Q_t)^{-1} C_t \bar{\Sigma}_t = -MM^T$
- and $\mathbf{x}^T MM^T \mathbf{x} = \mathbf{y}^T \mathbf{y} \geq 0$ for any \mathbf{x} .
- Alternatively: $\Sigma_t = (\bar{\Sigma}_t^{-1} + C_t^T Q_t^{-1} C_t)^{-1}$
- $C_t^T Q_t^{-1} C_t$ is positive semidefinite.

Question 3

- During update what is it that decides the weighing between measurements and belief?

Question 3

- During update what is it that decides the weighing between measurements and belief?
- $\mu_t = (I - W)\bar{\mu}_t + W\mu_z$
 $W = \Sigma_t C_t^T Q_t^{-1} C_t$
and $C_t \mu_z = (z_t - \bar{z}_t + C_t \bar{\mu}_t)$ shown in Lecture 5.
- So an answer is the relative size of Q and Σ , measurement error and uncertainty after the update is done.
- Just the measurement covariance, Q_t is an acceptable answer even if it is less precise.

Question 4

- What would be the result of using a too large a covariance (Q matrix) for the measurement model?

Question 4

- What would be the result of using a too large a covariance (Q matrix) for the measurement model?
- Estimates would be pessimistic (conservative) and convergence would be slower.

Question 5

- What would give the measurements an increased effect on the updated state estimate?

Question 5

- What would give the measurements an increased effect on the updated state estimate?
- Decreasing the measurement model uncertainty, Q , would increase the Kalman gain.

Question 6

- What happens to the belief uncertainty during prediction?
How can you show that?

Question 6

- What happens to the belief uncertainty during prediction?
How can you show that?
- It usually increases.
- Normally the dynamic model adds noise to the state.
- $\bar{\Sigma}_t = R_t + A_t \Sigma_{t-1} A_t^T$
- So if $A_t \geq I$ then this is adding a positive definite matrix to something bigger or equal to Σ .
- In general uncertainty could go down during predict but most real systems it increases. This is because A_t is often about I and R is significantly large.

Question 7

- How can we say that the Kalman filter is the optimal and minimum least square error estimator?

Question 7

- How can we say that the Kalman filter is the optimal and minimum least square error estimator?
- We know that the Kalman filter gives us the true posterior distribution for a linear Gaussian system. Thus the question is really asking is there any other mean μ_{better} for the state that has lower expected square error than the Gaussian mean μ .
- $\int_{-\infty}^{\infty} (\mathbf{x} - \mu_{better})^2 G(\mathbf{x}, \mu, \Sigma) d\mathbf{x}$.
- $\int_{-\infty}^{\infty} (\mathbf{x}^2 - 2\mathbf{x}\mu_{better} + \mu_{better}^2) G(\mathbf{x}, \mu, \Sigma) d\mathbf{x}$.
- $\Sigma + \mu^T \mu - \mu^T \mu_{better} - \mu_{better}^T \mu + \mu_{better}^T \mu_{better}$ minimize by differentiating.
- $0 = -2\mu + 2\mu_{better}$
- $\mu_{better} = \mu$

Question 8

- Is it (KF) a MLE estimator? MAP?

Question 8

- Is it (KF) a MLE estimator? MAP?
- MLE yes if we start with an update from no prior then the mean is the MLE of the Gaussian posterior (the top of the bell curve)
- MAP yes if we start from some Gaussian prior on the state.

Question 9

- How does the extended Kalman filter relate to the Kalman filter EKF?

Question 9

- How does the extended Kalman filter relate to the Kalman filter EKF?
- The EKF applies the KF to the linearized nonlinear system.
- $G_t \iff A_t$
- $\mathbf{g}(\mathbf{u}_t, \mu_{t-1}) \iff A_t \mu_{t-1} + B_t \mathbf{u}_t$
- $H_t \iff C_t$
- \mathbf{y}_t is defined non-linearly wrt. $\bar{\mu}_t$

Question 10

- Is the EKF guaranteed to converge to a consistent solution?

Question 10

- Is the EKF guaranteed to converge to a consistent solution?
- No. Consistency depends on the significance of the nonlinearity.

Question 11

- If our filter seems to diverge often can we change any parameter to try and reduce this?

Question 11

- If our filter seems to diverge often can we change any parameter to try and reduce this?
- Yes we might change our modeled uncertainties Q and R . Typically divergence occurs on update and increasing the relative size of the measurement covariance, Q , will help.
- If the divergence was due to a poor data association then we might try changing our matching threshold.

Question 12

- If a robot is completely unsure of its location and measures the range r to a known landmark with Gaussian noise what does its posterior belief of its location $p(x, y, \theta | r)$ look like?

Question 12

- If a robot is completely unsure of its location and measures the range r to a known landmark with Gaussian noise what does its posterior belief of its location $p(x, y, \theta | r)$ look like?
- It will have a uniform distribution over the heading θ between $-\pi$ and π . The position will be a sort of donut/ring. So a Gaussian on ρ in radial coordinate with uniform distribution on the angle ϕ .
- $x = \rho \cos \phi$
- $y = \rho \sin \phi$
- $\exp -(\rho - r)^2 / (2\sigma_r^2)$

Question 13

- If the above measurement also included a bearing how would the posterior look?

Question 13

- If the above measurement also included a bearing how would the posterior look?
- The same except that the heading and angle around the ring would be Gaussian with a completely correlated covariance.
- $\exp -[(\rho - r)^2/(2\sigma_r^2) + (b - \phi + \theta)^2/(2\sigma_b^2)]$

Question 14

- If the robot moves with relatively good motion estimation (prediction error is small) but a large initial uncertainty in heading θ how will the posterior look after traveling a long distance without seeing any features?

Question 14

- If the robot moves with relatively good motion estimation (prediction error is small) but a large initial uncertainty in heading θ how will the posterior look after traveling a long distance without seeing any features?
- It will look like a crescent/arc/C shape.
- The heading θ will be correlated with position along the arc.

Question 15

- If the above robot then sees a point feature and measures range and bearing to it how might the EKF update go wrong?

Question 15

- If the above robot then sees a point feature and measures range and bearing to it how might the EKF update go wrong?
- The linearization (Jacoboian) will produce an update 'direction' that is a straight line which can not move the estimate along the curved crescent.
- The Gaussian will not be able to represent the crescent shape and thus the update might diverge.