

# Natural Language Models and Interfaces

BSc Artificial Intelligence

Lecturer: Wilker Aziz

Institute for Logic, Language, and Computation

2018, week 3b — PGMs

# Quick intro to PGMs

Check the lecture notes on [PGMs](#)

# Tabular representation

Suppose  $A$ ,  $B$ , and  $C$  are binary rvs

1. How do we represent  $P_{A,B,C}$  (without making assumptions)?

# Tabular representation

Suppose  $A$ ,  $B$ , and  $C$  are binary rvs

1. How do we represent  $P_{A,B,C}$  (without making assumptions)?

Joint assignments			Probability values
$A$	$B$	$C$	$P_{A,B,C}$
0	0	0	$P_{A,B,C}(0,0,0)$
0	0	1	$P_{A,B,C}(0,0,1)$
0	1	0	$P_{A,B,C}(0,1,0)$
1	0	0	$P_{A,B,C}(1,0,0)$
0	1	1	$P_{A,B,C}(0,1,1)$
1	1	0	$P_{A,B,C}(1,1,0)$
1	0	1	$P_{A,B,C}(1,0,1)$
1	1	1	$P_{A,B,C}(1,1,1)$

Table : Tabular joint distribution over 3 binary rvs

# Tabular representation

Suppose  $A$ ,  $B$ , and  $C$  are binary rvs

1. How do we represent  $P_{A,B,C}$  (without making assumptions)?

Joint assignments			Probability values
$A$	$B$	$C$	$P_{A,B,C}$
0	0	0	$P_{A,B,C}(0,0,0)$
0	0	1	$P_{A,B,C}(0,0,1)$
0	1	0	$P_{A,B,C}(0,1,0)$
1	0	0	$P_{A,B,C}(1,0,0)$
0	1	1	$P_{A,B,C}(0,1,1)$
1	1	0	$P_{A,B,C}(1,1,0)$
1	0	1	$P_{A,B,C}(1,0,1)$
1	1	1	$P_{A,B,C}(1,1,1)$

Table : Tabular joint distribution over 3 binary rvs

2. How many probability values does it take in general for  $n$  variables with  $t$  outcomes each?

## Tabular representation

Suppose  $A$ ,  $B$ , and  $C$  are binary rvs

1. How do we represent  $P_{A,B,C}$  (without making assumptions)?

Joint assignments			Probability values
$A$	$B$	$C$	$P_{A,B,C}$
0	0	0	$P_{A,B,C}(0,0,0)$
0	0	1	$P_{A,B,C}(0,0,1)$
0	1	0	$P_{A,B,C}(0,1,0)$
1	0	0	$P_{A,B,C}(1,0,0)$
0	1	1	$P_{A,B,C}(0,1,1)$
1	1	0	$P_{A,B,C}(1,1,0)$
1	0	1	$P_{A,B,C}(1,0,1)$
1	1	1	$P_{A,B,C}(1,1,1)$

Table : Tabular joint distribution over 3 binary rvs

2. How many probability values does it take in general for  $n$  variables with  $t$  outcomes each?  $n^t$

# Directed graphical models or Bayesian networks

A directed acyclic graph (DAG)

- ▶ nodes represent rvs
- ▶ edges represent direct influence
- ▶ a set of conditional independence statements
  - ▶ an rv is conditionally independent of its **non-descendants** given its **parents**

## Conditional independence in BNs

Consider  $A$ ,  $B$ , and  $C$ , due to chain rule we can write

$$P_{A,B,C}(a, b, c) = P_A(a)P_{B|A}(b|a)P_{C|AB}(c|a, b) \quad (1)$$



## Conditional independence in BNs

Consider  $A$ ,  $B$ , and  $C$ , due to chain rule we can write

$$P_{A,B,C}(a, b, c) = P_A(a)P_{B|A}(b|a)P_{C|AB}(c|a, b) \quad (1)$$

But if we are given a particular set of assumptions




Figure : Examples of BN

then we can simplify it

$$P_{A,B,C}(a, b, c) = P_A(a)P_{B|A}(b|a)P_{C|AB}(c|a, b) \quad (2)$$

$$= P_A(a)P_{B|A}(b|a)P_{C|B}(c|b) \quad (3)$$

$C$  is independent of non-descendants  $\{A\}$  given its parents  $\{B\}$

# Chain rule for Bayesian networks

Chain rule (in general)

$$P_{X_1, \dots, X_n}(x_1, \dots, x_n) = \prod_{i=1}^n P_{X|X_{<i}}(x_i|x_{<i}) \quad (4)$$

Chain rule for Bayesian networks

$$P_{X_1, \dots, X_n}(x_1, \dots, x_n) = \prod_{i=1}^n P_{X|\text{Pa}_X}(x|\text{pa}_x) \quad (5)$$

where

- ▶  $\text{Pa}_X$  set of rvs parents of  $X$
- ▶  $\text{pa}_X$  assignments of parents of  $X$

# Representing BNs

Each variable (given its parents) gets a tabular CPD  
Thus for




Figure : Examples of BN

$A$	$P_A$	$A$	$B$	$P_{B A}$	$B$	$C$	$P_{C B}$
0	$P_A(0)$	0	0	$P_{B A}(0 0)$	0	0	$P_{C B}(0 0)$
1	$P_A(1)$	0	1	$P_{B A}(1 0)$	0	1	$P_{C B}(1 0)$
		1	0	$P_{B A}(0 1)$	1	0	$P_{C B}(0 1)$
		1	1	$P_{B A}(1 1)$	1	1	$P_{C B}(1 1)$

Representation cost

- ▶ from  $O(\prod_{i=1}^n |\text{supp}(X_i)|)$
- ▶ to  $O(\sum_{i=1}^n |\text{supp}(X_i)| \times |\text{supp}(\text{Pa}_{X_i})|)$

# Exercises




Figure : Write down the factorisation

Quiz

# Inferences

So the BN shows us what are the CPDs in the problem

- ▶ but what if we want to reason about something that's not a CPD?




Figure : Examples of BN

Here we have CPDs  $P_A$ ,  $P_{B|A}$ , and  $P_{C|B}$

- ▶ how do we reason about  $P_{B|C}$  or  $P_{A|B}$ ?
- ▶ or  $P_B$  or  $P_C$ ?
- ▶ or  $P_{BC|A}$ ?

# Inferences

So the BN shows us what are the CPDs in the problem

- ▶ but what if we want to reason about something that's not a CPD?




Figure : Examples of BN

Here we have CPDs  $P_A$ ,  $P_{B|A}$ , and  $P_{C|B}$

- ▶ how do we reason about  $P_{B|C}$  or  $P_{A|B}$ ?
- ▶ or  $P_B$  or  $P_C$ ?
- ▶ or  $P_{BC|A}$ ?

For whatever combination, we have rules of probability!

## Conditional probability and marginalisation

If we have CPDs  $P_A$ ,  $P_{B|A}$ , and  $P_{C|B}$ , infer  $P_{B|C}$

# Conditional probability and marginalisation

If we have CPDs  $P_A$ ,  $P_{B|A}$ , and  $P_{C|B}$ , **infer**  $P_{B|C}$

- ▶ start from the definition of conditional probability

$$P_{B|C}(b|c) = \frac{P_{BC}(b, c)}{P_C(c)}$$



# Conditional probability and marginalisation

If we have CPDs  $P_A$ ,  $P_{B|A}$ , and  $P_{C|B}$ , infer  $P_{B|C}$

- ▶ start from the definition of conditional probability

$$P_{B|C}(b|c) = \frac{P_{BC}(b, c)}{P_C(c)}$$

- ▶ marginalise  $A$  in the numerator

$$P_{B|C}(b|c) = \frac{\sum_a P_{ABC}(a, b, c)}{P_C(c)}$$

# Conditional probability and marginalisation

If we have CPDs  $P_A$ ,  $P_{B|A}$ , and  $P_{C|B}$ , **infer**  $P_{B|C}$

- ▶ start from the definition of conditional probability

$$P_{B|C}(b|c) = \frac{P_{BC}(b, c)}{P_C(c)}$$

- ▶ marginalise  $A$  in the numerator

$$P_{B|C}(b|c) = \frac{\sum_a P_{ABC}(a, b, c)}{P_C(c)}$$

- ▶ factorise the joint distribution to introduce the CPDs

$$P_{B|C}(b|c) = \frac{\sum_a P_A(a) P_{B|A}(b|a) P_{C|B}(c|b)}{P_C(c)}$$

# Conditional probability and marginalisation

If we have CPDs  $P_A$ ,  $P_{B|A}$ , and  $P_{C|B}$ , **infer**  $P_{B|C}$

- ▶ start from the definition of conditional probability

$$P_{B|C}(b|c) = \frac{P_{BC}(b, c)}{P_C(c)}$$

- ▶ marginalise  $A$  in the numerator

$$P_{B|C}(b|c) = \frac{\sum_a P_{ABC}(a, b, c)}{P_C(c)}$$

- ▶ factorise the joint distribution to introduce the CPDs

$$P_{B|C}(b|c) = \frac{\sum_a P_A(a) P_{B|A}(b|a) P_{C|B}(c|b)}{P_C(c)}$$

- ▶ rearrange the terms for convenience

$$P_{B|C}(b|c) = \frac{P_{C|B}(c|b) \sum_a P_A(a) P_{B|A}(b|a)}{P_C(c)}$$

# Conditional probability and marginalisation

If we have CPDs  $P_A$ ,  $P_{B|A}$ , and  $P_{C|B}$ , **infer**  $P_{B|C}$

- ▶ start from the definition of conditional probability

$$P_{B|C}(b|c) = \frac{P_{BC}(b, c)}{P_C(c)}$$

- ▶ marginalise  $A$  in the numerator

$$P_{B|C}(b|c) = \frac{\sum_a P_{ABC}(a, b, c)}{P_C(c)}$$

- ▶ factorise the joint distribution to introduce the CPDs

$$P_{B|C}(b|c) = \frac{\sum_a P_A(a) P_{B|A}(b|a) P_{C|B}(c|b)}{P_C(c)}$$

- ▶ rearrange the terms for convenience

$$P_{B|C}(b|c) = \frac{P_{C|B}(c|b) \sum_a P_A(a) P_{B|A}(b|a)}{P_C(c)}$$

- ▶ note that the last sum is the (inferred) marginal  $P_B(b)$

# Continuation

- ▶ we are here

$$P_{B|C}(b|c) = \frac{P_{C|B}(c|b) \sum_a P_A(a) P_{B|A}(b|a)}{P_C(c)}$$

# Continuation

- ▶ we are here

$$P_{B|C}(b|c) = \frac{P_{C|B}(c|b) \sum_a P_A(a) P_{B|A}(b|a)}{P_C(c)}$$

- ▶ now obtain the marginal in the denominator as a function of tabular CPDs

$$\begin{aligned} P_C(c) &= \sum_a \sum_b P_{ABC}(a, b, c) \\ &= \sum_a \sum_b P_A(a) P_{B|A}(b|a) P_{C|B}(c|b) \\ &= \sum_a P_A(a) \sum_b P_{B|A}(b|a) P_{C|B}(c|b) \end{aligned}$$

# Continuation

- ▶ we are here

$$P_{B|C}(b|c) = \frac{P_{C|B}(c|b) \sum_a P_A(a) P_{B|A}(b|a)}{P_C(c)}$$

- ▶ now obtain the marginal in the denominator as a function of tabular CPDs

$$\begin{aligned} P_C(c) &= \sum_a \sum_b P_{ABC}(a, b, c) \\ &= \sum_a \sum_b P_A(a) P_{B|A}(b|a) P_{C|B}(c|b) \\ &= \sum_a P_A(a) \sum_b P_{B|A}(b|a) P_{C|B}(c|b) \end{aligned}$$

- ▶ get back to the conditional

$$P_{B|C}(b|c) = \frac{P_{C|B}(c|b) \sum_a P_A(a) P_{B|A}(b|a)}{\sum_a P_A(a) \sum_b P_{B|A}(b|a) P_{C|B}(c|b)}$$

# References I