

Pizza Sales



Retrieve the total number of orders placed.

```
SELECT
```

```
    COUNT(order_id)as total_orders
```

```
FROM
```

```
orders;
```

	total_orders
	21350



Calculate the total revenue generated from pizza sales.

```
SELECT
    ROUND(SUM(order_details.quantity * pizzas.price),
          2) total_sales
FROM
    order_details
    JOIN
    pizzas ON order_details.pizza_id = pizzas.pizza_id;
```

	total_sales
	817860.05



Identify the highest-priced pizza.

```
SELECT
    pizza_types.name, pizzas.price
FROM
    pizza_types
    JOIN
    pizzas ON pizza_types.pizza_type_id = pizzas.pizza_type_id
ORDER BY pizzas.price DESC
LIMIT 1;
```

	name	price
▶	The Greek Pizza	35.95



Identify the most common pizza size ordered.

```
SELECT
    pizzas.size,
    COUNT(order_details.order_details_id) AS order_count
FROM
    pizzas
    JOIN
    order_details ON pizzas.pizza_id = order_details.pizza_id
GROUP BY pizzas.size
ORDER BY order_count DESC
LIMIT 1;
```

	size	order_count
▶	L	18526



List the top 5 most ordered pizza types along with their quantities.

```
SELECT
    pizza_types.name, SUM(order_details.quantity) AS quantity
FROM
    pizza_types
    JOIN
    pizzas ON pizza_types.pizza_type_id = pizzas.pizza_type_id
    JOIN
    order_details ON order_details.pizza_id = pizzas.pizza_id
GROUP BY pizza_types.name
ORDER BY quantity DESC
LIMIT 5;
```

	name	quantity
▶	The Classic Deluxe Pizza	2453
	The Barbecue Chicken Pizza	2432
	The Hawaiian Pizza	2422
	The Pepperoni Pizza	2418
	The Thai Chicken Pizza	2371

Join the necessary tables to find the total quantity of each pizza category ordered.

```
SELECT
    pizza_types.category,
    SUM(order_details.quantity) AS quantity
FROM
    pizza_types
    JOIN
    pizzas ON pizza_types.pizza_type_id = pizzas.pizza_type_id
    JOIN
    order_details ON order_details.pizza_id = pizzas.pizza_id
GROUP BY pizza_types.category
ORDER BY quantity DESC;
```

	category	quantity
►	Classic	14888
	Supreme	11987
	Veggie	11649
	Chicken	11050

Determine the distribution of orders by hour of the day.

```
SELECT
    HOUR(order_time) hour, COUNT(order_id) order_count
FROM
    orders
GROUP BY HOUR(order_time);
```

	hour	order_count
▶	11	1231
	12	2520
	13	2455
	14	1472
	15	1468
	16	1920
	17	2336
	18	2399
	19	2009
	20	1642
	21	1198
	22	663
	23	28
	10	8
	9	1

Join relevant tables to find the category-wise distribution of pizzas

```
SELECT
    category, COUNT(name) pizza_count
FROM
    pizza_types
GROUP BY pizza_types.category;
```

	category	pizza_count
▶	Chicken	6
	Classic	8
	Supreme	9
	Veggie	9

Determine the top 3 most ordered pizza types based on revenue.

```
SELECT
    pizza_types.name,
    Round(SUM(pizzas.price * order_details.quantity),2) AS revenue
FROM
    pizza_types
    JOIN
    pizzas ON pizza_types.pizza_type_id = pizzas.pizza_type_id
    JOIN
    order_details ON order_details.pizza_id = pizzas.pizza_id = order_details.pizza_id
GROUP BY pizza_types.name
ORDER BY revenue DESC
LIMIT 3;
```

	name	revenue
►	The Greek Pizza	5422207.2
	The Italian Vegetables Pizza	2487467.75
	The Chicken Pesto Pizza	2474391.75

Determine the top 3 most ordered pizza types based on revenue for each pizza category.

```
select category,revenue
from
(select category,name,revenue,
rank() over(partition by category order by revenue desc)as rn
from
(select pizza_types.category,pizza_types.name,
round(sum(pizzas.price*order_details.quantity),2)as revenue
from pizza_types join pizzas
on pizzas.pizza_type_id=pizza_types.pizza_type_id
join order_details
on order_details.pizza_id=pizzas.pizza_id
group by pizza_types.category,pizza_types.name) as a) as b
where rn<=3;
```

	category	revenue
▶	Chicken	43434.25
	Chicken	42768
	Chicken	41409.5
	Classic	38180.5
	Classic	32273.25
	Classic	30161.75
	Supreme	34831.25
	Supreme	33476.75
	Supreme	30940.5
	Veggie	32265.7
	Veggie	26780.75
	Veggie	26066.5

Thank You!

Questions? Let's connect!

