

WOUND

CALL 911 :

- Bleeding is severe
- You suspect [internal bleeding](#)
- There is an abdominal or chest wound
- Bleeding can't be stopped after 10 minutes of firm and steady pressure
- [Blood](#) spurts out of wound

FIRST AID:

1. Stop Bleeding

- Apply direct pressure on the cut or wound with a clean cloth, tissue, or piece of gauze until bleeding stops.
- If [blood](#) soaks through the material, don't remove it. Put more cloth or gauze on top of it and continue to apply pressure.
- If the wound is on the arm or leg, raise limb above the [heart](#), if possible, to help slow bleeding.
- [Wash your hands](#) again after giving [first aid](#) and before cleaning and dressing the wound.
- Do not apply a tourniquet unless the bleeding is severe and not stopped with direct pressure.

2. Clean Cut or Wound

- Gently clean with soap and warm water. Try to rinse soap out of wound to prevent irritation.
- Don't use [hydrogen peroxide](#) or iodine, which can damage tissue.

3. Protect the Wound

- Apply antibiotic cream to reduce risk of infection and cover with a sterile bandage.
- Change the bandage daily to keep the wound clean and dry.

4. When to Call a Doctor

- The wound is deep or the edges are jagged or gaping open.
- The wound is on the person's face.
- The wound has dirt or debris that won't come out.
- The wound shows signs of infection, such as redness, tenderness, or a thick discharge, or if the person runs a fever.
- The area around the wound feels numb.
- Red streaks form around the wound.
- The wound is a result of an animal or human bite.
- The person has a puncture wound or deep cut and hasn't had a [tetanus shot](#) in the past five years, or anyone who hasn't had a tetanus shot in the past 10 years.