



Test 2

Topic: Independent random variables

Period: 2016.1

Throughout this test $(\Omega, \mathcal{A}, \mathbf{P})$ is a probability space (Ω is a nonempty set) and $X, Y, U, V : (\Omega, \mathcal{A}) \rightarrow (\mathbb{R}, \mathcal{B}(\mathbb{R}))$ are random variables.

1. [5 pts.] Let X and Y be independent with $X \sim \exp_\alpha$ and $Y \sim \exp_\beta$ for certain $\alpha, \beta > 0$. Show that

$$\mathbf{P}[X < Y] = \frac{\alpha}{\alpha + \beta}.$$

2. [5 pts.] ([Box–Muller method](#)) Let U and V be i.i.d. with $U \sim \mathcal{N}_{[0,1]}$. Define

$$X := \sqrt{2 \log(U)} \cos(2\pi V) \quad \text{and} \quad Y := \sqrt{2 \log(U)} \sin(2\pi V).$$

Show that X and Y are independent with distribution $\mathcal{N}_{0,1}$.

3. [5 pts.] Let (X, Y) be uniformly distributed on the unit ball, i.e.,

$$f_{(X,Y)}(x, y) = \begin{cases} \frac{1}{\pi} & \text{if } x^2 + y^2 \leq 1 \\ 0 & \text{otherwise.} \end{cases}$$

Find the distribution of $R = \sqrt{X^2 + Y^2}$.

4. [5 pts.] Let X and Y be i.i.d. with $X \sim \mathcal{N}_{0,\sigma^2}$. Let

$$Z = \sqrt{X^2 + Y^2} \quad \text{and} \quad W = \arctan\left(\frac{X}{Y}\right).$$

Show that

- (a) Z has [Rayleigh distribution](#).
- (b) W is uniformly distributed on $] -\pi/2, \pi/2[$.
- (c) Z and W are independent.

April 30, 2016