

4 Moments

In the following $(\Omega, \mathcal{A}, \mathbf{P})$ will denote our canonical probability space and X, Y, T, X_1, X_2, \dots are real random variables.

Definition 4.1.

- (i) If $X \in \mathcal{L}^1(\mathbf{P})$, then X is called **integrable** and we call

$$\mathbf{E}[X] := \int X d\mathbf{P}$$

the **expectation** or **mean** of X . If $\mathbf{E}[X] = 0$, then X is called **centered**. More generally, we also write $\mathbf{E}[X] = \int X d\mathbf{P}$ if only X^- or X^+ is integrable.

- (ii) If $n \in \mathbb{N}$ and $X \in \mathcal{L}^n(\mathbf{P})$, then the quantities

$$m_k := \mathbf{E}[X^k], \text{ for any } k = 1, \dots, n,$$

are called the **k th moments** of X .

- (iii) If $X \in \mathcal{L}^2(\mathbf{P})$, then X is called **square integrable** and

$$\mathbf{Var}[X] := \mathbf{E}[X^2] - \mathbf{E}[X]^2$$

is the **variance** of X . The number

$$\sigma := \sqrt{\mathbf{Var}[X]}$$

is called the **standard deviation** of X .

- (iv) If $X, Y \in \mathcal{L}^2(\mathbf{P})$, then we define the **covariance** of X and Y by

$$\mathbf{Cov}[X, Y] := \mathbf{E}[(X - \mathbf{E}[X])(Y - \mathbf{E}[Y])].$$

X and Y are called **uncorrelated** if $\mathbf{Cov}[X, Y] = 0$ and **correlated** otherwise.

Remark 4.2.

- (i) The definition in (ii) is sensible since $\mathcal{L}^n(\mathbf{P}) \subset \mathcal{L}^k(\mathbf{P})$ for all $k = 1, \dots, n$.

- (ii) The standard deviation of X makes sense in definition (iii) since

$$\mathbf{Var}[X] = \mathbf{E}[(X - \mathbf{E}[X])^2] \geq 0.$$

- (iii) If $X, Y \in \mathcal{L}^2(\mathbf{P})$, then $XY \in \mathcal{L}^1(\mathbf{P})$ since $|XY| \leq X^2 + Y^2$. Hence the definition in (iv) makes sense and we have

$$\mathbf{Cov}[X, Y] = \mathbf{E}[XY] - \mathbf{E}[X]\mathbf{E}[Y].$$

In particular, $\mathbf{Var}[X] = \mathbf{Cov}[X, X]$.

Now, we collect the most important rules of expectations. All of these properties are direct consequences of the corresponding properties of the integral.

Property 4.3 (Rules for expectations). Suppose that X, Y, X_1, X_2, \dots are integrable.

(i) [Linearity] Let $a, b \in \mathbb{R}$. Then $aX + bY$ is integrable and

$$\mathbf{E}[aX + bY] = a\mathbf{E}[X] + b\mathbf{E}[Y].$$

(ii) If $X \geq 0$ a.s., then

$$\mathbf{E}[X] = 0 \quad \Leftrightarrow \quad X = 0 \text{ a.s.}$$

(iii) [Monotonicity] If $X \leq Y$ a.s., then $\mathbf{E}[X] \leq \mathbf{E}[Y]$, with equality iff $X = Y$ a.s.

(iv) [Triangle inequality] $|\mathbf{E}[X]| \leq \mathbf{E}[|X|]$.

(v) If $X_n \geq 0$ a.s. for all $n \in \mathbb{N}$, then

$$\mathbf{E} \left[\sum_{n=1}^{+\infty} X_n \right] = \sum_{n=1}^{+\infty} \mathbf{E}[X_n].$$

(vi) If $X_n \uparrow X$, then

$$\mathbf{E}[X] = \lim_{n \rightarrow +\infty} \mathbf{E}[X_n].$$

Proof.

(v) It follows from the monotone convergence theorem.

(vi) Again, it follows from applying the monotone convergence theorem to $X_n - X_1$. \square

Property 4.4. Let $h : (\mathbb{R}, \mathcal{B}(\mathbb{R})) \rightarrow (\mathbb{R}, \mathcal{B}(\mathbb{R}))$ be measurable. Then $h \circ X \in \mathcal{L}^1(\mathbf{P})$ iff $h \in \mathcal{L}^1(\mathbf{P}_X)$, and in this case:

$$\mathbf{E}[h \circ X] = \int h(x) \mathbf{P}_X(dx).$$

Proof. This follows from the image measure property of integrals. \square

Remark 4.5. If X and Y are indentially distributed, by virtue of the property above,

(i) $X, Y \in \mathcal{L}^1(\mathbf{P}) \Rightarrow \mathbf{E}[X] = \mathbf{E}[Y]$,

(ii) $X, Y \in \mathcal{L}^2(\mathbf{P}) \Rightarrow \mathbf{Var}[X] = \mathbf{Var}[Y]$.

Again probability theory comes into play when independence enters the stage; that is, when we exit the realm of linear integration theory.

Theorem 4.6. If X_1, \dots, X_n are independent and one of the following conditions holds:

$$(i) \ X_1, \dots, X_n \geq 0$$

$$(ii) \ X_1, \dots, X_n \in \mathcal{L}^1(\mathbf{P})$$

then

$$\mathbf{E}[X_1 \cdots X_n] = \mathbf{E}[X_1] \cdots \mathbf{E}[X_n].$$

Lemma 4.7. Let $h : \mathbb{R}^2 \rightarrow \mathbb{R}$ be measurable. If X and Y are independent and one of the following conditions holds:

$$(i) \ h \geq 0$$

$$(ii) \ h(X, Y) \in \mathcal{L}^1(\mathbf{P})$$

then

$$\mathbf{E}[h(X, Y)] = \int h \mathbf{P}_X \otimes \mathbf{P}_Y.$$

In particular, if $h(x, y) = f(x)g(y)$ where $f, g : \mathbb{R} \rightarrow \mathbb{R}$ are measurable functions and one of the following conditions holds:

$$(i) \ f, g \geq 0$$

$$(ii) \ f(X), g(Y) \in \mathcal{L}^1(\mathbf{P})$$

then

$$\mathbf{E}[f(X)g(Y)] = \mathbf{E}[f(X)]\mathbf{E}[g(Y)].$$

Proof. First, let us assume the validity of the first part and the hypothesis of the second part. If $f, g \geq 0$ then, by virtue of property 4.4,

$$\begin{aligned} \mathbf{E}[f(X)g(Y)] &= \mathbf{E}[h(X, Y)] \\ &= \int h \mathbf{P}_X \otimes \mathbf{P}_Y \\ &= \int \left(\int f(x)g(y) \mathbf{P}_X(dx) \right) \mathbf{P}_Y(dy) \\ &= \int f(x) \mathbf{P}_X(dx) \int g(y) \mathbf{P}_Y(dy) \\ &= \mathbf{E}[f(X)]\mathbf{E}[f(Y)]. \end{aligned}$$

Now, if $f(X), g(Y) \in \mathcal{L}^1(\mathbf{P})$ then, again by virtue of property 4.4, $f \in \mathcal{L}^1(\mathbf{P}_X)$ and $g \in \mathcal{L}^1(\mathbf{P}_Y)$, and thus

$$\begin{aligned} \int |h| \mathbf{P}_X \otimes \mathbf{P}_Y &= \int \left(\int |f(x)||g(y)| \mathbf{P}_X(dx) \right) \mathbf{P}_Y(dy) \\ &= \int |f(x)| \mathbf{P}_X(dx) \int |g(y)| \mathbf{P}_Y(dy) \\ &< +\infty. \end{aligned}$$

Hence, $h \in \mathcal{L}^1(\mathbf{P}_X \otimes \mathbf{P}_Y)$; then, using the fact that $\mathbf{P}_{X,Y} = \mathbf{P}_X \otimes \mathbf{P}_Y$ and the image measure property, $h(X, Y) \in \mathcal{L}^1(\mathbf{P})$, and the conclusion follows. Now, let us assume the hypothesis of the first part. If

□

Proof of the theorem 4.6.

□

In the following, an important identity that simplifies the calculation of the expected value of the sum of a random number of random quantities.

Theorem 4.8 (Wald's equation). Let $T, X_1, X_2, \dots : \Omega \rightarrow \mathbb{R}$ be independent random variables in $\mathcal{L}^1(\mathbf{P})$ with $\mathbf{P}(T \in \mathbb{N}_0) = 1$ and X_1, X_2, \dots identically distributed. Define:

$$S_T := \sum_{i=1}^T X_i.$$

Then $S_T \in \mathcal{L}^1(\mathbf{P})$ and

$$\mathbf{E}(S_T) = \mathbf{E}(T)\mathbf{E}(X_1).$$

Proof. Exercise.

□

Property 4.9 (Rules for variance and covariance). Let $X, Y, X_1, \dots, X_n : \Omega \rightarrow \mathbb{R}$ be square integrable random variables and $\alpha \in \mathbb{R}$, and let $E = \{Z : \Omega \rightarrow \mathbb{R} ; Z \in \mathcal{L}^2(\mathbf{P})\}$. Then:

- (i) $\mathbf{Var}[X] = 0 \iff X = \mathbf{E}[X] \text{ a.s.}$
- (ii) The map $f : \mathbb{R} \rightarrow \mathbb{R}, x \mapsto \mathbf{E}[(X - x)^2]$, is minimal at $\mathbf{E}[X]$ with $f(\mathbf{E}[X]) = \mathbf{Var}[X]$.
- (iii) $\mathbf{Var}[\alpha X] = \alpha^2 \mathbf{Var}[X]$.
- (iv) The map $\mathbf{Cov} : E \times E \rightarrow \mathbb{R}$ is a positive semidefinite symmetric bilinear form.
- (v) If $X_1 + \dots + X_n$ are uncorrelated, then

$$\mathbf{Var}[X_1 + \dots + X_n] = \mathbf{Var}[X_1] + \dots + \mathbf{Var}[X_n].$$

- (vi) [Cauchy-Schwarz inequality]

$$\mathbf{Cov}[X, Y]^2 \leq \mathbf{Var}[X]\mathbf{Var}[Y].$$

Equality holds iff there are $a, b, c \in \mathbb{R}$ with $|a| + |b| + |c| > 0$ and $aX + bY + c = 0$ a.s.

Proof.

(v)

(vi)

□

Example 4.10. Let $X : \Omega \rightarrow \mathbb{R}$ be a random variable.

- (i) Let $p \in [0, 1]$ and let $X \sim \text{Ber}_p$. Then $\mathbf{E}[X] = p$ and $\mathbf{Var}[X] = p(1 - p)$.
- (ii) Let $n \in \mathbb{N}$ and $p \in [0, 1]$, and let $X \sim \text{B}_{n,p}$. Then $\mathbf{E}[X] = np$ and $\mathbf{Var}[X] = np(1 - p)$.
- (iii) Let $\mu \in \mathbb{R}$ and $\sigma^2 > 0$, and let $X \sim \mathcal{N}_{\mu, \sigma^2}$. Then $\mathbf{E}[X] = \mu$ and $\mathbf{Var}[X] = \sigma^2$.
- (iv) Let $\theta > 0$ and let $X \sim \exp_\theta$. Then $\mathbf{E}[X] = \frac{1}{\theta}$ and $\mathbf{Var}[X] = \frac{1}{\theta^2}$.

In the following, an important identity that simplifies the calculation of the variance value of the sum of a random number of random quantities.

Theorem 4.11 ([Blackwell-Girshick equation](#)). Let $T, X_1, X_2, \dots : \Omega \rightarrow \mathbb{R}$ be independent random variables in $\mathcal{L}^2(\mathbf{P})$ with $\mathbf{P}[T \in \mathbb{N}_0] = 1$ and X_1, X_2, \dots identically distributed. Define:

$$S_T := \sum_{i=1}^T X_i.$$

Then $S_T \in \mathcal{L}^2(\mathbf{P})$ and

$$\mathbf{Var}[S_T] = \mathbf{Var}[T]\mathbf{E}[X_1]^2 + \mathbf{E}[T]\mathbf{Var}[X_1].$$

Proof. Exercise.

□

Exercise 4.1. Let $h : \mathbb{R} \rightarrow \mathbb{R}$ be measurable and let X and Y be identically distributed. Prove the following propositions.

- (i) $h(X)$ is integrable iff $h(Y)$ is integrable.
- (ii) In this case we have $\mathbf{E}[h(X)] = \mathbf{E}[h(Y)]$.

Exercise 4.2. Let X be integrable and with symmetric distribution. Prove the following propositions.

- (i) $\mathbf{E}[X] = 0$.
- (ii) If $h : \mathbb{R} \rightarrow \mathbb{R}$ is measurable and odd, then $h(X)$ has symmetric distribution.
- (iii) If $X \sim \mathcal{N}_{0,1}$, then $\mathbf{E}[X^k] = 0$ for every odd $k \geq 1$.

Exercise 4.3. Prove that if $X \in \mathcal{L}^1(\mathbf{P})$ and has density f , then

$$\mathbf{E}[X] = \int x f(x) \lambda(dx).$$

Exercise 4.4. Assume that (X, Y) are uniformly distributed on a circle with radius a , then

$$f_{(X,Y)}(x, y) = \begin{cases} \frac{1}{\pi a^2} & \text{if } x^2 + y^2 \leq a^2, \\ 0 & \text{elsewhere.} \end{cases}$$

Find $\mathbf{E}[X]$.

Exercise 4.5. Suppose that X and Y are independent with probability densities:

$$f_X(x) = \begin{cases} \frac{8}{x^3} & \text{if } x > 2, \\ 0 & \text{elsewhere,} \end{cases}$$

and

$$f_Y(y) = \begin{cases} \frac{2}{y} & \text{if } 0 < y < 1, \\ 0 & \text{elsewhere,} \end{cases}$$

Find $\mathbf{E}[XY]$.

Exercise 4.6. Let $X_1, X_2, \dots \geq 0$ be i.i.d. Prove the following propositions.

(i)

$$\limsup_{n \rightarrow +\infty} \frac{1}{n} X_n = \begin{cases} 0 \text{ a.s.} & \text{if } \mathbf{E}[X_1] < +\infty, \\ +\infty \text{ a.s.} & \text{if } \mathbf{E}[X_1] = +\infty. \end{cases}$$

(ii) For any $c \in]0, 1[$:

$$\sum_{n=1}^{+\infty} e^{X_n} c^n \begin{cases} < +\infty \text{ a.s.} & \text{if } \mathbf{E}[X_1] < +\infty, \\ = +\infty \text{ a.s.} & \text{if } \mathbf{E}[X_1] = +\infty. \end{cases}$$

Exercise 4.7. Let $\Omega =]0, 1[$, \mathcal{A} be the class of Borel sets and \mathbf{P} be the Lebesgue measure. If $X_n(\omega) = \sin(2\pi n\omega)$, $n = 1, 2, \dots$, then prove that X_1, X_2, \dots are uncorrelated but not independent.

Exercise 4.8. Prove that if $\mathbf{P}[X \in [0, 1]] = 1$, then $\mathbf{Var}[X] \leq 1/4$.

Exercise 4.9. By investing in a particular stock, a person can make a profit in one year of \$4,000 with probability 0.3 or take a loss of \$1,000 with probability 0.7.

(i) What is the person's expected gain?

(ii) What is the variance?

Exercise 4.10. Suppose that X represents the number of errors per 100 lines of software code and has the following probability distribution:

X	2	3	4	5	6
Probability	0.01	0.25	0.40	0.30	0.04

- (i) Find the variance of X
- (ii) Find the mean and variance of $3X - 2$.

Exercise 4.11. Let a six-sided die. Take the number on the die (call it T) and roll that number of six-sided dice to get the numbers X_1, \dots, X_T , and add up their values. What is the expected value of this sum?

Exercise 4.12. Let a particle in the x axis with probability $2/3$ to move one meter to the right and $1/3$ to move one meter to the left. Take a number on \mathbb{N}_0 and call it T ; suppose that $T \sim \text{Poi}_3$. Then starting at the origin, the particle performs T movements along the axis, say X_1, \dots, X_T . What is the expected final position of this particle?