

Lecture Notes · Arithmetic · Session 1 — Divisibility, Euclidean division and the Euclidean algorithm

All the session's material with complete proofs: division with remainder (existence via well-ordering, and uniqueness), the lemma that justifies the Euclidean algorithm, and Bézout's identity by its two routes — the structural one and the extended algorithm with a fully worked example. Exercises and a Going further section at the end.

1.1 Divisibility

Definition 1.1.1. For $a, b \in \mathbb{Z}$: a **divides** b , written $a \mid b$, if there exists $c \in \mathbb{Z}$ with $b = ac$.

Proposition 1.1.2. For arbitrary integers: 1. (*Transitivity*) $a \mid b$ and $b \mid c \Rightarrow a \mid c$. 2. (*Combinations*) if $a \mid b$ and $a \mid c$, then $a \mid (xb + yc)$ for all $x, y \in \mathbb{Z}$. 3. If $a \mid b$ and $b \neq 0$, then $|a| \leq |b|$.

Proof. (1) $b = au, c = bv \Rightarrow c = a(uv)$. (2) $b = au, c = av \Rightarrow xb + yc = a(xu + yv)$. (3) $b = ac$ with $c \neq 0$, so $|c| \geq 1$ and $|b| = |a||c| \geq |a|$. ■

Property (2), so modest, is the key to half the session: Bézout lives off it.

1.2 Euclidean division

Theorem 1.1.3 (division with remainder). For $a \in \mathbb{Z}$ and $b > 0$ there exist **unique** integers q (quotient) and r (remainder) with

$$a = bq + r, \quad 0 \leq r < b.$$

Proof. Existence. The set $S = \{a - bq : q \in \mathbb{Z}, a - bq \geq 0\}$ is nonempty (for q very negative, $a - bq$ becomes positive). By the **well-ordering principle** (M0: every nonempty subset of $\mathbb{N}$ has a least element), S has a least element $r = a - bq_0 \geq 0$. We claim $r < b$: if we had $r \geq b$, then $r - b = a - b(q_0 + 1) \geq 0$ would be in S and smaller than r , against minimality.

Uniqueness (the “subtract and bound” trick). If $a = bq + r = bq' + r'$ with $0 \leq r, r' < b$, subtracting: $b(q - q') = r' - r$. The right-hand side satisfies $|r' - r| < b$; the left-hand side is a multiple of b . The only multiple of b with absolute value less than b is 0: hence $r = r'$ and, since $b \neq 0$, $q = q'$. ■




Figure 1: Euclidean division on the number line: a falls between two consecutive multiples of b ; the quotient q counts them and the remainder r is what is left over ($0 \leq r < b$).

Remark 1.1.4 (the convention matters — and it *is* a convention). For $a = -7$, $b = 3$: $-7 = 3 \cdot (-3) + 2$ — the remainder is 2, not -1 . The convention $0 \leq r < b$ makes the expression unique and is the one congruences use (Session 3).

So why do we require $b > 0$? For convenience, not out of necessity. The general version holds for every $b \neq 0$: there exist **unique** q, r with $a = bq + r$ and $0 \leq r < |b|$. And it reduces to the positive case in one line: apply the theorem to $|b|$, which gives $a = |b|q' + r$; if $b < 0$ then $|b| = -b$, so $a = b(-q') + r$ and it suffices to take $q = -q'$. For instance, $7 = (-3) \cdot (-2) + 1$, with $0 \leq 1 < 3 = |-3|$. What we **do** have to exclude is $b = 0$: then $a = 0 \cdot q + r$ forces $r = a$, and the condition $0 \leq r < 0$ is impossible to satisfy — it is division by zero, and the **only** genuine exclusion in the theorem. We state it with $b > 0$ because in everything that follows the divisor is positive by nature (the modulus n of congruences, the remainders in Euclid’s algorithm, the gcd normalised to be ≥ 0), which spares the later statements a forest of absolute-value bars.

*(In $K[x]$ —Polynomials module— the analogous hypothesis is precisely “**nonzero divisor**”: $g \neq 0$ with $\deg r < \deg g$. There is no order there letting us ask for “positive”, only the degree; the dictionary is $|\cdot| \leftrightarrow \deg$. Stated with $b \neq 0$ and $|b|$, division in $\mathbb{Z}$ and division in $K[x]$ say literally the same thing: nonzero divisor, remainder smaller than the divisor in the measure proper to each world.)*

And uniqueness depends on the band, not on the sign of b . Fixing $0 \leq r < |b|$ is **one** choice among several, and all of them give existence and uniqueness: the remainder carrying the sign of the **dividend** (what C, C++, Java and Rust do, where $-7 \% 3$ is -1), or the **centred** remainder, $-|b|/2 < r \leq |b|/2$, convenient in cryptography and in lattice reduction. Ours is **Python**’s convention: $-7 \% 3$ returns 2. It is worth checking in the interpreter, because that disagreement between languages is a classic source of bugs.

Structural lens (a mention): $\mathbb{Z}$ is the model of a world “with division with remainder”. This single property generates everything that follows — and in the Polynomials module we will see that $K[x]$ shares it, with the **degree** in the role of the absolute value.

1.3 Greatest common divisor and the Euclidean algorithm

Definition 1.1.5. For a, b not both zero, $\gcd(a, b)$ is the largest of the common divisors. (It exists: the common divisors are bounded by 1.1.2.3 and the set is nonempty — it contains 1.)

On “the largest” and the sign. Divisors come in pairs $\pm d$, so the largest in the usual order of $\mathbb{Z}$ is the **positive** one: thus the $\gcd$ is an integer ≥ 0 , well defined and **unique** — “the largest” is literal. The characterization by divisibility (d such that every common divisor divides d ; exercise 8) determines it only **up to sign**, and the convention fixes $\gcd \geq 0$. (The case $a = b = 0$ is excluded on purpose: **every** integer divides 0, so there is no greatest common divisor.) *(In $K[x]$ — the Polynomials module — the $\gcd$ is unique up to a nonzero constant factor, and it is normalized by making it monic: the same “up to a unit” issue.)*

Lemma 1.1.6 (why the Euclidean algorithm works). If $a = bq + r$, then the **common divisors of (a, b) are exactly the common divisors of (b, r)** . In particular, $\gcd(a, b) = \gcd(b, r)$.

Proof. If $d \mid a$ and $d \mid b$: since $r = a - bq$ is a combination of a and b , $d \mid r$ (Prop. 1.1.2.2) — a common divisor of (b, r) . Conversely, if $d \mid b$ and $d \mid r$: $a = bq + r$ is a combination of b and r , so $d \mid a$. The two pairs have *the same set* of common divisors; in particular, the same maximum. ■

Algorithm 1.1.7 (Euclid). Divide a by b , replace $(a, b) \mapsto (b, r)$, repeat until the remainder is 0; the $\gcd$ is **the last nonzero remainder**.

Justification. Each step preserves the $\gcd$ (Lemma 1.1.6). The remainders form a strictly decreasing sequence of nonnegative integers ($0 \leq r < b$), which by well-ordering **cannot be infinite**: the algorithm terminates. On reaching $(d, 0)$, $\gcd(d, 0) = d$. ■

Example 1.1.8. $\gcd(252, 198)$:

$$252 = 1 \cdot 198 + 54 \quad 198 = 3 \cdot 54 + 36 \quad 54 = 1 \cdot 36 + 18 \quad 36 = 2 \cdot 18 + 0$$

$\gcd(252, 198) = 18$.

1.4 Bézout’s identity

Theorem 1.1.9 (Bézout). For a, b **not both zero**, there exist $x, y \in \mathbb{Z}$ such that $\gcd(a, b) = ax + by$.

Proof (structural). Let $D = \{ax + by : x, y \in \mathbb{Z}\}$ and let d be the **least positive element** of D (it exists: since a, b are not both zero, D contains some positive element — if $a \neq 0$, $|a| \in D$; if $a = 0$, then $b \neq 0$ and $|b| \in D$; well-ordering). We claim that $d = \gcd(a, b)$:

- **d divides a :** dividing, $a = dq + r$ with $0 \leq r < d$; then $r = a - dq$ is a

combination of a and b (because d is one), that is, $r \in D$ with $0 \leq r < d$. By minimality of d , $r = 0$. Likewise for b .

- **It is the largest:** any common divisor c of a and b divides every combination $ax + by$ (Prop. 1.1.2.2), in particular d ; since $d > 0$, this forces $|c| \leq d$, and therefore $c \leq d$. Thus d is the largest of the common divisors. ■

Method 1.1.10 (extended Euclidean algorithm). The coefficients x, y are **constructed** by working back up the divisions of Example 1.1.8, solving for each remainder:

$$\begin{aligned} 18 &= 54 - 36 = 54 - (198 - 3 \cdot 54) = 4 \cdot 54 - 198 \\ &= 4(252 - 198) - 198 = 4 \cdot 252 - 5 \cdot 198. \end{aligned}$$

Check: $4 \cdot 252 - 5 \cdot 198 = 1008 - 990 = 18$ ✓.

Remark 1.1.11 (the coefficients are not unique — and which ones are all of them). Method 1.1.10 hands you **one** pair (x, y) , and there are **infinitely many**. Fixing $d = \gcd(a, b)$ and a pair (x_0, y_0) with $ax_0 + by_0 = d$, the complete family is

$$x = x_0 + k \frac{b}{d}, \quad y = y_0 - k \frac{a}{d}, \quad k \in \mathbb{Z}.$$

They all work (a direct computation — the k terms cancel):

$$a(x_0 + k \frac{b}{d}) + b(y_0 - k \frac{a}{d}) = ax_0 + by_0 + k(\frac{ab}{d} - \frac{ab}{d}) = d.$$

And there are no others. If $ax + by = d$, subtracting from $ax_0 + by_0 = d$ leaves $a(x - x_0) = -b(y - y_0)$; dividing by d ,

$$\frac{a}{d}(x - x_0) = -\frac{b}{d}(y - y_0).$$

So $\frac{b}{d}$ divides $\frac{a}{d}(x - x_0)$, and **it is coprime to $\frac{a}{d}$** (exercise 7), whence $\frac{b}{d} \mid (x - x_0)$. (*This step is the generalised Euclid lemma, and it follows from Bézout itself: from $u \frac{b}{d} + v \frac{a}{d} = 1$, multiplying by $x - x_0$, both summands turn out to be multiples of $\frac{b}{d}$.)* Hence $x - x_0 = k \frac{b}{d}$ for some k , and substituting, $y - y_0 = -k \frac{a}{d}$. ■

In Example 1.1.8 ($a = 252$, $b = 198$, $d = 18$): $\frac{b}{d} = 11$ and $\frac{a}{d} = 14$, so the family is $(4 + 11k, -5 - 14k)$. With $k = 1$: $15 \cdot 252 - 19 \cdot 198 = 3780 - 3762 = 18$ ✓.

Non-uniqueness turns out to be harmless in every later use: in Session 3 **one** x with $ax \equiv 1 \pmod{n}$ is enough, and every member of the family gives **the same** inverse mod n — precisely because they differ by multiples of $\frac{b}{d}$.




Figure 2: The mental picture of the extended algorithm, with Example 1.1.8: **Euclid goes down** (each remainder becomes the divisor, until the last nonzero remainder — the gcd) and **Bézout climbs back up** (solve for the gcd in the next-to-last division and substitute the remainders upwards, until it is written as a combination of the original inputs).

What Bézout is for (an announcement that will be honored):
 in Session 2 it proves Euclid’s lemma (the heart of unique factorization);
 in Session 3 it **decides and computes inverses** modulo n ;
 in Session 4 it builds the solutions of the Chinese remainder theorem.
 It is, quite possibly, the most profitable result of the module.

Exercises

- ★ Compute $\gcd(1001, 273)$ by the Euclidean algorithm, recording each division, and work back up to write it as $1001x + 273y$. Check numerically.
- ★ Prove from the definition (choose and declare the **type of proof** — M0 connection): if $a \mid b$ and $b \mid a$, then $a = \pm b$.
- ★ Find the quotient and remainder when dividing -100 by 7 (mind the convention $0 \leq r < 7$!) and when dividing 100 by 7 . What is the relation between the two remainders?
- Justify from the definition: $\gcd(a, 0) = |a|$ (for $a \neq 0$) and $\gcd(a, 1) = 1$.
- Prove that $\gcd(a, b) = \gcd(a, b - a)$ **without** using Euclidean division (only Prop. 1.1.2.2), and explain why this already “almost” justifies the Euclidean algorithm.
- Give the uniqueness proof of Theorem 1.1.3 **by contradiction**, and compare it with the direct one in these notes (which do you find clearer? — a connection with M0).
- Prove: if $\gcd(a, b) = d$, then $\gcd(a/d, b/d) = 1$. (Hint: Bézout, divided by d .)
- The set $D = \{ax + by\}$ from the proof of Bézout: prove that D is exactly the set of **multiples** of d . (The combinations of two numbers are the

multiples of their gcd — the fine-grained version of Bézout.)

Going further and references

Going further (gcd of several numbers, and efficiency). $\gcd(a, b, c) = \gcd(\gcd(a, b), c)$, and the Euclidean algorithm is **genuinely fast**: the number of steps is proportional to the number of digits (the worst case is given by the Fibonacci numbers — Lamé’s theorem). That efficiency, set against the cost of factoring, is the crack on which the cryptography of Session 4 is built. See Childs, *A Concrete Introduction to Higher Algebra*, ch. 3–4.

Going further (Euclidean domains). Any “world” with a reasonable division with remainder (a *Euclidean domain*) repeats this entire session word for word: gcd, the Euclidean algorithm, Bézout. The star example of the course will be $K[x]$ (Polynomials module); others, such as the Gaussian integers $\mathbb{Z}[i]$, in Aluffi, *Algebra: Chapter 0*, ch. V.

References. Childs, ch. 2–4; Dorronsoro–Hernández, *Números, grupos y anillos*, ch. 2.

Lecture Notes · Arithmetic · Session 2 — Primes and unique factorization

The session’s complete proofs: the infinitude of the primes (with the preliminary lemma the classical proof usually takes for granted), Euclid’s lemma via Bézout, and the fundamental theorem of arithmetic with existence **and uniqueness** in full detail. Exercises and a Going further section at the end.

2.1 Prime numbers

Definition 1.2.1. An integer $p > 1$ is **prime** if its only positive divisors are 1 and p . An integer > 1 that is not prime is **composite**. (*The number 1 is neither one thing nor the other — the reason for excluding it becomes visible in Theorem 1.2.6.*)

Lemma 1.2.2 (the least divisor is prime). Every integer $n > 1$ has some prime divisor; in fact, its **least** divisor greater than 1 is prime.

Proof. The set of divisors of n greater than 1 is nonempty (n is in it); let p be its minimum (well-ordering). If p had a divisor d with $1 < d < p$, that d would divide n (transitivity, Prop. 1.1.2.1) and would be a divisor of n greater than 1 and smaller than p — against minimality. Hence p is prime. ■

Theorem 1.2.3 (Euclid: infinitude of the primes). There are infinitely many prime numbers.

Proof (by contradiction — the most famous in mathematics). Suppose there were only finitely many: $p_1, p_2, \dots, p_k$. Consider

$$N = p_1 p_2 \cdots p_k + 1.$$

Since $N > 1$, it has a prime divisor p (Lemma 1.2.2) — which must be one from the list. But each p_i leaves **remainder** 1 when dividing N (it divides the product, not the 1; Prop. 1.1.2.2 with signs). Contradiction: p is a prime outside the list, which was supposed to be complete. ■

M0 connection: this is the canonical example of proof by contradiction (and a good place to recall, in passing, the remark on the excluded middle). Note also the role of Lemma 1.2.2: the “popular-science” version of this proof usually skips it.

The proof is constructive (no excluded middle?). Although presented “by contradiction”, at bottom it **exhibits** a new prime: given any finite list $p_1, \dots, p_k$, the number $N = p_1 \cdots p_k + 1$ has (Lemma 1.2.2) a prime divisor, and that prime is **not** in the list (each p_i leaves remainder 1). It is a procedure that, from known primes, produces another — it does not require the excluded middle. The wording “suppose there are finitely many...” is a convenient wrapper, not a logical necessity. (*A nod to the constructivism of M0.*)

2.2 Euclid’s lemma — where Bézout pays off

Lemma 1.2.4 (general version). If $\gcd(c, a) = 1$ and $c \mid ab$, then $c \mid b$.

Proof. Bézout (Theorem 1.1.9): $1 = cx + ay$. Multiplying by b :

$$b = cbx + aby.$$

c divides the first summand (obvious) and the second (because $c \mid ab$); hence it divides the sum (Prop. 1.1.2.2): $c \mid b$. ■

Corollary 1.2.5 (Euclid’s lemma). If p is prime and $p \mid ab$, then $p \mid a$ or $p \mid b$.

Proof. If $p \nmid a$, then $\gcd(p, a) = 1$ (the positive divisors of p are 1 and p , and p does not divide a), and Lemma 1.2.4 gives $p \mid b$. ■

Structural counterfactual (the hypothesis is essential): $6 \mid 4 \cdot 9 = 36$, but $6 \nmid 4$ and $6 \nmid 9$. Composites do not penetrate products: their ability to “split themselves” between the factors ($6 = 2 \cdot 3$, with the 2 going to the 4 and the 3 to the 9) is exactly what the lemma forbids the primes. Being prime is not just “not factoring”: it is this property.




Figure 3: Via unique factorization, the primes of ab are those of a together with those of b ; if $p \mid ab$, p is one of them, hence it is “in” a or “in” b . For a composite like 6, its two primes can split between the factors.

Heuristic, not proof. It is a good mental image, but **circular as a proof** (the uniqueness of factorization is proved *with* this lemma): the rigorous argument is Bézout’s (Lemma 1.2.4).

2.3 The fundamental theorem of arithmetic

Theorem 1.2.6 (fundamental theorem of arithmetic). Every integer $n > 1$ can be written as a product of primes, $n = p_1 p_2 \cdots p_r$, and the expression is **unique up to the order** of the factors.

Proof. Existence (strong induction; see Preliminaries). For $n = 2$: prime, a product of one factor. Let $n > 2$ and suppose the claim true for all integers between 2 and $n - 1$. If n is prime, done. If not, $n = ab$ with $1 < a, b < n$; by the induction hypothesis, a and b are products of primes, and concatenating, so is n . (Here the **strong** version of induction is essential: a and b are smaller than n but not necessarily $n - 1$, so ordinary induction — which only grants the case $n - 1$ — would not suffice.)

Uniqueness. Suppose $n = p_1 \cdots p_r = q_1 \cdots q_s$ with all the p_i, q_j prime; we argue by induction on r . Since $p_1 \mid q_1(q_2 \cdots q_s)$, Euclid’s lemma (applied repeatedly) forces p_1 to divide **some** q_j ; reordering, $p_1 \mid q_1$, and since both are prime, $p_1 = q_1$. Cancelling (n/p_1) : $p_2 \cdots p_r = q_2 \cdots q_s$, a **smaller** number, to which the induction hypothesis applies: $r - 1 = s - 1$ and the factors coincide up to order. (Base case $r = 1$: $n = p_1$ prime admits no factorizations other than itself.) ■

Remark 1.2.7 (why 1 is not prime). If 1 counted as a prime, uniqueness would die on the spot: $6 = 2 \cdot 3 = 1 \cdot 2 \cdot 3 = 1 \cdot 1 \cdot 2 \cdot 3 = \cdots$. The exclusion is not a whim of nomenclature: it is the condition for Theorem 1.2.6 to be true as stated.

Remark 1.2.7b (negative integers and “units”). Theorem 1.2.6 is stated for $n > 1$ with positive primes, and then “up to order” is exact. If negative integers are allowed, uniqueness becomes “up to order **and a sign** ± 1 ”: the ± 1

are the **units** of $\mathbb{Z}$ (its invertible elements), and the general statement of unique factorization — in any unique factorization domain — is “up to order and units”. Note, in passing, that under Definition 1.2.1 **prime** and **composite** only classify the integers > 1 : a negative number such as -4 is neither prime nor composite — its arithmetic is read through its positive part, $-4 = (-1) \cdot 2^2$, with the sign set aside as a unit — and 0 and ± 1 also fall outside the classification. Here the positive case suffices; the term “unit” is formalized in ring theory (references in the Going further section).

Structural lens: $\mathbb{Z}$ has **atoms** (the primes) and unique composition. The complete skeleton — division with remainder $\rightarrow$ the Euclidean algorithm $\rightarrow$ Bézout $\rightarrow$ Euclid’s lemma $\rightarrow$ unique factorization — will be rebuilt, piece by piece and with the same proofs, in $K[x]$ (Polynomials module). When it arrives, it will not be a new topic: it will be this one, in disguise.

2.4 Practical consequences

Proposition 1.2.8 (gcd and lcm via exponents). For nonzero a, b , writing $|a| = \prod p_i^{\alpha_i}$ and $|b| = \prod p_i^{\beta_i}$ (exponents ≥ 0 , common primes), then

$$\gcd(a, b) = \prod p_i^{\min(\alpha_i, \beta_i)}, \quad \text{lcm}(a, b) = \prod p_i^{\max(\alpha_i, \beta_i)}, \quad \gcd \cdot \text{lcm} = |ab|.$$

(The sign of a, b is irrelevant for divisibility — hence the $|\cdot|$ and the $|ab|$, not ab , valid also with negatives. *Proof:* a divisor of a has, by the uniqueness in the fundamental theorem of arithmetic, exponents $\leq \alpha_i$; the rest is taking minima/maxima and the identity $\min + \max = \alpha + \beta$. Details: exercise 5.)

But beware the practical trap: this formula requires **factoring**, and factoring is computationally expensive; the Euclidean algorithm computes the gcd **without factoring**, and blazingly fast. That asymmetry — multiplication/gcd cheap, factoring expensive — is not an anecdote: it is the basis of the cryptography of Session 4.

Proposition 1.2.9 (primality test up to $\sqrt{n}$). If $n > 1$ has no prime divisor $p \leq \sqrt{n}$, then n is prime.

Proof. If $n = ab$ with $1 < a \leq b < n$, then $a^2 \leq ab = n$, so $a \leq \sqrt{n}$, and the least prime divisor of a (Lemma 1.2.2) is a prime divisor of n less than or equal to $\sqrt{n}$. ■

Exercises

1. ★ Factor 360 and 756; compute the gcd and lcm via exponents, and **redo the gcd with the Euclidean algorithm**. Compare the work of the two methods and explain which would scale better with 100-digit numbers.

2. ★ (Counterfactual.) Find **your own** counterexample to Euclid’s lemma with a **composite number** (different from $6 \mid 4 \cdot 9$) and point out how the composite “splits” between the factors.
3. ★ Is 221 prime? And 223? Decide with the test of Proposition 1.2.9, listing exactly which primes must be tried and why those suffice.
4. ★ Answer with the fundamental theorem of arithmetic: why is 1 not prime? (The correct answer is not “because the definition says so”.)
5. Complete the proof of Proposition 1.2.8 (divisors via exponents; the min + max identity).
6. Prove that $\sqrt{2}$ is irrational using the fundamental theorem of arithmetic: in $a^2 = 2b^2$, count the parity of the exponent of 2 on each side. (*Compare with the “classical” proof by contradiction from M0 — same conclusion, finer tool.*)
7. Prove that there are infinitely many primes of the form $4k + 3$. (*Guide: adapt Euclid’s proof with $N = 4p_1 \cdots p_k - 1$; you will need: a product of numbers of the form $4k + 1$ is of the form $4k + 1$.*)
8. If p is prime and $p \mid a^n$, prove that $p^n \mid a^n$. (*Iterated Euclid’s lemma + fundamental theorem of arithmetic.*)

Going further and references

Going further (the distribution of the primes). Knowing there are infinitely many opens the question of *how many*: the **prime number theorem** says that up to x there are approximately $x/\ln x$. Between the arithmetic of this session and that result lie a century and a half of analysis. A readable overview: Childs, or ch. 1 of Hardy–Wright, *An Introduction to the Theory of Numbers*.

Going further (unique factorization that fails). There are “worlds” where the fundamental theorem of arithmetic is false: in $\mathbb{Z}[\sqrt{-5}]$, $6 = 2 \cdot 3 = (1 + \sqrt{-5})(1 - \sqrt{-5})$ are two genuinely different factorizations. Uniqueness is not free: it depends on Euclid’s lemma, which depends on Bézout, which depends on division with remainder. Aluffi, ch. V.

References. Childs, ch. 5–6; Dorronsoro–Hernández, ch. 2–3.

Lecture Notes · Arithmetic · Session 3 — Congruences and $\mathbb{Z}/n\mathbb{Z}$

The heart of the module, with everything proved: the equivalence relation, the *well-definedness* of the quotient operations, the invertibility criterion (with the effective computation of inverses), the **zero divisor / non-invertible dichotomy** — which prepares the notion of a **field**, to be formalized later in the course —, $\mathbb{Z}/p\mathbb{Z}$ as a first field,

and Fermat's little theorem with its rearrangement proof. Exercises and a Going further section at the end.

3.1 Congruences

Definition 1.3.1. Fix $n \geq 2$: $a \equiv b \pmod{n}$ if $n \mid (a - b)$.

Proposition 1.3.2. $a \equiv b \pmod{n} \iff a$ and b leave the **same remainder** when divided by n .

Proof. Writing $a = nq_1 + r_1$, $b = nq_2 + r_2$ with $0 \leq r_i < n$ (Theorem 1.1.3): $a - b = n(q_1 - q_2) + (r_1 - r_2)$. If $r_1 = r_2$, then $n \mid (a - b)$. Conversely, if $n \mid (a - b)$, then $n \mid (r_1 - r_2)$ (Prop. 1.1.2.2), and since $|r_1 - r_2| < n$, it must be that $r_1 = r_2$ (the bounding trick from Session 1). ■

Proposition 1.3.3. Congruence modulo n is an **equivalence relation** on $\mathbb{Z}$ (the concept from M0, at work!).

Proof. *Reflexive:* $n \mid 0$. *Symmetric:* if $n \mid (a - b)$ then $n \mid (b - a)$ (the negative). *Transitive:* if $n \mid (a - b)$ and $n \mid (b - c)$, then $n \mid ((a - b) + (b - c)) = (a - c)$ (Prop. 1.1.2.2). ■

Definition 1.3.4. The **class** of a is $\bar{a} = \{b : b \equiv a \pmod{n}\} = \{a, a \pm n, a \pm 2n, \dots\}$. By Proposition 1.3.2 there are exactly n **classes**, one per remainder: $\bar{0}, \bar{1}, \dots, \overline{n-1}$.




Figure 4: Congruence modulo n : wrap the number line around a circle with n marks; integers congruent modulo n land on the same mark (here $n = 5$).

3.2 $\mathbb{Z}/n\mathbb{Z}$: operating with classes (and the toll of the quotient)

Definition 1.3.5. $\mathbb{Z}/n\mathbb{Z} = \{\bar{0}, \bar{1}, \dots, \overline{n-1}\}$, with

$$\bar{a} + \bar{b} := \overline{a + b}, \quad \bar{a} \cdot \bar{b} := \overline{a \cdot b}.$$

Proposition 1.3.6 (the operations are well defined). If $a \equiv a' \pmod{n}$ and $b \equiv b' \pmod{n}$, then $a + b \equiv a' + b'$ and $ab \equiv a'b' \pmod{n}$.

Proof. Sum: $(a + b) - (a' + b') = (a - a') + (b - b')$, a sum of two multiples of n . Product: the trick of **adding and subtracting the cross term**:

$$ab - a'b' = ab - a'b + a'b - a'b' = (a - a')b + a'(b - b'),$$

a combination of multiples of n (Prop. 1.1.2.2). ■

Why this is THE fine point: Definition 1.3.5 computes with **representatives** of the classes. If the result depended on the chosen representative, the definition would be nonsense. Proposition 1.3.6 is the **mandatory toll of every quotient construction** — a pattern worth internalizing here, in the most concrete example possible (M0: equivalence classes).

The properties of the operations (associative, commutative, distributive, identity elements $\bar{0}, \bar{1}$, negatives) are **inherited** from $\mathbb{Z}$ representative by representative. *Structural lens*: a set with these properties is called a **ring** — we drop the name in passing; its general theory is not part of this course.

3.3 Invertibles and zero divisors

Definition 1.3.7. $\bar{a} \in \mathbb{Z}/n\mathbb{Z}$ is **invertible** if there exists $\bar{x}$ with $\bar{a}\bar{x} = \bar{1}$. And $\bar{a} \neq \bar{0}$ is a **zero divisor** if there exists $\bar{b} \neq \bar{0}$ with $\bar{a}\bar{b} = \bar{0}$.

Theorem 1.3.8 (invertibility criterion). $\bar{a}$ is invertible in $\mathbb{Z}/n\mathbb{Z} \iff \gcd(a, n) = 1$. Moreover, in that case the inverse is **computed** with the extended Euclidean algorithm.

Proof. ($\Leftarrow$) Bézout: $1 = ax + ny$ for some x, y . Modulo n : $\bar{a}\bar{x} = \bar{1}$, that is, $\bar{a}\bar{x} = \bar{1}$ — and the Bézout coefficients **are** the inverse. ($\Rightarrow$) If $\bar{a}\bar{x} = \bar{1}$, then $n \mid (ax - 1)$, i.e. $ax - 1 = ny$: $1 = ax - ny$. Every common divisor of a and n divides that combination (Prop. 1.1.2.2), hence divides 1: the gcd is 1. ■

Example 1.3.9. Inverse of $\bar{7}$ in $\mathbb{Z}/30\mathbb{Z}$: the Euclidean algorithm gives $30 = 4 \cdot 7 + 2$, $7 = 3 \cdot 2 + 1$; working back up, $1 = 7 - 3 \cdot 2 = 7 - 3(30 - 4 \cdot 7) = 13 \cdot 7 - 3 \cdot 30$. Hence $\bar{7}^{-1} = \bar{13}$. *Check*: $7 \cdot 13 = 91 = 3 \cdot 30 + 1$ ✓.

Proposition 1.3.9b (linear congruences, the general case). The congruence $ax \equiv b \pmod{n}$ has a solution **if and only if** $d = \gcd(a, n)$ divides b ; and in that case it has **exactly d solutions** modulo n : if x_0 is the unique solution of $\frac{a}{d}x \equiv \frac{b}{d} \pmod{\frac{n}{d}}$ (unique because $\frac{a}{d}$ is invertible modulo $\frac{n}{d}$), the solutions are

$$x_0, x_0 + \frac{n}{d}, x_0 + 2\frac{n}{d}, \dots, x_0 + (d-1)\frac{n}{d} \pmod{n}.$$

Proof. Solving $ax \equiv b \pmod{n}$ is solving $ax + ny = b$ over the integers, which has a solution $\iff d \mid b$ (it is Bézout's equation, scaled: Session 1). If $d \mid b$, dividing **everything** by d yields the equivalent congruence $\frac{a}{d}x \equiv \frac{b}{d} \pmod{\frac{n}{d}}$, where $\gcd(\frac{a}{d}, \frac{n}{d}) = 1$: by Theorem 1.3.8 there is a unique solution x_0 modulo $\frac{n}{d}$. And one class modulo $\frac{n}{d}$ splits into exactly d classes modulo n : the numbers $x_0 + k\frac{n}{d}$ with $0 \leq k < d$. ■

Example 1.3.9c. $6x \equiv 9 \pmod{15}$: $d = \gcd(6, 15) = 3$ divides 9 ✓. Dividing by 3: $2x \equiv 3 \pmod{5}$, and since $\bar{2}^{-1} = \bar{3}$ in $\mathbb{Z}/5\mathbb{Z}$ ($2 \cdot 3 = 6 \equiv 1$), $x \equiv 9 \equiv 4 \pmod{5}$. The **three** solutions modulo 15: $x \in \{4, 9, 14\}$. *Check*: $6 \cdot 9 = 54 = 3 \cdot 15 + 9$ ✓.

Theorem 1.3.10 (the dichotomy). Let $\bar{a} \neq \bar{0}$ in $\mathbb{Z}/n\mathbb{Z}$. Then:

$$\bar{a} \text{ is a zero divisor} \iff \bar{a} \text{ is not invertible} \iff \gcd(a, n) > 1.$$

Proof. The second equivalence is Theorem 1.3.8 negated. We prove the first in both directions.

(Zero divisor $\Rightarrow$ not invertible.) Let $\bar{a}\bar{b} = \bar{0}$ with $\bar{b} \neq \bar{0}$. If $\bar{a}$ had an inverse, multiplying: $\bar{b} = (\bar{a}^{-1}\bar{a})\bar{b} = \bar{a}^{-1}(\bar{a}\bar{b}) = \bar{a}^{-1}\bar{0} = \bar{0}$ — contradiction.

(Not invertible $\Rightarrow$ zero divisor.) Let $d = \gcd(a, n) > 1$ and take $\bar{b} = \overline{n/d}$. Then $\bar{b} \neq \bar{0}$ (because $0 < n/d < n$, since $d > 1$), and

$$a \cdot \frac{n}{d} = \frac{a}{d} \cdot n \equiv 0 \pmod{n}$$

(it is an integer multiple of n : $a/d \in \mathbb{Z}$ because $d \mid a$). Hence $\bar{a}\bar{b} = \bar{0}$: a zero divisor. ■

Remark 1.3.11 (the dichotomy is special to $\mathbb{Z}/n\mathbb{Z}$). In $\mathbb{Z}$, the number 2 is **not** invertible and is **not** a zero divisor either: outside the $\mathbb{Z}/n\mathbb{Z}$'s the two notions need not fill the whole space. (This is exactly the caution that will reappear later, when the **fields** are presented in Linear Algebra: here it is already proved in the most concrete example.)




Figure 5: The reflex this section dismantles: in $\mathbb{Z}$ (or in $\mathbb{R}$!) a zero product betrays a zero factor, but in $\mathbb{Z}/6\mathbb{Z}$ we have $\bar{2} \cdot \bar{3} = \bar{0}$ with both factors nonzero — zero divisors, exactly what Theorem 1.3.10 detects via $\gcd > 1$.

Corollary 1.3.12 (the thread of fields begins). 1. If p is **prime**, every nonzero element of $\mathbb{Z}/p\mathbb{Z}$ is invertible — a finite world where **one can divide**. It gets a *name*: $\mathbb{Z}/p\mathbb{Z}$ is a **field** (the formal definition and its theory, in the Linear Algebra module). 2. If n is **composite**, $\mathbb{Z}/n\mathbb{Z}$ has zero divisors and is **not** a field.

Proof. (1) $0 < a < p$ and p prime $\Rightarrow \gcd(a, p) = 1$ (the only divisors of p are 1 and p) $\Rightarrow$ invertible (1.3.8). (2) $n = ab$ with $1 < a, b < n$ gives $\bar{a}\bar{b} = \bar{0}$ with both nonzero. ■

×	0	1	2	3
0	0	0	0	0
1	0	1	2	3
2	0	2	0	2
3	0	3	2	1

$\mathbb{Z}/4\mathbb{Z}$: there are zero divisors
($2 \cdot 2 = 0$, $2 \neq 0$)

×	0	1	2	3	4
0	0	0	0	0	0
1	0	1	2	3	4
2	0	2	4	1	3
3	0	3	1	4	2
4	0	4	3	2	1

$\mathbb{Z}/5\mathbb{Z}$: a field
(no product of nonzero elements is 0)

Figure 6: Multiplication tables of $\mathbb{Z}/4\mathbb{Z}$ and $\mathbb{Z}/5\mathbb{Z}$: in $\mathbb{Z}/4\mathbb{Z}$ there are zero divisors ($2 \cdot 2 = 0$); $\mathbb{Z}/5\mathbb{Z}$, since 5 is prime, is a field.

3.4 Fermat's little theorem


Theorem 1.3.13 (Fermat). If p is prime and $p \nmid a$, then

$$a^{p-1} \equiv 1 \pmod{p}.$$

Proof (the rearrangement). Consider the nonzero classes $\bar{1}, \bar{2}, \dots, \overline{p-1}$ and multiply them all by $\bar{a}$:

$$\bar{a} \cdot \bar{1}, \bar{a} \cdot \bar{2}, \dots, \bar{a} \cdot \overline{p-1}.$$

Claim: this list is the same as before, rearranged. Indeed, the map $\bar{x} \mapsto \bar{a}\bar{x}$ is **injective** ($\bar{a}\bar{x} = \bar{a}\bar{y} \Rightarrow \bar{x} = \bar{y}$, multiplying by $\bar{a}^{-1}$, which exists by Corollary 1.3.12.1) and never produces zero ($\bar{a}\bar{x} = \bar{0}$ with $\bar{a}$ invertible would force $\bar{x} = \bar{0}$); an injective map from a **finite** set to itself is bijective (M0). Equating the products of the two lists:



the same set, shuffled

Figure 7: For $p = 5$, $a = 2$: the row $(\bar{1}, \bar{2}, \bar{3}, \bar{4})$ is transformed into $(\bar{2}, \bar{4}, \bar{1}, \bar{3})$ by $x \mapsto 2x$ ($1 \rightarrow 2$, $2 \rightarrow 4$, $3 \rightarrow 1$, $4 \rightarrow 3$) — the same set, shuffled.

That is why the product of the whole row does not change, and that is what gets cancelled.

$$\overline{(p-1)!} = \overline{a^{p-1}} \cdot \overline{(p-1)!}.$$

And $\overline{(p-1)!}$ is invertible (a product of invertibles); cancelling it: $\overline{a^{p-1}} = \bar{1}$. ■

Note what the proof uses: that $\mathbb{Z}/p\mathbb{Z}$ is a **field** (all nonzero elements invertible, twice) and a **counting** argument (injective on a finite set = bijective, M0). **What it is for:** Fermat powers the modular exponentiation that sustains public-key cryptography — Session 4 collects the debt.

Exercises

1. ★ In $\mathbb{Z}/30\mathbb{Z}$, compute $\overline{17}^{-1}$ with the extended algorithm (not by trial and error) and check.
2. ★ Classify **all** the nonzero elements of $\mathbb{Z}/12\mathbb{Z}$ into invertibles and zero divisors; verify the dichotomy of Theorem 1.3.10 and exhibit, for each zero divisor, the partner $\bar{b}$ that betrays it.
3. ★ Compute the remainder of 3^{100} when divided by 7 (Fermat: $3^6 \equiv 1$) and that of 2^{1000} when divided by 11.

4. ★ **(The Linear Algebra counterfactual, now with theory.)** In $\mathbb{Z}/6\mathbb{Z}$: $\bar{2} \cdot \bar{3} = \bar{0}$. Locate the gcd that explains it via Theorem 1.3.10, and construct the “canonical” zero divisor n/d for $\bar{a} = \bar{2}$.
5. Prove that $\mathbb{Z}$ has no zero divisors, and conclude that the dichotomy 3.10 **fails** in $\mathbb{Z}$ (with 2 as the witness).
6. Prove the **divisibility rule for 9**: a number is divisible by 9 if and only if the sum of its digits is. (*Hint*: $10 \equiv 1 \pmod{9}$, so $10^k \equiv 1$.)
7. Is it true that $a \equiv b \pmod{n}$ implies $a^2 \equiv b^2 \pmod{n}$? And the converse? Prove or refute (minimal counterexample).
8. **(Python, asynchronous activity.)** Generate the multiplication tables of $\mathbb{Z}/5\mathbb{Z}$ and $\mathbb{Z}/6\mathbb{Z}$ (`[[a*b % n for b in range(n)] for a in range(n)]`), spot the zero divisors visually, compute inverses with `pow(a, -1, n)` (built in; it raises an error exactly when no inverse exists — why?), and check Fermat with `pow(a, p-1, p)` for several primes.

Going further and references

Optional going further (rings). A set with two operations $+, \cdot$ satisfying the properties inherited here (both associative, addition commutative, distributive, with identity elements $\bar{0}, \bar{1}$ and negatives) is a **commutative ring with identity**; $\mathbb{Z}$, $\mathbb{Z}/n\mathbb{Z}$ and (Polynomials module) $K[x]$ are examples. The **zero divisors** and the **units** (invertibles) of this session are general notions of ring theory; a commutative ring without zero divisors is an **integral domain**, and one where every nonzero element is invertible is a **field** (M4). To expand, optional: Dorronsoro–Hernández, *Números, grupos y anillos*, ch. 3; Childs, ch. 7–9; with a categorical flavor, Aluffi, *Algebra: Chapter 0*, ch. III. Not assessed.

Going further (Euler’s φ function). How many invertibles does $\mathbb{Z}/n\mathbb{Z}$ have? Exactly $\varphi(n) = \#\{1 \leq a \leq n : \gcd(a, n) = 1\}$. **Euler’s** theorem generalizes Fermat’s: $a^{\varphi(n)} \equiv 1 \pmod{n}$ if $\gcd(a, n) = 1$ — and the same rearrangement proof works, restricted to the invertibles. For $n = pq$ (the RSA case): $\varphi(pq) = (p-1)(q-1)$. Childs, ch. 9; Dorronsoro–Hernández.

References. Childs, ch. 7–9 (congruences, $\mathbb{Z}/n\mathbb{Z}$, Fermat); Dorronsoro–Hernández, ch. 3. The dichotomy and fields: Linear Algebra notes, Session 1 (where what is proved here gets used).

Lecture Notes · Arithmetic · Session 4 — The Chinese remainder theorem and cryptography (RSA)

The module's closing session with everything proved: the constructive CRT (with the uniqueness lemma), fast modular exponentiation with the why behind it, and the theorem that **RSA decrypts** — Fermat and the CRT glued together, with the toy example verified by hand and the code of the demo. Exercises and a Going further section at the end.

4.1 The Chinese remainder theorem

Theorem 1.4.1 (CRT). If $\gcd(m, n) = 1$, the system of congruences

$$x \equiv a \pmod{m}, \quad x \equiv b \pmod{n}$$

has a solution, and it is **unique modulo** mn (any two solutions are congruent modulo mn).

Proof. **Existence (constructive: the switches).** Bézout: $1 = mx_0 + ny_0$. Define

$$e_1 = ny_0, \quad e_2 = mx_0.$$

Then $e_1 + e_2 = 1$, and: $e_1 \equiv 1 \pmod{m}$ (since $e_1 = 1 - mx_0$) and $e_1 \equiv 0 \pmod{n}$; symmetrically $e_2 \equiv 0 \pmod{m}$, $e_2 \equiv 1 \pmod{n}$. They are **switches**: each is worth 1 in one modulus and 0 in the other. The combination

$$x = a e_1 + b e_2$$

satisfies $x \equiv a \cdot 1 + b \cdot 0 = a \pmod{m}$ and $x \equiv b \pmod{n}$. ■ (*existence*)




Figure 8: The CRT “switches”: e_1 is worth 1 modulo m and 0 modulo n , and e_2 the other way around; combining them, $x = a e_1 + b e_2$ solves the system.

Lemma 1.4.2 (for the uniqueness). If $m \mid x$, $n \mid x$ and $\gcd(m, n) = 1$, then $mn \mid x$.

Proof. $x = mt$; since $n \mid mt$ and $\gcd(n, m) = 1$, Lemma 1.2.4 gives $n \mid t$, i.e. $t = ns$: $x = mns$. ■

Uniqueness of the CRT. If x and x' solve the system, m and n divide $x - x'$, and by Lemma 1.4.2, $mn \mid (x - x')$. ■

Corollary 1.4.2b (CRT for k moduli). If $m_1, \dots, m_k$ are **pairwise coprime**, the system $x \equiv a_i \pmod{m_i}$ ($i = 1, \dots, k$) has a solution, unique modulo $m_1 m_2 \cdots m_k$.

Proof. Induction on k . The case $k = 2$ is Theorem 1.4.1. For the step, $M = m_1 \cdots m_{k-1}$ is coprime with m_k (a prime of m_k divides no m_i , $i < k$, by pairwise coprimality; fundamental theorem of arithmetic): solve the first $k - 1$ congruences by hypothesis (one congruence modulo M) and combine it with the k -th one via the two-moduli case. (*Switch version:* $e_i \equiv 1 \pmod{m_i}$ and $\equiv 0$ in the others, built with $\widehat{m}_i = \prod_{j \neq i} m_j$, which is invertible modulo m_i ; $x = \sum_i a_i e_i$.

Exercise 5.) ■

Example 1.4.3 (the riddle). $x \equiv 2 \pmod{3}$, $x \equiv 3 \pmod{5}$. Bézout for $(3, 5)$: $1 = 2 \cdot 3 - 1 \cdot 5$, so $e_1 = -5 \pmod{15} \equiv 10$ ($\equiv 1 \pmod{3}, \equiv 0 \pmod{5}$) and $e_2 = 6$. Solution: $x = 2(-5) + 3(6) = 8$, unique modulo 15: $x \in \{8, 23, 38, \dots\}$. Check: $8 = 2 + 2 \cdot 3$ ✓, $8 = 3 + 5$ ✓.

Non-coprime moduli (the hypothesis is not decorative). With $\gcd(m, n) > 1$ the system has a solution **if and only if** $\gcd(m, n) \mid (b - a)$ (exercise 6); when there is one, it is unique modulo $\text{lcm}(m, n)$.

- *Unsolvable:* $x \equiv 0 \pmod{2}$, $x \equiv 1 \pmod{4}$: $\gcd(2, 4) = 2 \nmid (1 - 0) = 1$ — no solution (x even and at the same time $\equiv 1 \pmod{4}$, impossible).
- *Solvable despite non-coprimality:* $x \equiv 2 \pmod{4}$, $x \equiv 4 \pmod{6}$: $\gcd(4, 6) = 2 \mid (4 - 2) = 2$ — there is a solution, unique modulo $\text{lcm}(4, 6) = 12$. Trying: $x = 10$ satisfies $10 \equiv 2 \pmod{4}$ and $10 \equiv 4 \pmod{6}$; the solutions are $x \equiv 10 \pmod{12}$.

4.2 Fast modular exponentiation

For $a^k \pmod{n}$ with k huge, the key is to **square repeatedly**, reducing modulo n at every step, and multiply the squares dictated by the **binary expansion** of k :

$$k = 13 = 8 + 4 + 1 = 1101_2 \implies a^{13} = a^8 \cdot a^4 \cdot a^1.$$

Example 1.4.4. $3^{13} \pmod{7}$: $3^2 = 9 \equiv 2$; $3^4 \equiv 2^2 = 4$; $3^8 \equiv 4^2 = 16 \equiv 2$. Then $3^{13} = 3^8 \cdot 3^4 \cdot 3 \equiv 2 \cdot 4 \cdot 3 = 24 \equiv 3 \pmod{7}$.

Cost: on the order of the **number of bits of k** — about $2 \log_2 k$ multiplications, and $\log_2 k$ is precisely that number of bits —: for a thousand-bit k , **a few thousand** operations, not 2^{1000} . Without this algorithm, RSA would be theory without practice.

4.3 RSA

What it solves: public-key cryptography

Before the scheme, the problem. To **encrypt** a message is to transform it with a key so that only whoever holds the right key can recover it. *Classical* (symmetric) cryptography uses the **same** secret key to encrypt and to decrypt: sender and receiver must **share it beforehand** over a secure channel — but if they already had a secure channel, a good part of the original problem would already be solved. That vicious circle — the **key-exchange problem** — limited cryptography for centuries.

The idea of **public-key** cryptography breaks it. Each user has a mathematically linked **pair** of keys: a **public** one, published openly, which anyone uses to encrypt messages for them, and a **private** one, kept secret, which is the only one capable of decrypting. No secret ever needs to be shared in advance. **RSA** (Rivest–Shamir–Adleman, 1978) was the first practical system to realize this idea, and it still protects — among many other things — the web connections with a “padlock”. What keeps the public key from betraying the private one is a **computational asymmetry** (multiplying is easy, factoring is not), which we make precise in “*Why it is secure*”.

The scheme

1. **Keys.** Choose primes $p \neq q$ (large); $n = pq$; $\varphi = (p-1)(q-1)$; choose e with $\gcd(e, \varphi) = 1$, and compute $d = e^{-1} \bmod \varphi$ — **the extended algorithm from S1/S3, once again.** **Public** key: (n, e) . **Private** key: d .
2. **Encrypt** (anyone): $c = m^e \bmod n$. **Decrypt** (only whoever holds d): $m = c^d \bmod n$. (Messages $0 \leq m < n$.)

Theorem 1.4.5 (RSA decrypts). With the keys built this way, for every m : $(m^e)^d \equiv m \pmod{n}$.

Proof. Since $ed \equiv 1 \pmod{\varphi}$, we write $ed = 1 + k(p-1)(q-1)$. **We work modulo p and modulo q separately, and glue with the CRT.**

Modulo p : if $p \mid m$, both sides are $\equiv 0$. If $p \nmid m$, Fermat (Theorem 1.3.13) gives $m^{p-1} \equiv 1$, so

$$m^{ed} = m \cdot (m^{p-1})^{k(q-1)} \equiv m \cdot 1 = m \pmod{p}.$$

Modulo q : identical. Thus p and q both divide $m^{ed} - m$; since $\gcd(p, q) = 1$, **Lemma 1.4.2** gives $pq \mid (m^{ed} - m)$, that is, $m^{ed} \equiv m \pmod{n}$. ■

Inventory of the proof: **Bézout** (built d), **Fermat** (the engine), **CRT/Lemma 1.4.2** (the glue). The entire module, in a ten-line proof.

Why it is secure (honesty)

To decrypt one needs d ; to compute d one needs $\varphi = (p-1)(q-1)$; and knowing φ together with n **is equivalent to factoring n** (exercise 8: from n and φ one recovers p and q by solving a quadratic). Security rests, then, on the **computational asymmetry** of Session 2: multiplying is instantaneous; factoring a 2048-bit n , infeasible with known technology.

Multiplying versus factoring (the asymmetry, with intuition). It is worth seeing why one direction is easy and the other is not. **Multiplying** two large numbers is fast: the work grows *smoothly* with the number of **digits** — the school algorithm uses on the order of (number of digits)² operations; one says it is **polynomial** in the size of the input (the “size” is the number of digits, not the value of the number). For the reverse path — given n , **finding its divisors** — no fast method is known: in essence one must **search** among candidates, and that search space grows **explosively** (exponentially) with the number of digits. The gap becomes abysmal as soon as the numbers are large: with two primes of about 300 digits each (≈ 1024 bits), multiplying them is instantaneous, while factoring the ~ 600 -digit product would exceed the age of the universe using all of today’s computers at once. *That* distance — trivial in one direction, intractable in the other — is the security of RSA. And it is robust against hardware progress: if the machines improve, **enlarging the keys** makes encryption slightly more expensive and factoring vastly more so.



Figure 9: The asymmetry, in one image: the forward task barely registers on the meter; the inverse one blows it up.

Note (quantum computing). “Infeasible” refers to **classical** computers. In 1994 Peter Shor discovered an algorithm that factors **efficiently**... on a **quantum** computer. That RSA still stands today only means that quantum machines large enough to factor real keys do not yet exist. That is why **post-quantum cryptography** — based

on problems conjectured to be hard even for quantum computers — is already being standardized. (Going further; cf. Childs.)

(Honest fine print: “equivalent de facto” — real-world RSA also carries random padding and implementation precautions without which there are attacks that do not require factoring. Ours is the mathematical core.)

Example 1.4.6 (a toy one, verified by hand)

$p = 3, q = 11$: $n = 33, \varphi = 20$. We take $e = 3$ ($\gcd(3, 20) = 1$); the extended algorithm gives $d = 7$ ($3 \cdot 7 = 21 \equiv 1 \pmod{20}$). **Encrypt** $m = 5$: $c = 5^3 = 125 \equiv 26 \pmod{33}$. **Decrypt**: $26^7 \pmod{33}$, by squaring ($26 \equiv -7$): $(-7)^2 = 49 \equiv 16$; $(-7)^4 \equiv 16^2 = 256 \equiv 25$; $26^7 \equiv 25 \cdot 16 \cdot (-7) \equiv 4 \cdot (-7) = -28 \equiv 5 \pmod{33}$ ✓ — the message comes back.

The Python demo (skeleton)

Everything is standard Python except `nextprime`, which comes from `sympy` (free: `pip install sympy`); modular exponentiation and the modular inverse are **built in** (three-argument `pow` and `pow(e, -1, phi)`).

```
from sympy import nextprime
from math import gcd

p = nextprime(10**6); q = nextprime(2 * 10**6)
n = p*q; phi = (p-1)*(q-1)
e = 65537; assert gcd(e, phi) == 1
d = pow(e, -1, phi)           # modular inverse (Bézout under the hood)
m = 271828                   # the "message"
c = pow(m, e, n)              # encrypt (public key)
assert pow(c, d, n) == m     # decrypt (private key) - m comes back
```

(Seed of the SE3 mini-project: each student generates their own keys and messages are exchanged; guided asynchronous activity.)

Exercises

- ★ Solve with the switches: $x \equiv 3 \pmod{7}, x \equiv 5 \pmod{11}$. Give the general solution and the least positive one; check.
- ★ RSA by hand with $p = 5, q = 11, e = 3$: build n, φ, d ; encrypt $m = 4$ and decrypt the result, all by fast exponentiation.
- ★ Compute $7^{45} \pmod{11}$ combining **Fermat** (reduce the exponent) and successive squaring.
- ★ (**Python — SE3 seed.**) Set up the full scheme with primes of 6–8 digits; encrypt a numeric message and verify the decryption. Then **break**

- someone else's toy RSA with `sympy.factorint(n)` and discuss: up to what size of n does factoring succeed on your machine within a second?
5. Complete the **switch construction** of Corollary 1.4.2b: for pairwise coprime $m_1, \dots, m_k$, define $\widehat{m}_i = \prod_{j \neq i} m_j$, find its inverse modulo m_i , form the e_i and verify that $x = \sum_i a_i e_i$ solves the system. Apply it to $x \equiv 2 \pmod{3}$, $x \equiv 3 \pmod{5}$, $x \equiv 2 \pmod{7}$.
 6. (**Counterfactual completed.**) Prove: $x \equiv a \pmod{m}$, $x \equiv b \pmod{n}$ has a solution **if and only if** $\gcd(m, n) \mid (b - a)$. (This explains exactly when the non-coprime case fails.)
 7. Why does RSA require $\gcd(e, \varphi) = 1$? Point out the step of the construction that would break without it.
 8. Prove that knowing $n = pq$ and $\varphi = (p - 1)(q - 1)$ allows one to recover p and q . (*Hint: $p + q = n - \varphi + 1$; then p, q are the roots of $t^2 - (p + q)t + n$.*)

Going further and references

Going further (from Fermat to Euler, and general RSA). Our Theorem 1.4.5 avoids the general φ function by using Fermat + CRT. The alternative route: **Euler's** theorem ($a^{\varphi(n)} \equiv 1$ if $\gcd(a, n) = 1$; Going further of Session 3) gives the case $\gcd(m, n) = 1$ in one line — but it does not cover the messages not coprime with n , which our proof does. Childs, ch. 9–10.

Going further (digital signatures). Swapping the roles of the keys (d to “encrypt”, e to verify), RSA produces **signatures**: only the owner of d could have generated the signature, and anyone can verify it with the public key. The same mathematics, read backwards. Childs, ch. 10.

References. CRT: Childs, ch. 8; Dorronsoro–Hernández. RSA: Childs, ch. 10 (the reference closest to the course); the original Rivest–Shamir–Adleman paper (1978) is surprisingly readable.