

## Question 3: Betting on Roulette

A casino offers a House Special bet on roulette, which is a bet on five pockets (00, 0, 1, 2, 3) out of 38 total pockets. The bet pays out 6 to 1. In other words, a losing bet yields -\$1 and a successful bet yields \$6. A gambler wants to know the chance of losing money if he places 500 bets on the roulette House Special.

The following 7-part question asks you to do some calculations related to this scenario.

### Question 3a

1/1 point (graded)

What is the expected value of the payout for one bet?

✓ Answer: -0.0789

### Explanation

The expected value can be calculated using the following code:

```
p <- 5/38
a <- 6
b <- -1
mu <- a*p + b*(1-p)
mu
```

You have used 1 of 10 attempts

❗ Answers are displayed within the problem

### Question 3b

1/1 point (graded)

What is the standard error of the payout for one bet?

✔ Answer: 2.37

### Explanation

The standard error can be calculated using the following code:

```
sigma <- abs(b-a) * sqrt(p*(1-p))  
sigma
```

You have used 1 of 10 attempts

---

**i** Answers are displayed within the problem

---

## Question 3c

1/1 point (graded)

What is the expected value of the average payout over 500 bets?

Remember there is a difference between expected value of the average and expected value of the sum.

✔ Answer: -0.0789

### Explanation

The expected value can be calculated using the following code:

```
mu
```

You have used 1 of 10 attempts

---

**i** Answers are displayed within the problem

---

## Question 3d

1/1 point (graded)

What is the standard error of the average payout over 500 bets?

Remember there is a difference between the standard error of the average and standard error of the sum.

✔ Answer: 0.106

### Explanation

The standard error can be calculated using the following code:

```
n <- 500  
sigma/sqrt(n)
```

Submit

You have used 1 of 10 attempts

**i** Answers are displayed within the problem

## Question 3e

1/1 point (graded)

What is the expected value of the sum of 500 bets?

-39.47368

✓ Answer: -39.5

-39.47368

### Explanation

The expected value can be calculated using the following code:

```
n*mu
```

Submit

You have used 1 of 10 attempts

**i** Answers are displayed within the problem

## Question 3f

1/1 point (graded)

What is the standard error of the sum of 500 bets?

52.91045

✓

52.91045

Submit

You have used 1 of 10 attempts

✓ Correct (1/1 point)

## Question 3g

1/1 point (graded)

Use `pnorm` with the expected value of the sum and standard error of the sum to calculate the probability of losing money over 500 bets,  $\Pr(X \leq 0)$ .

0.7721805

✓ Answer: 0.772

0.7721805

### Explanation

The standard error can be calculated using the following code:

```
pnorm(0, n*mu, sqrt(n)*sigma)
```

Submit

You have used 1 of 10 attempts

---

**i** Answers are displayed within the problem