

ID 2335100

Exam for Mathematical and Logical  
Foundations of Computer Science

Math

Date.  $4^0 = 1$   $1 \bmod 3 = 1$

Q.  $4^1 = 4$   $4 \bmod 3 = 1$

or  $4^2 = 16$   $16 \bmod 3 = 1$   $P(n) = 4^n \bmod 3$

$4^3 = 64$   $64 \bmod 3 = 1$

$\therefore 1 \in \mathbb{N}$

$4^4 = 256$   $256 \bmod 3 = 1$

$\therefore 4^n \bmod 3$

Base case:  $4^n \bmod 3 = 1, 1 \in \mathbb{N}$

induction case: assume  $4^2 \bmod 3 = n \in \mathbb{N}$  is true

show the  $4^n \bmod 3 = n \in \mathbb{N}$  is true too

$P(n)$

(ii)

$\square [x \in \mathbb{N} \mid x \geq 0] \rightarrow [y \in \mathbb{R} \mid y \geq 0]$

$\text{sqrt}(x) = y$   $x = y^2$   $x^2 > 0$

every element only has one output

$\text{sqrt}$  is injective because  $x = y^2 = x$  two different value has same answer

$\text{sqrt}$  is ~~not~~ surjective because ~~every possible output is obtained~~

$\text{sqrt}$  is ~~not~~ injective and is surjective so is not bijective

(iii)  $\text{sqrt}$  has infinitely many elements,  $\text{int}$  has  $2^{32}$  many

(1) if the parameter is ~~not~~ <sup>positive</sup> infinity, then the result is ~~not~~ <sup>positive</sup> infinity

(2) if the argument passed is positive or negative zero the result will be same as the of the argument but

(4) for normal positive numbers, to return the double value closest to the true math square root of the argument value if the input was perfect square  $\text{int}$ , the result will be an integer valued double that is in  $\text{int}$  range

(5)  $\text{sqrt}$  contains uncomputable elements, whereas every total is the possible result of computation

for  $\text{sqrt} : \mathbb{R}$

Q (b)

i)

$$P_1 \begin{pmatrix} 1 \\ 2 \\ 4 \end{pmatrix} \quad P_2 = \begin{pmatrix} -1 \\ 0 \\ 4 \end{pmatrix} \quad P_3 \begin{pmatrix} -1 \\ 2 \\ 2 \end{pmatrix}$$

$$\overrightarrow{P_1 P_2} \quad \overrightarrow{P_2 P_3} \quad \overrightarrow{P_1 P_3}$$

$$\begin{pmatrix} -2 \\ -2 \\ 0 \end{pmatrix} \quad \begin{pmatrix} 0 \\ 2 \\ -2 \end{pmatrix} \quad \begin{pmatrix} -2 \\ 0 \\ -2 \end{pmatrix}$$

$$|\overrightarrow{P_1 P_2}| = \sqrt{(-2)^2 + (-2)^2 + 0^2} = 2\sqrt{2}$$

$$|\overrightarrow{P_2 P_3}| = \sqrt{0^2 + 2^2 + (-2)^2} = 2\sqrt{2}$$

$$|\overrightarrow{P_1 P_3}| = \sqrt{(-2)^2 + 0^2 + (-2)^2} = 2\sqrt{2}$$

so that these is equilateral triangle

ii)

$$\vec{r} = \begin{pmatrix} 1 \\ 2 \\ 4 \end{pmatrix} + s \begin{pmatrix} -2 \\ -2 \\ 0 \end{pmatrix} + t \begin{pmatrix} -2 \\ 0 \\ -2 \end{pmatrix}$$

parametric represent

normal form:

$$ax_1 + bx_2 + cx_3 = d$$

$$a = \begin{matrix} 4 & -0 \\ 0 & -4 \end{matrix} = 4$$

$$b = \begin{matrix} 0 & -4 \\ 0 & -4 \end{matrix} = -4$$

$$c = 0 - 4 = -4$$

$$d = \begin{pmatrix} 1 \\ 2 \\ 4 \end{pmatrix} \cdot \begin{pmatrix} 4 \\ -4 \\ -4 \end{pmatrix}$$

$$= 4 - 8 - 16 = -20$$

$$4x_1 - 4x_2 - 4x_3 = -20$$

$$x_1 - x_2 - x_3 = -5$$

Date: / /

(ii)

$$N \begin{pmatrix} 1 \\ 0 \\ 3 \end{pmatrix} + q \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} + r \begin{pmatrix} 2 \\ 0 \\ 1 \end{pmatrix}$$

$$M \begin{pmatrix} 1 \\ 2 \\ 4 \end{pmatrix} + s \begin{pmatrix} -2 \\ -2 \\ 0 \end{pmatrix} + t \begin{pmatrix} -2 \\ 0 \\ -2 \end{pmatrix}$$

$$M = \begin{pmatrix} 1 & -2s & -2t \\ 2 & -2s \\ 4 & -2t \end{pmatrix} \quad N = \begin{pmatrix} 1+q+2r \\ q \\ 3+q+r \end{pmatrix}$$

1:  $1 - 2s - 2t = 1 + q + 2r$  3:  $4 - 2t = 3 + q + r$   
 $-2s - 2t - q - 2r = 0$   $q + r + 2t = 1$

2:  $2 - 2s = q$   
 $q + 2s = 2$

3:  $4 - 2t = 3 + q + r$   
 $q + r + 2t = 1$

row 2 + row 3

$$\begin{bmatrix} -2 & -2 & -1 & -2 & 0 \\ 0 & -2 & 0 & -2 & 2 \\ 0 & 0 & 1 & -2 & 3 \end{bmatrix} \xleftarrow{\text{row 1} + \text{row 2}} \begin{bmatrix} -2 & -2 & -1 & -2 & 0 \\ 0 & -2 & 0 & -2 & 2 \\ 0 & 0 & 1 & -2 & 3 \end{bmatrix}$$

row 2 + row 1

$$\begin{bmatrix} -2 & 0 & -1 & 0 & 2 \\ 0 & -2 & 0 & -2 & 2 \\ 0 & 0 & 1 & -2 & 3 \end{bmatrix} \xrightarrow{\text{row 3} - \text{row 1}} \begin{bmatrix} -2 & 0 & 0 & -2 & 1 \\ 0 & -2 & 0 & -2 & 2 \\ 0 & 0 & 1 & -2 & 3 \end{bmatrix}$$

$\begin{cases} -2s - 2r = 1 \\ -2t - 2r = 2 \\ q - 2r = 3 \end{cases} \Rightarrow \begin{cases} -2s = 1 & s = -\frac{1}{2} \\ -2t = 2 & t = -1 \\ q = 3 & q = 3 \end{cases}$



Q2 when  $q=3$

$$l = \begin{pmatrix} 1 \\ 0 \\ 3 \end{pmatrix} + 3 \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} + r \begin{pmatrix} 2 \\ 0 \\ 1 \end{pmatrix}$$
$$l = \begin{pmatrix} 4 \\ 3 \\ 6 \end{pmatrix} + r \begin{pmatrix} 2 \\ 0 \\ 1 \end{pmatrix}$$



Q3

(i)

$$\begin{array}{l} \frac{\forall x. X < succ(x)}{X = 0} \\ \frac{X = 0}{\exists y. X < y} [I] \\ \frac{\exists y. X < y}{\neg \exists x. \neg (X < x)} [E] \\ \frac{\neg \exists x. \neg (X < x)}{\neg \exists x. \forall y. X < y} [I] \\ S_1 \rightarrow \neg \exists x. \forall y. X < y \end{array}$$

$$(i) \quad (M,)$$

$$\models M_1 \rightarrow (\forall x. \forall y. x < y \rightarrow x \leq y)$$

$$M_1: \langle \mathbb{Q}, \langle F_{\text{zero}}, F_{\text{succ}} \rangle, R_1 \rangle$$

$$D = \mathbb{N} \quad F_{\text{zero}} = 0 \quad F_{\text{succ}} = \langle n \rangle \rightarrow n+$$

$$R \leq = \{ \langle n, m \rangle \mid n \leq m \}$$

$$M = \langle N, 0, +1 \rangle \langle n, m \rangle \mid \cancel{n \leq m} \wedge n < m \rangle$$

where  $+1$  is the function that give a number increment it by 1  
for all  $n, m \in \mathbb{D}$  if  $(n, m) \in R$  then  $F_{succ}(n) = m \in R$


Therefore for all  $n, m \in \mathbb{N}$  if  $(n, m) \in (n, m) \{ (n, m) | n < m \}$  then

$$\in \{(n, m) \mid n \leq m\} \quad (n+1, m)$$

for all  $n, m \in \mathbb{N}$  if  $n < m$  then  $n+1 < m$

which is false for Model  $M_7$  ( $\forall x, y, x \leq y \rightarrow x = y$ )





### Statement of good academic conduct

By submitting this assignment, I understand that I am agreeing to the following statement of good academic conduct:

- I confirm that this assignment is **my own work** and I have not worked with others in preparing this assignment.
- I confirm this assignment was written by me and is in my own words, except for any materials from published or other sources which are clearly indicated and acknowledged as such by appropriate referencing.
- I confirm that this work is not copied from any other person's work (published or unpublished), web site, book or other source, and has not previously been submitted for assessment either at the University of Birmingham or elsewhere.
- I confirm that I have not asked, or paid, others to prepare any part of this work for me.
- I confirm that I have read and understood the University regulations on plagiarism (<https://intranet.birmingham.ac.uk/as/registry/policy/conduct/plagiarism/index.aspx>).