TWO HIDDEN LAYERS SUFFICE: THE CATERPILLAR NET

Jakob Paul Zimmermann
FU Berlin, TU Berlin
jakob.paul.zimmermann@campus.tu-berlin.de
https://jakob-paul-zimmermann.com

ABSTRACT

Every continuous piecewise linear (CPWL) function is a finite signed sum of maxima of affine functions, making maximum representation a central problem in the study of the expressiveness of ReLU networks. Earlier work disproved the conjectured base-two depth bound, leaving open whether a constant number of hidden layers suffices. Inspired by the proof by Brunck et al. (2026) that two hidden layers suffice for general CPWL functions, we give the *Caterpillar Net*, a nonrecursive construction for the maximum of N inputs with two hidden layers of widths $O(N^2)$ and $2^{O(N \log N)}$. The construction has a direct geometric interpretation.[1]

1 INTRODUCTION

In this paper we investigate the representation capabilities of ReLU networks: What functions are representable by such networks of different depth and widths? As ReLU neural networks represent continuous piecewise-linear (CPWL) functions, a sharp question is: given any CPWL function, what is the ReLU network of smallest depth or smallest width representing this function? The maximum is a universal building block for such CPWL functions. Write

$$\max_N(x) := \max_{0 \le i < N} x_i. \tag{1}$$

Theorem (Wang and Sun, 2005)**.** *Let $d \ge 1$ and let $f : \mathbb{R}^d \to \mathbb{R}$ be continuous and affine on each member of a finite polyhedral cover. There are $R \in \mathbb{N}$, affine maps and signs*

$$A_t : \mathbb{R}^d \longrightarrow \mathbb{R}^{d+1}, \qquad \varepsilon_t \in \{-1, 1\} \qquad (1 \le t \le R) \tag{2}$$

such that

$$f(x) = \sum_{t=1}^{R} \varepsilon_t \max_{d+1}(A_t(x)) \qquad \textit{for every } x \in \mathbb{R}^d. \tag{3}$$

Thus a two-hidden-layer representation of $\max_{d+1}$ gives one for every CPWL function on $\mathbb{R}^d$: precompose with the affine maps and add the networks in parallel.

Historical conjecture (Hertrich et al., 2023, Conjecture 1.5, disproved in 2025)**.** *For every integer $k \ge 1$, the function $x \mapsto \max\{0, x_1, \ldots, x_{2^k}\}$ cannot be represented by a ReLU network with k hidden layers, regardless of their widths.*

The case $k = 2$ was already disproved in the May 2025 preprint of Bakaev et al. (2026), later published at STOC 2026, by a two-hidden-layer representation of the maximum of five inputs. Their improved general upper bound still grows logarithmically, so this did not settle whether a weaker $\Omega(\log N)$ lower bound holds, or even whether the required depth grows with N at all. Their concluding discussion explicitly left open whether constant depth, perhaps two hidden layers, suffices for all CPWL functions (Bakaev et al., 2026). Here constant depth means a layer count independent of N, with widths allowed to grow.

[1] Our results were obtained through proof search in Lean using Claude Code and Codex. The Lean code is available in the accompanying repository https://github.com/JayPiZimmermann/Max_Caterpillar. Links to the Lean formalization appear in theorem titles and equation numbers; equation numbers are clickable.

This reduction made the geometry of the maximum a central approach to the representation problem well before the work of Brunck et al. (2026). Polyhedral approaches to depth include the Newton-polytope framework of Hertrich et al. (2023), the simplex subdivisions of Bakaev et al. (2026), and the depth lower bounds for braid-fan-compatible networks of Grillo et al. (2025). For width, Safran (2026) used extremal graph theory to prove lower bounds for exact maximum representation, while Lu (2026) studied first hidden layer width thresholds for approximation and exact representation. Brunck et al. (2026) then constructed two-hidden-layer ReLU representations for arbitrary CPWL functions in every dimension.

Theorem 1 (Brunck et al., 2026)**.** *Every continuous piecewise affine function $f : \mathbb{R}^d \to \mathbb{R}$ with finitely many affine pieces can be represented exactly by a ReLU network with two hidden layers.*

Our contribution is to specialize their general construction to $\max_N$ and simplify it into the *Caterpillar Net*. Taking $2^{\Theta(N^2)}$ as the second hidden layer width baseline via applying the Brunck et al. (2026) recursive construction, we pursue two aims:

1. Give a nonrecursive construction for $\max_N$ with a direct geometric interpretation.
2. Reduce its second hidden layer width.

Write $\sigma(t) := \max(0, t)$ applied coordinatewise to vectors. A network has an input layer of N coordinates, first and second hidden layers of widths h_1 and h_2, respectively, and a scalar output layer. Its parameters are

$$\begin{aligned} &W_1 \in \mathbb{R}^{h_1 \times N}, \quad b_1 \in \mathbb{R}^{h_1}, \\ &W_2 \in \mathbb{R}^{h_2 \times h_1}, \quad b_2 \in \mathbb{R}^{h_2}, \\ &w_3 \in \mathbb{R}^{h_2}, \qquad b_3 \in \mathbb{R}, \end{aligned} \tag{4}$$

and evaluates as the function

$$x \longmapsto w_3^\top \sigma\big(W_2 \sigma(W_1 x + b_1) + b_2\big) + b_3. \tag{5}$$

The first and second hidden layers apply ReLU, and the output layer is affine. There are no skip connections, and equality is required on all of $\mathbb{R}^N$.

Theorem 2 (Caterpillar Net, `Lean_1`, `Lean_2`, proof)**.** *For every $N \geq 3$, define*

$$C_N = \sum_{m=2}^{N-1} \binom{N}{m} \frac{m!}{2} (N - m) m^{N-m-1}. \tag{6}$$

There is a two-hidden-layer ReLU network computing $\max_N$ with widths

$$h_1 = \binom{N}{2} + N + 1, \qquad h_2 = 3^{N-2} C_N - N(N-1) + 2. \tag{7}$$

In particular, these widths satisfy, as $N \to \infty$,

$$h_1 = O(N^2), \qquad h_2 = 2^{O(N \log N)}. \tag{8}$$

Remark (small arities). The linked theorem also covers $N = 1$ and $N = 2$ by separate networks of widths $(2, 2)$ and $(4, 2)$. At $N = 3$, formula (7) gives widths $(7, 5)$.

2 The Caterpillar Net

Two one-line identities drive the construction. For reals s, t and every linear function ℓ on $\mathbb{R}^N$,

$$\max(s, t) = t + \sigma(s - t), \tag{9}$$

$$\sigma(\ell(x)) = \sigma(-\ell(x)) + \ell(x). \tag{10}$$

The first suggests computing the maximum along a path with attached leaves. Alternating sums of support functions of caterpillar polytopes give the correct support-function class; support functions of their proper faces, together with linear functions, correct the remaining error to pointwise equality. The second reverses an inner hinge at the cost of a linear term; it merges equal units in Section 4.

Fix $N \geq 3$ and write $x = (x_0, \ldots, x_{N-1}) \in \mathbb{R}^N$.

Definition 3 (Labelled caterpillar, `Lean_3`)**.** A *caterpillar description* is a permutation–set pair

$$\pi \in \mathrm{Sym}\{0, \ldots, N-1\}, \qquad \varnothing \neq S \subseteq \{1, \ldots, r\}, \qquad r := N - 2. \tag{11}$$

Position j carries the label $\pi(j+1)$, for $0 \leq j \leq r$, and the label $\pi(0)$ is the *apex*. The *spine positions* are $\{0\} \cup S$. We also call S the *cut set*: a cut at $j \in S$ separates positions $j-1$ and j in the ordered list. The last spine position, and the *parent* of a position $j \geq 1$, the nearest spine position before it, are

$$q := \max S, \qquad p(j) := \max\big(\{0\} \cup \{i \in S : i < j\}\big) \qquad (1 \leq j \leq r). \tag{12}$$

A graph on the set of labels with an edge from each position $j \geq 1$ to its parent gives a directed tree, the *labelled caterpillar*: its spine is the path through the spine positions, and every other label is a leaf attached to the preceding spine vertex (Figure 1). The apex is separate from this tree.

Definition 4 (Caterpillar function and face functions, `Lean_4`, `Lean_5`)**.** For labels i, j, for the tree edges, and for the head, put

$$\ell_{ij}(x) := x_i - x_j, \tag{13}$$

$$L_{\pi,S} := \big\{\ell_{\pi(p(j)+1),\pi(j+1)} : 1 \leq j \leq r\big\}, \tag{14}$$

$$h_{\pi,S} := \ell_{\pi(q+1),\pi(0)}. \tag{15}$$

The *edge functions* $L_{\pi,S}$ subtract each child's coordinate from its parent's, one for each of the r tree edges, and the *head function* $h_{\pi,S}$ compares the last spine vertex with the apex. For a linear function h, a finite set L of linear functions, and a *state* $\tau(\ell) \in \{0,\ \ell,\ \sigma \circ \ell\}$ chosen for every $\ell \in L$, define

$$U^{\tau}_{h,L}(x) := \sigma\Big(h(x) + \sum_{\ell \in L} \tau(\ell)(x)\Big). \tag{16}$$

Thus each $\ell \in L$ is omitted, kept linearly, or kept as a hinge. Write $U_{h,L} := U^{\tau}_{h,L}$ for the all-hinge state $\tau(\ell) = \sigma \circ \ell$. We call (h, L) a *function description* of $U_{h,L}$: it records the linear head and the inner hinges. The caterpillar description (π, S) specifies the labelled tree from which such functions are generated. The *caterpillar function* of (π, S) and its *face functions* are

$$U_{\pi,S} := U_{h_{\pi,S},L_{\pi,S}}, \qquad U^{\tau}_{\pi,S} := U^{\tau}_{h_{\pi,S},L_{\pi,S}}. \tag{17}$$





Figure 1: A labelled caterpillar with $N = 10$ and $S = \{3, 6\}$. The spine $\pi(1) \to \pi(4) \to \pi(7)$ has two leaves at each vertex; all arrows point from parent to child. The apex $\pi(0)$ is separate from the tree. The last spine position is $q = 6$, so the head function is $h_{\pi,S}(x) = x_{\pi(7)} - x_{\pi(0)}$.

The term *face function* comes from the following geometry. Identify a linear function $a(x) = \sum_i a_i x_i$ with its coefficient vector $a = (a_0, \ldots, a_{N-1})$, so that $a(x) = \langle a, x \rangle$. In particular, ℓ_{ij} is the vector $e_i - e_j$: it can be drawn as an arrow from e_j to e_i. For a caterpillar, $h = h_{\pi,S} = e_{\pi(q+1)} - e_{\pi(0)}$ points from the apex label to the last spine label, and each $\ell \in L = L_{\pi,S}$ points from a child label to its parent label. In the polytope construction these are free vectors: h places one corner of the base, and the vectors ℓ generate its edges. The polytope apex is the origin, distinct from the labelled point $e_{\pi(0)}$ used to define the head. For a polytope P, its *support function*, and the polytope whose support function is $U_{h,L}$, are

$$h_P(x) := \max_{p \in P} \langle p, x \rangle, \tag{18}$$

$$Q_{h,L} := \mathrm{conv}\Big(\{0\} \cup \Big(h + \sum_{\ell \in L} [0, \ell]\Big)\Big), \qquad U_{h,L} = h_{Q_{h,L}}. \tag{19}$$

Indeed, the segment $[0, \ell]$ has support function $\sigma \circ \ell$, Minkowski sums add support functions, and adjoining 0 applies the outer ReLU. The base is the translated zonotope $h + \sum_{\ell \in L} [0, \ell]$. For a state τ in (16), set $t_\ell = \mathbb{1}_{\{\tau(\ell) = \ell\}}$, where $\mathbb{1}$ denotes the indicator. Thus $t_\ell = 1$ for an edge kept linearly and $t_\ell = 0$ for an omitted edge or a hinge; the hinges contribute segments separately. The corresponding polytope is $Q^\tau_{h,L} := \mathrm{conv}\big(\{0\} \cup (h + \sum_{\ell \in L} t_\ell \ell + \sum_{\tau(\ell) = \sigma \circ \ell} [0, \ell])\big)$, and $U^\tau_{h,L} = h_{Q^\tau_{h,L}}$. Thus a generator drawn between two labelled unit vectors is translated to the base corner h when drawing the zonotope. Figure 2 illustrates this for $N = 4$ in the coordinates obtained by restricting to $x_0 = 0$: a coefficient vector a becomes (a_1, a_2, a_3), so e_0 is represented by 0.



(a) Coordinate differences (b) Original description (c) Normalized description

Figure 2: Coordinate differences and the caterpillar polytope for $\pi = (0, 3, 2, 1)$ and $S = \{1, 2\}$. (a) The head $h = e_1 - e_0$ and edge functions $L = \{\ell_{32} = e_3 - e_2, \ell_{21} = e_2 - e_1\}$ drawn between their labelled endpoints. (b) The same generators translated to the base corner h of $Q_{h,L}$. (c) Normalization (Section 4) gives $h^+ = h + \ell_{32} + \ell_{21} = e_3 - e_0$ and $L^+ = \{\ell_{12} = e_1 - e_2, \ell_{23} = e_2 - e_3\}$, leaving the polytope unchanged. Purple arrows represent the head (the apex–last spine connection before normalization); blue arrows represent the spine edge functions. Orange dots mark the polytope apex and purple dots the chosen head corner. The grey coordinate axes meet at the origin, representing e_0 after restriction to $x_0 = 0$; the positive unit points are labelled e_1, e_2, e_3.

The head and the $N - 2$ edge functions are jointly linearly independent: their label edges form a tree on all N labels, since the head connects the apex to the last spine vertex. Thus the base is an $(N-2)$-dimensional zonotope whose affine hull does not contain 0, and $Q_{h,L}$ is an $(N-1)$-dimensional pyramid. A face of $Q_{h,L}$ containing the apex 0 and another point is the convex hull of 0 and a face of the base. Since the base is a translated sum of segments, the face exposed by a linear functional is obtained by maximizing on each segment separately, which selects the endpoint 0, the endpoint ℓ, or the whole segment. These have support functions 0, ℓ, and $\sigma \circ \ell$, and the outer ReLU adjoins the apex. Hence the face functions (16) include the support function of every face of $Q_{h,L}$ containing the apex.

Definition 5 (Dictionary, `Lean_6`)**.** The *dictionary* $\mathcal{D}$ is the finite set of functions $\mathbb{R}^N \to \mathbb{R}$ consisting of every face function $U^\tau_{\pi,S}$, over all caterpillar descriptions (π, S) and all states τ, together with the *auxiliary functions*

$$\begin{aligned} &x \mapsto \sigma(x_i), \quad x \mapsto \sigma(-x_i) \quad (0 \le i < N), \\ &x \mapsto \sigma(x_i - x_j) \qquad\qquad\quad (0 \le i < j < N). \end{aligned} \tag{20}$$

Heads and edge functions are coordinate differences, so face functions and the functions $x \mapsto \sigma(x_i - x_j)$ are translation invariant: with $\mathbf{1} := (1, \dots, 1) \in \mathbb{R}^N$,

$$U^\tau_{\pi,S}(x + t\mathbf{1}) = U^\tau_{\pi,S}(x) \qquad (t \in \mathbb{R}). \tag{21}$$

Only the coordinate features $\sigma(\pm x_i)$ change under $x \mapsto x + t\mathbf{1}$. Theorem 7 below proves $\max_N \in \mathrm{span}\, \mathcal{D}$; the following rule then fixes one real coefficient for each dictionary function.

A *tag* is a finite description of a dictionary feature, retaining the data needed to construct it in the network. The tag set is $\mathcal{A} = \mathcal{A}_{\mathrm{face}} \sqcup \mathcal{A}_{\mathrm{aux}}$, with two types: a *face tag* records a pair (h', M) representing $\sigma(h' + \sum_{\ell \in M} \sigma \circ \ell)$, after absorbing linear edges into h' and retaining only hinges in M as in (90); there is one *auxiliary tag* for each indexed feature in (20). The evaluation map sends

each description to its function:

$$v : \mathcal{A} \longrightarrow \mathcal{D}, \qquad \mathcal{D} = v(\mathcal{A}). \tag{22}$$

Different tags may represent the same function, so v need not be injective.

An exact coefficient rule (`Lean_7`) Let

$$\mathcal{B} := \{S \subseteq \mathcal{D} : S \text{ is a basis of } \operatorname{span} \mathcal{D}\}. \tag{23}$$

This set is nonempty because the finite spanning set $\mathcal{D}$ contains a basis. For $S \in \mathcal{B}$, form

$$G_S := \big(\langle g, h\rangle\big)_{g,h\in S}, \tag{24}$$

$$b_S := \big(\langle g, \max_N\rangle\big)_{g\in S}. \tag{25}$$

using the inner product

$$\langle f, g\rangle := \int_{[-1,1]^N} f(x)g(x)\,dx. \tag{26}$$

Every dictionary function is continuous and positively homogeneous. If a linear combination of the basis functions has zero squared integral, continuity makes it zero throughout the cube, and positive rescaling then makes it zero on all of $\mathbb{R}^N$. Linear independence forces every coefficient to vanish, so G_S is positive definite and invertible. Write G_S^{-1} for its inverse matrix. For each $g \in \mathcal{D}$, average over the bases containing g:

$$c_g := \frac{1}{|\mathcal{B}|} \sum_{\substack{S\in\mathcal{B}\\ g\in S}} (G_S^{-1} b_S)_g. \tag{27}$$

Lemma 6 (Canonical expansion, `Lean_8`)**.** *Assuming Theorem 7, the coefficients* (27) *satisfy*

$$\max_N = \sum_{g\in\mathcal{D}} c_g g. \tag{28}$$

Proof of Lemma 6. Fix $S \in \mathcal{B}$. By Theorem 7, there is $d \in \mathbb{R}^S$ such that

$$\max_N = \sum_{g\in S} d_g g. \tag{29}$$

Taking the inner product with each $h \in S$ gives

$$\begin{aligned} (b_S)_h &= \sum_{g\in S} \langle h, g\rangle d_g = (G_S d)_h, \\ d &= G_S^{-1} b_S. \end{aligned} \tag{30}$$

Interchanging the finite sums in (27) therefore yields

$$\sum_{g\in\mathcal{D}} c_g g = \frac{1}{|\mathcal{B}|} \sum_{S\in\mathcal{B}} \sum_{g\in S} (G_S^{-1} b_S)_g\, g = \frac{1}{|\mathcal{B}|} \sum_{S\in\mathcal{B}} \max_N = \max_N . \tag{31}$$

This finishes the proof of Lemma 6. □

For each tag, split the coefficient of its represented function equally over its nonempty finite fiber:

$$\widetilde{c}_a := \frac{c_{v(a)}}{|\{b \in \mathcal{A} : v(b) = v(a)\}|}. \tag{32}$$

While computationally intractable to extract explicit weights for large N, averaging over all bases avoids arbitrary symmetry-breaking and provides a canonical coefficient map ideal for formal verification. Regrouping by the fibers of v gives the literal tagged expansion

$$\max_N = \sum_{a\in\mathcal{A}} \widetilde{c}_a v(a). \tag{33}$$

Thus duplicate face and auxiliary tags neither duplicate nor lose any coefficient.

3 WHY THE DICTIONARY SPANS THE MAXIMUM

The maximum $\max_N$ is the support function of $\mathrm{conv}\{e_0, \dots, e_{N-1}\}$. To work with a full dimensional polytope, let us do the following substitution. First, let us define

$$\Delta := \mathrm{conv}\{0, e_1, \dots, e_{N-1}\} \subseteq H_0 := \{x \in \mathbb{R}^N : x_0 = 0\}. \tag{34}$$

Observe that for arbitrary input $x \in \mathbb{R}^N$

$$\max_N(x) = x_0 + h_\Delta(x - x_0\mathbf{1}). \tag{35}$$

Hence, to prove that $\max_N \in \operatorname{span} \mathcal{D}$ it suffices to show that h_Δ lies in the span of the restrictions of $\mathcal{D}$ to H_0. Indeed, $x_0 = \sigma(x_0) - \sigma(-x_0)$ lies in the span of the auxiliary functions (20), and by (21) the substitution $x \mapsto x - x_0\mathbf{1}$ fixes every dictionary function except the coordinate features $\sigma(\pm x_i)$, which become $\sigma(\pm(x_i - x_0))$; by (10) these are linear combinations of auxiliary functions. Hence $x \mapsto g(x - x_0\mathbf{1})$ belongs to $\operatorname{span} \mathcal{D}$ for every $g \in \mathcal{D}$, and (35) gives the claimed implication.

All polytopes in the geometric argument below lie in the $(N-1)$-dimensional space H_0. We restrict heads and edge functions to H_0 and identify them with their coefficient vectors there. Let $\mathcal{L}_k$ denote the real span of support functions of polytopes in H_0 of affine dimension at most k. In particular, $\mathcal{L}_0$ is the space of linear functions. For a direction

$$u \in H_0, \tag{36}$$

the exposed face of a polytope $P \subseteq H_0$ with normal u is the set of points of P maximizing the linear function $p \mapsto \langle p, u\rangle$:

$$P^u := \{p \in P : \langle p, u\rangle = h_P(u)\}. \tag{37}$$

The opposite face P^{-u} consists of the minimizers; for $u = 0$, both faces are P itself. Define the function

$$D_u h_P(x) := h_{P^u}(x) - h_{P^{-u}}(x) \qquad (x \in H_0). \tag{38}$$

Here u selects the two faces, while x is the input to their support functions. In particular, $D_0 = 0$. Geometrically, D_u acts as a derivative that replaces the support function of a polytope by the difference of the support functions of its opposite exposed faces. If $\langle u, \cdot\rangle$ is nonconstant on P, these are proper boundary faces of smaller dimension; if it is constant, the two support functions cancel. This removes the full-dimensional contribution and explains the dimension drop in (40). For a simplex and a facet normal u, D_u is a *facet-isolating operator*: it returns that facet's support function minus the linear support function of the opposite vertex. This is the mechanism used in Step 7.2.

For a finite linear combination g of support functions, define its one-sided directional derivative at u by

$$\begin{aligned} \partial_u g(x) &:= \lim_{t \downarrow 0} \frac{g(u + tx) - g(u)}{t}, \\ \partial_u h_P &= h_{P^u}, \qquad D_u g = \partial_u g - \partial_{-u} g. \end{aligned} \tag{39}$$

Thus the map $h_P \mapsto h_{P^u}$ extends linearly to linear combinations of support functions, independently of their decomposition. If $\dim P \le k$, either $\langle u, \cdot\rangle$ is constant on P, so $D_u h_P = 0$, or both selected faces have dimension at most $k - 1$. Consequently,

$$D_u \mathcal{L}_k \subseteq \mathcal{L}_{k-1} \qquad (k \ge 1). \tag{40}$$

The signed caterpillar sum. We seek a combination that agrees with h_Δ up to a function in $\mathcal{L}_{N-2}$. Adding $x_{\pi(0)}$ makes the caterpillar function the following maximum

$$x_{\pi(0)} + U_{\pi,S}(x) = \max\left\{ x_{\pi(0)},\, x_{\pi(q+1)} + \sum_{\ell \in L_{\pi,S}} \sigma(\ell(x)) \right\}. \tag{41}$$

Averaging over permutations gives the required symmetry

$$F_N(x) := \frac{1}{N!} \sum_{\pi} \sum_{\varnothing \ne S \subseteq \{1,\dots,r\}} (-1)^{N+|S|} \big(x_{\pi(0)} + U_{\pi,S}(x)\big), \tag{42}$$

where the signs $(-1)^{N+|S|}$ are chosen to cancel the extra contributions from support functions of exposed caterpillar faces, modulo $\mathcal{L}_{N-3}$. Establishing the required cancellations is a main step of the proof below.

Theorem 7 (Dictionary sufficiency, `Lean_9`, proof)**.** *For every $N \geq 3$, the dictionary of Definition 5 satisfies*

$$\max_N \in \operatorname{span} \mathcal{D}. \tag{43}$$

Proof of Theorem 7. Recall from this section's introduction that it suffices to show that h_Δ lies in the span of the restrictions of $\mathcal{D}$ to H_0. Define the error by

$$\begin{aligned} f : H_0 &\longrightarrow \mathbb{R}, \\ f(x) &:= F_N(x) - h_\Delta(x) \\ &= F_N(0, x_1, \ldots, x_{N-1}) - \max\{0, x_1, \ldots, x_{N-1}\}. \end{aligned} \tag{44}$$

Step 7.1 reduces dictionary membership to $f \in \mathcal{L}_{N-2}$. Step 7.2 shows that it suffices to prove $D_u f \in \mathcal{L}_{N-3}$ for every direction u. Step 7.3 proves these bounds by induction on N: dimension bounds and cancellation handle all directions except those with one coordinate different from all the others, which share a common value. In that case, deleting the exceptional coordinate invokes the induction hypothesis.

Step 7.1 (Reduce to a lower-dimensional error). In this step we reduce the problem to proving

$$f = F_N - h_\Delta \in \mathcal{L}_{N-2}. \tag{45}$$

Assume this relation temporarily, and write $Q_{\pi,S} := Q_{h_{\pi,S}, L_{\pi,S}}$, so that $U_{\pi,S} = h_{Q_{\pi,S}}$ by (19). Since the apex terms in (42) are linear, the assumed relation gives

$$\frac{1}{N!} \sum_{\pi} \sum_{\varnothing \neq S \subseteq \{1,\ldots,N-2\}} (-1)^{N+|S|} h_{Q_{\pi,S}} - h_\Delta \in \mathcal{L}_{N-2}. \tag{46}$$

We will correct this error by differentiating its support-function terms via ∂_u to lower the residual dimension until the residual is linear. The resulting support functions of proper faces of the $Q_{\pi,S}$ already lie in the dictionary span: the support function of the convex hull of the apex with a base face is a face function from Definition 4, given by (16), while the support function of a base face is a linear function plus inner hinges. Coordinate features represent the linear function, and (10) puts either orientation of each hinge in the span of (20). The apex alone has zero support function, and $h_{Q_{\pi,S}} = U_{\pi,S}$ is itself a dictionary function. Differentiating the simplex support-function term also produces support functions of proper simplex faces. Their membership in the dictionary span is supplied by induction on the number of vertices.

Claim 7.1 [Simplex faces, `Lean_10`, `Lean_11`]. For every nonempty $T \subseteq \{0, e_1, \ldots, e_{N-1}\}$, the support function of $\Delta_T := \operatorname{conv} T$ lies in the dictionary span.

Start ($|T| = 1$). The support function of a singleton is linear and lies in the dictionary span.

Step ($|T| \geq 2$). Fix T and assume that $h_{\Delta_A} \in \operatorname{span} \mathcal{D}$ for every nonempty vertex set A with $|A| < |T|$, in particular for every nonempty proper subset of T. Consider a current vertex set $A \subseteq \{0, e_1, \ldots, e_{N-1}\}$ with $|A| \geq 2$. To remove $v \in A$, choose a direction whose value at v is strictly larger than its common value at all other vertices. The positive exposed face is then the singleton $\{v\}$, and the negative exposed face is the simplex on the remaining vertices. Writing $\Delta_A := \operatorname{conv} A$, we have

$$\begin{aligned} u_v &:= \begin{cases} -\sum_{i=1}^{N-1} e_i & v = 0, \\ v & v \neq 0, \end{cases} \\ (\Delta_A)^{u_v} &= \{v\}, \qquad (\Delta_A)^{-u_v} = \Delta_{A \setminus \{v\}}, \\ D_{u_v} h_{\Delta_A}(x) &= \langle v, x \rangle - h_{\Delta_{A \setminus \{v\}}}(x). \end{aligned} \tag{47}$$

Choose an ordering $v_1, \ldots, v_{N-|T|}$ of the vertices outside T. Denote the error associated with the remaining vertex set T by

$$\varepsilon_T := D_{u_{v_{N-|T|}}} \cdots D_{u_{v_1}} \left(\frac{1}{N!} \sum_{\pi} \sum_{\varnothing \neq S \subseteq \{1,\ldots,N-2\}} (-1)^{N+|S|} h_{Q_{\pi,S}} - h_\Delta \right). \tag{48}$$

The operators act from right to left, removing v_1, then v_2, and so on. In (47), the support function of the remaining simplex carries a minus sign, while the removed singleton contributes the linear function $x \mapsto \langle v, x\rangle$. Every subsequent D_u annihilates linear functions. Thus the $N - |T|$ deletions give

$$D_{u_{v_{N-|T|}}} \cdots D_{u_{v_1}} h_\Delta = (-1)^{N-|T|} h_{\Delta_T} + \ell_T, \tag{49}$$

where ℓ_T is linear. The original error contains $-h_\Delta$, so the coefficient of h_{Δ_T} in ε_T is $-(-1)^{N-|T|} = (-1)^{N-|T|+1}$. All other terms are linear functions or support functions of faces of the $Q_{\pi,S}$, since each D_u replaces a support function by a difference of support functions of exposed faces. The coefficient of h_{Δ_T} in ε_T is ± 1, and all other terms already lie in the dictionary span. Thus proving $\varepsilon_T \in \operatorname{span} \mathcal{D}$ completes the induction step. To prove this membership, we use the additional information $\varepsilon_T \in \mathcal{L}_{|T|-2}$, obtained by repeated application of (40). This bound starts the inner induction below: each reduction lowers the dimension bound by one while subtracting a function in the dictionary span, until the remaining error lies in $\mathcal{L}_0$ and is therefore linear. For the full vertex set, the operator composition is empty and means the identity, so ε_T is exactly the error in (46).

For this fixed T, define the remaining errors as functions $\varepsilon_T^{(j)} : H_0 \to \mathbb{R}$ recursively by

$$\begin{aligned} \varepsilon_T^{(0)}(x) &:= \varepsilon_T(x), \\ \varepsilon_T^{(j+1)}(x) &:= \varepsilon_T^{(j)}(x) - \sum_{u \in E_{T,j}} \big(\partial_u \varepsilon_T^{(j)}\big)(x). \end{aligned} \tag{50}$$

Here $x \in H_0$ and $0 \le j < |T| - 2$. At stage j, the finite set of directions $E_{T,j} \subseteq H_0$ is chosen from the current error by the construction below. Thus $\varepsilon_T^{(j)}$ is the error left after j derivative sums have been subtracted from ε_T; the superscript counts stages and is not a power.

Claim 7.2 [Residual reduction, `Lean_12`]. The sets $E_{T,j}$ can be chosen successively so that, for $0 \le j \le |T| - 2$,

$$\begin{aligned} \varepsilon_T^{(j)} &\in \mathcal{L}_{|T|-2-j}, \\ \varepsilon_T - \varepsilon_T^{(j)} &\in \operatorname{span} \mathcal{D}. \end{aligned} \tag{51}$$

Each remaining error is a linear combination of h_{Δ_T}, support functions of proper faces of Δ_T, support functions of faces of the $Q_{\pi,S}$, and linear functions. The second condition says that the part already removed is represented by the dictionary.

Start ($j = 0$). By definition, $\varepsilon_T^{(0)} = \varepsilon_T$. Its dimension bound and support-function expression were established above, and the removed part is zero.

Step ($j < |T| - 2$). Assume the directions for the first j stages have been chosen and the claim holds at stage j. Put $k := |T| - 2 - j \ge 1$. We require each new direction u to be nonorthogonal to the simplex direction space $\operatorname{dir} \Delta_T := \operatorname{span}\{v - w : v, w \in T\}$. Its values on T are then not all equal, so the maximizing vertices form a nonempty proper subset:

$$\begin{aligned} A_u &:= \{v \in T : \langle u, v\rangle = h_{\Delta_T}(u)\}, \qquad \varnothing \ne A_u \subsetneq T, \\ \partial_u h_{\Delta_T} &= h_{\Delta_{A_u}} \in \operatorname{span} \mathcal{D}. \end{aligned} \tag{52}$$

The last membership uses the outer induction hypothesis, since $|A_u| < |T|$. An orthogonal direction would leave h_{Δ_T} unchanged and would not allow this use of the hypothesis.

Choose an auxiliary expression for $\varepsilon_T^{(j)}$ by support functions of polytopes of dimension at most k, and collect their distinct k-dimensional direction spaces. For each such space V, choose $u_V \in V^\perp$ not orthogonal to $\operatorname{dir} \Delta_T$ or to any other collected space V'. This is possible because $\dim \Delta_T = |T| - 1 > k$: each of $\operatorname{dir} \Delta_T$ and V' contains a vector outside V, whose nonzero projection onto $V^\perp$ is not orthogonal to that vector. Adding sufficiently small multiples of these projections avoids each orthogonality condition in turn while preserving those already avoided. Choose $E_{T,j} := \{u_V : V \text{ is a collected direction space}\}$, with $E_{T,j} = \varnothing$ if there are none. In the sum of derivatives over $E_{T,j}$, each term coming from a k-dimensional polytope is preserved exactly once. Every other

derivative of that term is the support function of a proper face, while terms of lower dimension remain in $\mathcal{L}_{k-1}$. Subtracting this derivative sum therefore cancels all k-dimensional terms. By the recursive definition (50), the resulting error satisfies $\varepsilon_T^{(j+1)} \in \mathcal{L}_{k-1} = \mathcal{L}_{|T|-3-j}$.

Now compute these same derivatives using the support-function expression in the claim. The derivative of h_{Δ_T} lies in the dictionary span by (52). Differentiating the support function of an already proper simplex face still gives a support function of a proper face of Δ_T, so the same induction hypothesis applies. Differentiating the support function of a face of $Q_{\pi,S}$ gives a support function of a face of the same polytope, whose support functions lie in the dictionary span, and linear functions remain linear. Thus the derivative sum lies in the dictionary span. By (50), $\varepsilon_T - \varepsilon_T^{(j+1)}$ is the sum of the previously removed part and this derivative sum, so it also lies in the dictionary span. Subtracting the derivative sum preserves the stated types of support-function terms. This completes the proof step.

At stage $j = |T| - 2$, the remaining error lies in $\mathcal{L}_0$ and is linear. Together with (51), this gives $\varepsilon_T \in \operatorname{span} \mathcal{D}$. Its vertex-deletion expression has coefficient $(-1)^{N-|T|+1} \in \{-1, 1\}$ on h_{Δ_T} and all other terms in the dictionary span, so $h_{\Delta_T} \in \operatorname{span} \mathcal{D}$. This completes the induction step on $|T|$. Taking T to be the full vertex set proves $h_\Delta \in \operatorname{span} \mathcal{D}$ under the temporary assumption (45). This finishes the proof of Step 7.1.

Step 7.2 (Why the facet bounds imply the class identity). For $N \geq 4$ and $f = F_N - h_\Delta$, we show that $f \in \mathcal{L}_{N-2}$ follows from

$$D_u f \in \mathcal{L}_{N-3} \qquad \text{for every } u \in H_0. \tag{53}$$

The idea is to express f modulo $\mathcal{L}_{N-2}$ using support functions of full-dimensional simplices. The facet bounds (53) then force their weighted contributions to cancel modulo $\mathcal{L}_{N-2}$.

The base of each $Q_{\pi,S}$ is an affine cube. Order its edge vectors by a bijection $\omega : \{1, \dots, N-2\} \to L_{\pi,S}$, add them one at a time, and adjoin the apex:

$$K_{\pi,S,\omega} := \operatorname{conv}\left(\{0\} \cup \left\{ h_{\pi,S} + \sum_{a=1}^{k} \omega(a) : 0 \leq k \leq N-2 \right\}\right). \tag{54}$$

These $(N-2)!$ simplices are obtained by coning the *Freudenthal–Kuhn triangulation* of the base to the apex (Freudenthal, 1942; Kuhn, 1960): in cube coordinates, each base simplex occupies one ordering region $1 \geq t_{\omega(1)} \geq \cdots \geq t_{\omega(N-2)} \geq 0$. For a hyperplane cut $P = P_+ \cup P_-$, the support-function identity

$$h_P + h_{P_+ \cap P_-} = h_{P_+} + h_{P_-} \tag{55}$$

shows that subdivision changes the sum of support functions only by support functions of the cutting sections. Hence

$$h_{Q_{\pi,S}} - \sum_{\omega} h_{K_{\pi,S,\omega}} \in \mathcal{L}_{N-2}. \tag{56}$$

Substituting (56) into (42) expresses f as a finite linear combination of support functions of $(N-1)$-dimensional simplices in H_0, plus a remainder in $\mathcal{L}_{N-2}$. The simplex Δ contributes its support function with coefficient -1, and the linear apex terms are included in the remainder.

For a simplex P, an outward facet normal ν_F exposes the facet F on one side and its opposite vertex v_F on the other. Thus

$$D_{\nu_F} h_P(x) = h_F(x) - \langle v_F, x \rangle. \tag{57}$$

Directions exposing no facet on either side give support functions of faces of dimension at most $N-3$. Consequently, (53) says that each weighted, signed sum of parallel-facet support functions belongs to $\mathcal{L}_{N-3}$.

Our assumption controls the weighted facet contributions, whereas our conclusion concerns f. To recover the original sum modulo $\mathcal{L}_{N-2}$, we use the same linear operator for every simplex, so cancellations between facet contributions survive reconstruction. Define

$$(\mathcal{T} g)(x) := \begin{cases} g(x) & x_1 \geq 0, \\ g(x - x_1 e_1) & x_1 < 0. \end{cases} \tag{58}$$

The sign in this definition concerns the input x. To interpret it geometrically, first suppose that every point $p \in P$ satisfies $p_1 \geq 0$, and write $\mathrm{pr}(p) := p - p_1 e_1$ for projection onto $p_1 = 0$. Since $\langle p, x\rangle - \langle \mathrm{pr}(p), x\rangle = p_1 x_1$, each point dominates its projection when $x_1 \geq 0$, and its projection dominates when $x_1 < 0$. Consequently,

$$\mathcal{T} h_P(x) = \max\{h_P(x), h_{\mathrm{pr}\,P}(x)\} = h_{\mathrm{conv}(P \cup \mathrm{pr}\,P)}(x). \tag{59}$$

This convex hull fills the segments with endpoints p and its projection, for $p \in P$, so its dimension is at most $\dim P + 1$. For an arbitrary P, choose $c \geq 0$ such that $P + ce_1$ lies in $p_1 \geq 0$. Translation adds cx_1 to the support function, and $\mathcal{T}(x_1) = \sigma(x_1)$, giving

$$\mathcal{T} h_P(x) = h_{\mathrm{conv}((P+ce_1) \cup \mathrm{pr}(P+ce_1))}(x) - c\sigma(x_1). \tag{60}$$

The correction $\sigma(x_1)$ is the support function of the segment $[0, e_1]$. Thus linearity gives

$$\mathcal{T}\mathcal{L}_k \subseteq \mathcal{L}_{k+1}. \tag{61}$$

Claim 7.3 [Simplex reconstruction, `Lean_13`]. Every $(N-1)$-dimensional simplex $P \subseteq H_0$ satisfies

$$h_P - \mathcal{T}\left(\sum_{F \text{ facet of } P} \mathrm{sgn}\langle \nu_F, e_1\rangle \, h_F\right) \in \mathcal{L}_{N-2}. \tag{62}$$

Here ν_F is an outward normal of F. Its sign against e_1 records whether the facet bounds lines parallel to e_1 from above or below; parallel facets contribute zero. Let $k = |\{F \text{ facet of } P : \langle \nu_F, e_1\rangle \neq 0\}|$ count facets not parallel to e_1. Boundedness ensures both nonzero signs occur, so $k \geq 2$.

Start ($k = 2$). There is one upper facet F^+ and one lower facet F^-. Every line parallel to e_1 that meets P has its two intersection endpoints on these facets. Thus $h_P(x) = h_{F^+}(x)$ when $x_1 \geq 0$, and $h_P(x) = h_{F^-}(x)$ when $x_1 < 0$. When $x_1 \geq 0$, $\mathcal{T}$ evaluates at x, so subtracting $\mathcal{T}(h_{F^+} - h_{F^-})$ from h_P leaves h_{F^-}. When $x_1 < 0$, $\mathcal{T}$ evaluates both facet support functions at $x - x_1 e_1$, where they agree because the facets have the same projection. Their difference is therefore zero, again leaving h_{F^-}. The reconstruction error is therefore $h_{F^-} \in \mathcal{L}_{N-2}$, proving the claim for $k = 2$.

Step ($k > 2$). Assume the claim for $k-1$, and let P have k facets not parallel to e_1. Two of them have the same nonzero sign of $\langle \nu_F, e_1\rangle$. Their intersection is an $(N-3)$-dimensional face C. Let $W := \mathrm{span}\{p - q : p, q \in C\}$, and project orthogonally within H_0 onto $W^\perp$. This is a two-dimensional plane, since $\dim H_0 = N-1$ and $\dim W = N-3$. All points of C have the same projection c, because their differences lie in W. There are just two vertices of P outside C, so the projected simplex is a triangle with vertices c and the projections of those two vertices. The chosen facets project to its two sides meeting at c.

The chosen facet normals are orthogonal to W, so their signs against e_1 equal their signs against its projection. This projected direction is nonzero, since the chosen facets are not parallel to e_1. If both sides are upper, moving from c opposite to this direction enters the triangle; if both are lower, moving in this direction enters it. Continuing meets the interior of the opposite edge. The inverse image of the line through c in this direction is a hyperplane in H_0 containing C and parallel to e_1. It cuts the edge whose endpoints are the two vertices outside C at an interior point and divides P into two $(N-1)$-dimensional simplices. Each keeps one of the two chosen facets and replaces the other by the new facet parallel to e_1. Its other facets retain their normal directions, so each new simplex has exactly $k-1$ facets not parallel to e_1.

Write $E(Q)$ for the reconstruction error on the left of (62). Let P_+ and P_- be the two new simplices, let $R := P_+ \cap P_-$ be their common cutting facet, and write F_1, F_2 for the chosen facets and $R_F := R \cap F$ for every other facet of P. With $s_F := \mathrm{sgn}\langle \nu_F, e_1\rangle$, applying (55) to P and to every split facet gives the exact residual identity

$$E(P) = E(P_+) + E(P_-) - h_R + \mathcal{T}\left(\sum_{\substack{F \text{ facet of } P \\ F \notin \{F_1, F_2\}}} s_F h_{R_F}\right). \tag{63}$$

Indeed, the inherited facets retain their signs, while the two copies of R are parallel to e_1 and therefore have sign zero. By the induction hypothesis, $E(P_+), E(P_-) \in \mathcal{L}_{N-2}$; moreover, $\dim R = N-2$ and $\dim R_F \leq N-3$. Equation (61) therefore puts every term on the right of (63) in $\mathcal{L}_{N-2}$. This proves the claim for k and completes the induction.

Conclusion for f. Apply (62) to every simplex support-function term in the expansion of f, keeping its original coefficient. A facet F parallel to e_1 has $\langle \nu_F, e_1 \rangle = 0$, so its term in the reconstruction formula is zero and can be omitted. For each remaining occurrence of a facet, choose ν_F outward from its simplex and put

$$u_F := \frac{\nu_F}{\langle \nu_F, e_1 \rangle}, \qquad \langle u_F, e_1 \rangle = 1. \tag{64}$$

Thus the coordinate with index 1 is normalized to 1; the coordinate with index 0 remains zero because $u_F \in H_0$. Oppositely oriented parallel normals give the same u_F. Only the normal is rescaled: the support function and its coefficient are unchanged.

Let $\mathcal{U}$ be the finite set of vectors u_F obtained this way. For each $u \in \mathcal{U}$, define G_u by adding all facet support-function terms with $u_F = u$, retaining their signs and weights. An occurrence of F as a facet of $K_{\pi,S,\omega}$ contributes $(-1)^{N+|S|} \operatorname{sgn}\langle \nu_F, e_1 \rangle \, h_F/N!$, while an occurrence as a facet of Δ contributes $-\operatorname{sgn}\langle \nu_F, e_1 \rangle \, h_F$. These are precisely the terms with the same normalized normal, including all repetitions.

More explicitly, define ρ to be the sum of the linear apex terms and the weighted subdivision errors in (56). Then

$$f = \frac{1}{N!} \sum_{\pi} \sum_{\varnothing \neq S \subseteq \{1,\dots,N-2\}} (-1)^{N+|S|} \sum_{\omega} h_{K_{\pi,S,\omega}} - h_\Delta + \rho, \qquad \rho \in \mathcal{L}_{N-2}. \tag{65}$$

For a simplex P, collect its facets with normalized normal u as

$$B_u(P) := \sum_{\substack{F \text{ facet of } P \\ \langle \nu_F, e_1 \rangle \neq 0,\ u_F = u}} \operatorname{sgn}\langle \nu_F, e_1 \rangle h_F. \tag{66}$$

An empty sum is zero. If $\langle \nu_F, e_1 \rangle > 0$, the direction u_F exposes F; if it is negative, $-u_F$ exposes F. The opposite face is a single vertex, so (57) gives the signed facet term up to a linear function. If neither u nor $-u$ exposes a facet of P, both exposed faces have dimension at most $N-3$. Thus $D_u h_P - B_u(P) \in \mathcal{L}_{N-3}$. Applying this to the full weighted expansion, with the same permutations and nonempty cut sets as in (42), gives

$$\begin{aligned} D_u f - G_u = \sum_{\pi, S \neq \varnothing} \frac{(-1)^{N+|S|}}{N!} \sum_{\omega} \big(D_u h_{K_{\pi,S,\omega}} - B_u(K_{\pi,S,\omega}) \big) \\ - \big(D_u h_\Delta - B_u(\Delta) \big) + D_u \rho \in \mathcal{L}_{N-3} \qquad (u \in \mathcal{U}). \end{aligned} \tag{67}$$

Together with (53), relation (67) puts every G_u in $\mathcal{L}_{N-3}$. By linearity, the sum of the reconstructed facet terms is $\mathcal{T}(\sum_{u \in \mathcal{U}} G_u)$, which lies in $\mathcal{L}_{N-2}$ by (61). Its difference from f consists of the reconstruction and subdivision remainders, also in $\mathcal{L}_{N-2}$. Hence $f \in \mathcal{L}_{N-2}$. This finishes the proof of Step 7.2.

Step 7.3 (Prove the class identity by induction).

Claim 7.4 [Class identity, `Lean_14`]. The class identity (45) holds for every $N \geq 3$.

Start ($N = 3$). Here $r = 1$, so the nonempty cut set S must be $\{1\}$. The tree has the single edge $\pi(1) \to \pi(2)$, with $q = 1$ and $p(1) = 0$. Equation (41) gives

$$\begin{aligned} x_{\pi(0)} + U_{\pi,\{1\}}(x) &= \max\big\{ x_{\pi(0)}, x_{\pi(2)} + \sigma(x_{\pi(1)} - x_{\pi(2)}) \big\} \\ &= \max\{ x_{\pi(0)}, x_{\pi(1)}, x_{\pi(2)} \} = \max_3(x). \end{aligned} \tag{68}$$

Each term has coefficient $(-1)^{3+1}/3! = 1/6$. For each apex label there are two orders of the remaining labels, and both give $\max_3$. Grouping the six positive terms by their apex gives

$$F_3(x) = \frac{1}{6} \left(\underbrace{2 \max_3(x)}_{\pi(0)=0} + \underbrace{2 \max_3(x)}_{\pi(0)=1} + \underbrace{2 \max_3(x)}_{\pi(0)=2} \right) = \max_3(x). \tag{69}$$

On H_0, $x_0 = 0$ and $\max_3(x) = \max\{0, x_1, x_2\} = h_\Delta(x)$. Hence $f = 0 \in \mathcal{L}_1$, proving (45) for $N = 3$.

Step ($N \geq 4$). Assume the class identity for $N-1$. By Step 7.2, it suffices to prove $D_u f \in \mathcal{L}_{N-3}$ for every $u \in H_0$. By (38), applying D_u replaces each support function h_P by $h_{P^u} - h_{P^{-u}}$: we select the maximizing and minimizing faces in direction u and subtract their support functions. The linear apex terms disappear. For Δ, these faces contain exactly the vertices with largest and smallest coordinate values u_i, including $u_0 = 0$. For a caterpillar base, the segment $[0, \ell_{ij}]$ remains free when $u_i = u_j$; its maximizing endpoint is ℓ_{ij} when $u_i > u_j$ and 0 when $u_i < u_j$. The minimizing face uses the opposite choices.

We discard support-function contributions already in $\mathcal{L}_{N-3}$ and cancel the remaining ones. Induction is needed only when one of $u_0, \ldots, u_{N-1}$ differs from the common value of all the others: the exposed simplex facet and the surviving caterpillar sum then give the same problem with $N-1$ inputs.

Discarding contributions in $\mathcal{L}_{N-3}$. Fix $u \in H_0 \setminus \{0\}$, since $D_0 f = 0$, and fix a caterpillar polytope $Q_{\pi,S}$ occurring in the signed sum. We first analyze its exposed faces. A face F of $Q_{\pi,S}$ lying entirely in the base zonotope $B_{\pi,S} := h_{\pi,S} + \sum_{\ell \in L_{\pi,S}} [0, \ell]$, and thus not containing the apex 0, is itself a translated sum of segments. Its support function is therefore a linear function plus a sum of segment support functions, and belongs to $\mathcal{L}_1 \subseteq \mathcal{L}_{N-3}$. If the exposed face is the singleton apex $\{0\}$, its support function is identically zero. If at least two tree edges have endpoints whose labels i, j satisfy $u_i \neq u_j$, at most $N-4$ segments remain free, so even adjoining the apex gives dimension at most $N-3$. Thus we need only faces containing the apex and an exposed base face, with at most one tree edge whose endpoint labels i, j satisfy $u_i \neq u_j$.

We consider an exposed face in either direction u or $-u$. For the latter, replace u by $-u$, so that in both cases we analyze the face on which $\langle u, \cdot \rangle$ is maximized. Let $a := u_{\pi(0)}$ be the coordinate value at the apex label. The exposed face contains both the apex 0 and points of the base precisely when the maximum of $\langle u, p \rangle$ over the base is zero. There are three cases.

Case 1: one spine edge with unequal endpoint values. Removing this edge splits the tree into two components with constant coordinate values $c \neq d$. The maximum over the base is $\max\{c, d\} - a$, so the required condition is

$$a = \max\{c, d\}. \tag{70}$$

The apex label therefore adds no new coordinate value, and u has exactly two distinct coordinate values.

Case 2: one leaf edge with unequal endpoint values. Let d be the leaf's coordinate value and $c \neq d$ the common value at all other tree vertices. The maximum over the base is $c + \sigma(c-d) - a$, so the required condition is

$$a = c + \sigma(c-d) = \begin{cases} c, & d > c, \\ 2c - d, & d < c. \end{cases} \tag{71}$$

If $d > c$, the apex label has value c, giving exactly two distinct coordinate values. If $d < c$, put $t := c - d > 0$. The leaf has value $c - t$, the apex label has value $c + t$, and all other $N-2$ labels have value c. Up to a permutation of labels, this gives the three-value pattern $(c+t, c-t, c, \ldots, c)$.

Case 3: equal values at every tree-edge endpoint pair. If $u_i = u_j$ for the endpoint labels of every tree edge, all tree values equal some c. The maximum over the base is $c - a$, so we would also need $a = c$, making every coordinate of u equal. Since $u \in H_0$, this would force $u = 0$, which we excluded.

Replacing u by $-u$ preserves the two types of pattern from Cases 1 and 2. Therefore every remaining contribution has either exactly two distinct coordinate values or the three-value pattern $(c+t, c-t, c, \ldots, c)$; all other directions have been excluded. Since $Q_{\pi,S}$ was arbitrary, this classification applies to every summand. We now consider their contributions to the full signed sum.

Cancellation for three distinct values. Let α, β be the exceptional labels, with $u_\alpha = c + t$, $u_\beta = c - t$, and $u_i = c$ for every other label, where $t \neq 0$. Let τ exchange α and β and fix all other labels. Define the linear map $R : H_0 \to H_0$ explicitly by

$$(Rx)_i := x_{\tau(i)} - x_{\tau(0)} \qquad (0 \leq i < N,\ x \in H_0). \tag{72}$$

Thus R first swaps the two exceptional coordinates and then subtracts the new zeroth coordinate from every coordinate; in particular, $(Rx)_0 = 0$. Since $u_0 = 0$, this formula also gives $Rx = x - \frac{x_\alpha - x_\beta}{t}u$. With the explicit linear functional $\langle b, x\rangle := (x_\alpha - x_\beta)/(2t)$, we therefore have $Rx = x - 2\langle b, x\rangle u$, $\langle b, u\rangle = 1$, and $Ru = -u$. Hence R is a rank-one reflection. Permutation symmetry and the common coordinate shift $-x_{\tau(0)}$ give $F_N(Rx) = F_N(x) + \ell(x)$ for a linear function ℓ. Write $G := F_N|_{H_0}$. Applying the reflection identity to $u + \varepsilon x$ and to u, subtracting, dividing by $\varepsilon > 0$, and taking $\varepsilon \downarrow 0$ gives $\partial_{-u}G(Rx) = \partial_u G(x) + \ell(x)$, because $R(u + \varepsilon x) = -u + \varepsilon Rx$. To remove R from the derivative argument, positive homogeneity gives the identity $\partial_v G(x + \lambda v) = \partial_v G(x) + \lambda G(v)$ for every real λ. Indeed, for sufficiently small $\varepsilon > 0$, write $v + \varepsilon(x + \lambda v) = (1 + \varepsilon\lambda)\big(v + \frac{\varepsilon}{1+\varepsilon\lambda}x\big)$ and apply the definition (39). Using this identity with $v = -u$ and $\lambda = 2\langle b, x\rangle$, we obtain

$$\begin{aligned} \partial_u G(x) + \ell(x) &= \partial_{-u}G(Rx) \\ &= \partial_{-u}G\big(x + 2\langle b, x\rangle(-u)\big) \\ &= \partial_{-u}G(x) + 2\langle b, x\rangle G(-u), \\ D_u G(x) &= \partial_u G(x) - \partial_{-u}G(x) = 2F_N(-u)\langle b, x\rangle - \ell(x). \end{aligned} \tag{73}$$

The right-hand side is linear in x, so

$$D_u(F_N|_{H_0}) \in \mathcal{L}_0. \tag{74}$$

The two exposed simplex faces are vertices, so their support functions are linear as well.

Handling two distinct values. First suppose that each value occurs at least twice. After the preceding dimension bound, every term still requiring cancellation has a unique split $u_{\pi(1)} = \cdots = u_{\pi(k)} \neq u_{\pi(k+1)} = \cdots = u_{\pi(r+1)}$ with $k \in S$. If the apex label has the latter value, toggle membership of $k - 1$ in S. Otherwise, exchange the equal-valued labels $\pi(k + 1), \pi(k + 2)$ and toggle membership of $k + 1$. Repetition of both values gives $k \geq 2$ in the first case and $k + 2 \leq r + 1$ in the second, so the operations are defined; the unchanged cut at k keeps the cut set nonempty. The transformed term remains in the same case, and repeating the operation returns the original term. If Φ denotes this involution on the active terms and $w_{\pi,S} := (-1)^{N+|S|}/N!$, then

$$w_{\Phi(\pi,S)} = -w_{\pi,S}, \qquad D_u h_{Q_{\pi,S}} - D_u h_{Q_{\Phi(\pi,S)}} \in \mathcal{L}_1. \tag{75}$$

Inactive fixed points of the global transformation are already in $\mathcal{L}_{N-3}$ by the dimension bound. Summing the active pairs therefore puts $D_u F_N$ in $\mathcal{L}_{N-3}$. Each exposed simplex face has at most $N - 2$ vertices and hence dimension at most $N - 3$, so $D_u f \in \mathcal{L}_{N-3}$ without using induction.

One exceptional coordinate: the induction hypothesis. We are still working in H_0, so $u_0 = x_0 = 0$. It remains to consider two values, one occurring only once. Choose a permutation τ such that $\tau(0)$ is the exceptional label. Relabeling must be followed by subtracting the new zeroth coordinate to stay in H_0:

$$B_\tau : H_0 \longrightarrow H_0, \qquad (B_\tau x)_i := x_{\tau(i)} - x_{\tau(0)} \qquad (0 \leq i < N). \tag{76}$$

Its zeroth coordinate is zero, and its inverse is $B_{\tau^{-1}}$. It sends u to $(0, t, \ldots, t)$ with $t \neq 0$. This change preserves every $\mathcal{L}_k$ and changes f only by a linear term, annihilated by D; we transform both the input and the direction. Rescaling the direction, and reversing the sign of D_u if necessary, reduces to $u = (0, 1, \ldots, 1)$ within H_0. It remains to show that $D_u f \in \mathcal{L}_0$.

This direction exposes the $(N - 2)$-dimensional facet $\Delta' := \operatorname{conv}\{e_1, \ldots, e_{N-1}\}$ and the opposite vertex 0. Hence

$$D_u h_\Delta(x) = h_{\Delta'}(x) = \max\{x_1, \ldots, x_{N-1}\} \qquad (x \in H_0). \tag{77}$$

Here is the exact use of the induction hypothesis. Set $z := (x_1, \ldots, x_{N-1})$ and $y := z - x_1\mathbf{1} = (0, x_2 - x_1, \ldots, x_{N-1} - x_1)$, where $\mathbf{1} \in \mathbb{R}^{N-1}$ has every coordinate equal to 1. Every caterpillar maximum increases by a common added constant. Averaging over the $(N - 1)!$ lower label orders cancels the factorial in the total signed coefficient, giving

$$F_{N-1}(z) - \max_{N-1}(z) = F_{N-1}(y) - \max_{N-1}(y) + (W_{N-1} - 1)x_1 \tag{78}$$

for

$$W_{N-1} := \sum_{\varnothing \neq T \subseteq \{1,\ldots,N-3\}} (-1)^{N-1+|T|}. \tag{79}$$

The remaining $N-1$ inputs z need not start with zero, but y lies in the lower-arity slice $\{y \in \mathbb{R}^{N-1} : y_0 = 0\}$, exactly the domain of the induction hypothesis. Apply the $(N-1)$-input class identity to y on the right. Linear precomposition preserves $\mathcal{L}_{N-3}$, and the last term is linear. Since $(N-1)-2 = N-3$, this gives

$$\big(x \mapsto F_{N-1}(x_1, \dots, x_{N-1}) - \max\{x_1, \dots, x_{N-1}\}\big) \in \mathcal{L}_{N-3}. \tag{80}$$

The remaining task is to match the exposed caterpillar support functions with the lower-arity sum.

Claim 7.5 [Coordinate deletion, `Lean_15`]. For $N \geq 4$,

$$D_{(0,1,\dots,1)}(F_N|_{H_0}) - \big(x \mapsto F_{N-1}(x_1, \dots, x_{N-1})\big) \in \mathcal{L}_{N-3}. \tag{81}$$

The proof below checks the surviving terms and their coefficients. Readers may skip it and continue with the completion of the induction.

Proof of Claim 7.5. We now determine which terms of the signed sum must be retained for $u = (0, 1, \dots, 1)$. Only label 0 has a different coordinate value from the others. Consequently, a tree-edge generator is nonorthogonal to u exactly when one of its endpoint labels is 0.

If label 0 is the apex label, all tree-edge generators are orthogonal to u. The maximizing face is the whole base $B_{\pi,S}$ and the minimizing face is $\{0\}$, so $D_u h_{Q_{\pi,S}} = h_{B_{\pi,S}} \in \mathcal{L}_1 \subseteq \mathcal{L}_{N-3}$. If label 0 is a tree vertex of degree at least two, at least two segment generators are nonorthogonal to u. The preceding dimension bound then gives $D_u h_{Q_{\pi,S}} \in \mathcal{L}_{N-3}$ as well. We can therefore discard both kinds of term: it suffices to retain descriptions in which label 0 is a tree vertex of degree one.

To express this condition in the permutation notation, let $p := \pi^{-1}(0)$ be the index of label 0 in the full order $(\pi(0), \dots, \pi(N-1))$. When $p \geq 1$, its tree position is $p-1$. A degree-one vertex is either an endpoint of the spine with no attached leaves or a leaf outside the spine. The retained descriptions are thus exactly

$$\begin{aligned} &p = 1, \quad 1 \in S && \text{(root endpoint)},\\ &2 \leq p \leq N-1, \quad p-1 \notin S && \text{(leaf outside the spine)},\\ &p = N-1, \quad N-2 \in S && \text{(last spine endpoint)}. \end{aligned} \tag{82}$$

Indeed, the root has only one child precisely when $1 \in S$; a nonroot spine vertex has a parent and also a child unless it occupies the final tree position. Deleting label 0 from the full order now gives an order $\bar{\pi}$ mapping positions $\{0, \dots, N-2\}$ bijectively to labels $\{1, \dots, N-1\}$.

For $0 \leq i \leq N-3$, let ι_i insert a gap at the prospective pendant position $i+1$. Given a lower cut set $T \subseteq \{1, \dots, N-3\}$, the three active insertion rules are

$$\begin{aligned} \iota_i(t) &:= \begin{cases} t, & t < i+1,\\ t+1, & t \geq i+1, \end{cases}\\ \text{first end:} \quad & (p, S) = \big(1, \{1\} \cup \{t+1 : t \in T\}\big),\\ \text{pendant } i\text{:} \quad & (p, S) = \big(i+2, \{\iota_i(t) : t \in T\}\big) \qquad (0 \leq i \leq N-3),\\ \text{last end:} \quad & (p, S) = \big(N-1, T \cup \{N-2\}\big). \end{aligned} \tag{83}$$

In every row the upper permutation is uniquely obtained by inserting label 0 at position p in $\bar{\pi}$. Conversely, with t restricted to $\{1, \dots, N-3\}$, the deletion rules are $T = \{t : t+1 \in S\}$ for the first end, $T = \{t : \iota_i(t) \in S\}$ for pendant i, and $T = S \setminus \{N-2\}$ for the last end. These operations are inverse. The three cases in (82) correspond respectively to the first-end row, the pendant row with $i = p-2$, and the last-end row. Thus a nonempty T has $N-2$ pendant and two endpoint preimages, whereas $T = \varnothing$ has only the two endpoints because upper cut sets are nonempty.

Extend the lower caterpillar notation to $T = \varnothing$ by setting its last spine position and every parent equal to 0, so its head begins at the root $\bar{\pi}(1)$. For every lower label order $\bar{\pi}$ put

$$\begin{aligned} C_{\bar{\pi},T}(x) &:= x_{\bar{\pi}(0)} + U_{\bar{\pi},T}(x),\\ q_T &:= \max(\{0\} \cup T), \qquad p_T(j) := \max(\{0\} \cup \{t \in T : t < j\}). \end{aligned} \tag{84}$$

For each upper preimage (π, S) of $(\bar{\pi}, T)$, the insertion identities and the literal factorial coefficients are

$$\begin{aligned} \text{pendant:} \quad & \frac{(-1)^{N+|S|}}{N!} = -\frac{1}{N}\frac{(-1)^{N-1+|T|}}{(N-1)!}, \quad D_u\big(x_{\pi(0)} + U_{\pi,S}\big) + C_{\bar{\pi},T} \in \mathcal{L}_1, \\ \text{endpoint:} \quad & \frac{(-1)^{N+|S|}}{N!} = \frac{1}{N}\frac{(-1)^{N-1+|T|}}{(N-1)!}, \qquad D_u\big(x_{\pi(0)} + U_{\pi,S}\big) - C_{\bar{\pi},T} \in \mathcal{L}_1. \end{aligned} \tag{85}$$

Thus every one of the $N - 2 + 2 = N$ preimages contributes $1/N$ of the lower summand modulo $\mathcal{L}_{N-3}$, and together they recover its coefficient in F_{N-1}.

The extra terms are the two endpoint fibres whose deletion leaves $T = \varnothing$, hence a rooted star. By (85), each contributes $1/N$ of the weighted lower summand modulo $\mathcal{L}_1$. For a fixed lower order $\bar{\pi}$, the nonroot positions are $J = \{0, 2, \ldots, N-2\}$, including the original apex position 0. For $a \in J$, let $\bar{\pi}^{(a)}$ exchange positions 0 and a, keeping the root at position 1; the exchange at $a = 0$ is the identity. Put

$$\begin{aligned} q := |J| = N - 2, \qquad & d_a := x_{\bar{\pi}(1)} - x_{\bar{\pi}(a)} \quad (a \in J), \\ T_A := \sigma\left(\sum_{a\in A} d_a + \sum_{a\in J\setminus A} \sigma(d_a)\right) \quad & (A \subseteq J), \\ C_{\bar{\pi}^{(a)},\varnothing} = x_{\bar{\pi}(a)} + T_{\{a\}} \quad & (a \in J). \end{aligned} \tag{86}$$

The Boolean alternating identity gives

$$\sum_{A\subseteq J} (-1)^{|A|} T_A = 0 \quad \Longrightarrow \quad \sum_{a\in J} T_{\{a\}} = T_\varnothing + \sum_{\substack{A\subseteq J \\ |A|\geq 2}} (-1)^{|A|} T_A. \tag{87}$$

The empty term is a sum of support functions of segments. For $|A| \geq 2$, the polytope with support function T_A has one head direction and $q - |A|$ segment directions, hence dimension at most $1 + q - |A| \leq q - 1 = N - 3$. Thus the singleton sum lies in $\mathcal{L}_{N-3}$; the sum of the linear apex coordinates lies in $\mathcal{L}_0$. Each fixed-position exchange is a bijection on lower label orders, so

$$q \sum_{\bar{\pi}} C_{\bar{\pi},\varnothing} = \sum_{\bar{\pi}} \sum_{a\in J} C_{\bar{\pi}^{(a)},\varnothing} \in \mathcal{L}_{q-1}. \tag{88}$$

Since $q > 0$, the full empty-cut sum belongs to $\mathcal{L}_{q-1} = \mathcal{L}_{N-3}$. The empty-cut coefficient $(-1)^{N-1}/(N-1)!$ and the factor $2/N$ from the two endpoint fibres are scalars, so the extra star sum lies there as well. This proves Claim 7.5. □

Completion of the induction. Combining (77), (80), and Claim 7.5 gives

$$\begin{aligned} D_u f(x) = \big[D_u(F_N|_{H_0})(x) - F_{N-1}(x_1, \ldots, x_{N-1})\big] \\ + \big[F_{N-1}(x_1, \ldots, x_{N-1}) - \max\{x_1, \ldots, x_{N-1}\}\big]. \end{aligned} \tag{89}$$

The first bracket belongs to $\mathcal{L}_{N-3}$ by coordinate deletion Claim 7.5, the second by the induction hypothesis. This proves the bound for every direction. Step 7.2 gives $f \in \mathcal{L}_{N-2}$, completing the induction, and Step 7.1 gives the dictionary representation. This finishes the proof of Theorem 7. □

4 NETWORK SIZE

The widths in Theorem 2 are worst-case upper bounds for this construction. To count second hidden layer units, we absorb linear edges into the head, orient each remaining hinge as $x_i - x_j$ with $i < j$, and merge descriptions with the same adjusted head and hinge set. We count all possible resulting descriptions, without relying on any coefficient being zero. We combine the contributions $x \mapsto \sigma(x_i - x_j)$ with the auxiliary terms and represent their sum using two additional second hidden layer neurons. The first hidden layer width comes from a fixed set of shared features sufficient for every such description. Discarding zero aggregate coefficients can yield a smaller network; zero padding gives the stated widths. We first count these descriptions, then construct the shared first hidden layer and account for both widths.

Proof of Theorem 2. Fix $N \geq 3$. Theorem 7 and Lemma 6, followed by (33), give the exact tagged expansion of $\max_N$.

Step 2.1 (Count the caterpillar face functions). We first rewrite each face function using only its adjusted head and the hinges that remain. This lets us recognize when different face descriptions require the same second hidden layer neuron. An edge kept linearly is added to the head, an omitted edge disappears, and an edge kept as a hinge stays in the set M. Thus

$$U^{\tau}_{h,L} = U_{h',M}, \qquad h' := h + \sum_{\tau(\ell)=\ell} \ell, \qquad M := \{\ell \in L : \tau(\ell) = \sigma \circ \ell\}. \tag{90}$$

The resulting function description is (h', M) in the sense of Definition 4: h' is the adjusted head, and M contains precisely the retained hinges. When two resulting pairs have the same adjusted head and the same hinge set, they can use one second hidden layer neuron, with their output coefficients added. Distinct pairs may still represent the same function, so counting these pairs gives an upper bound. The shared first hidden layer is constructed in the next step. We next orient the remaining hinges consistently. By (10), applied to each $\ell_{ij} \in M$ with $i > j$, the same face function $U_{h',M}$ has the *normalized description* (h^+, M^+), where

$$\begin{aligned} M^+ &:= \{\ell_{\min(i,j),\max(i,j)} : \ell_{ij} \in M\}, \\ h^+ &:= h' + \sum_{\ell_{ij}\in M,\ i>j} \ell_{ij}. \end{aligned} \tag{91}$$

Every remaining hinge now has orientation $x_i - x_j$ with $i < j$, and the additional linear terms have been absorbed into h^+. No edge function of a caterpillar is the negative of another one, so M and M^+ have the same size. Geometrically, $[0, \ell] = \ell + [0, -\ell]$: reversing a generator requires shifting the head by the old generator. Applying this to every reversed edge leaves the translated base, and hence $Q_{h',M} = Q_{h^+,M^+}$, exactly unchanged. Normalization chooses a consistent orientation and a corresponding base corner; it does not rescale vectors or change the shape (Figure 2).

For a caterpillar description (π, S), the spine endpoints are labelled $\pi(1)$ and $\pi(q+1)$, and every label other than $\pi(1)$ and the apex is the child of exactly one edge. Hence the normalized description of $U_{\pi,S}$ is

$$\begin{aligned} h^+_{\pi,S}(x) &= \sum_{\ell_{ij}\in L_{\pi,S}} x_{\max(i,j)} + x_{\pi(1)} + x_{\pi(q+1)} - \sum_{i=0}^{N-1} x_i, \\ L^+_{\pi,S} &= \{\ell_{\min(i,j),\max(i,j)} : \ell_{ij} \in L_{\pi,S}\}. \end{aligned} \tag{92}$$

It depends only on the undirected tree, the apex, and the unordered pair of spine endpoints, not on the order of the leaves at a spine vertex or on the orientation of the spine. For a spine with m vertices, choose its vertex set in $\binom{N}{m}$ ways, its order up to reversal in $m!/2$ ways, its apex in $N - m$ ways, and the attachment of each of its $N - m - 1$ leaves in m ways. Reversal has no fixed point since $m \geq 2$. Every caterpillar has such data, so by (6)

$$\left|\{(h^+_{\pi,S}, L^+_{\pi,S}) : (\pi, S) \text{ a caterpillar description}\}\right| \leq C_N. \tag{93}$$

Reversing an edge function swaps its omitted and linear states, so normalizing before or after choosing the states gives the same descriptions. Every face tag therefore has a normalized description obtained from $(h^+_{\pi,S}, L^+_{\pi,S})$ by choosing one of three states for each of its $N - 2$ edge functions. Let $\mathcal{K}_0$ be the set of these normalized descriptions and let $\nu : \mathcal{A}_{\text{face}} \to \mathcal{K}_0$ be normalization. For $k = (h, M) \in \mathcal{K}_0$, write $U_k := U_{h,M}$. With (93), $|\mathcal{K}_0| \leq 3^{N-2} C_N$. Only equal normalized descriptions are merged; distinct members of $\mathcal{K}_0$ are not identified merely because they represent the same function.

Among them are the descriptions $(\ell_{ij}, \varnothing)$ representing $x \mapsto \sigma(x_i - x_j)$, with $i \neq j$: for the all-cut path $S = \{1, \dots, r\}$ with apex $\pi(0) = j$ and last spine vertex $\pi(r+1) = i$, omitting every edge function gives

$$U^0_{\pi,S} = \sigma \circ \ell_{ij} \qquad (0 \leq i \neq j < N). \tag{94}$$

The linear heads ℓ_{ij} are distinct for distinct ordered pairs. The corresponding functions $x \mapsto \sigma(x_i - x_j)$ are distinct as well: equality of two such ReLUs makes their positive half-spaces equal, so

their heads are positive scalar multiples, and equality on that half-space makes the scalar one. Let $\mathcal{P}$ be the subset of $\mathcal{K}_0$ consisting of every normalized description with empty hinge set and head equal to a coordinate difference; we combine their contributions with the auxiliary terms. The descriptions above show that $|\mathcal{P}| \geq N(N-1)$. Aggregate the tagged coefficients only along equal normalized face descriptions:

$$d_k := \sum_{\substack{a \in \mathcal{A}_{\mathrm{face}} \\ \nu(a)=k}} \widetilde{c}_a. \tag{95}$$

Discard zero fiber sums, then remove the absorbable keys:

$$\mathcal{K}_{\mathrm{act}} := \{k \in \mathcal{K}_0 : d_k \neq 0\}. \tag{96}$$

$$K := \mathcal{K}_{\mathrm{act}} \setminus \mathcal{P}. \tag{97}$$

By (10), every unit indexed by $\mathcal{P}$ lies in the span of the auxiliary functions and needs no separate second hidden layer neuron. Since $K \subseteq \mathcal{K}_0 \setminus \mathcal{P}$,

$$|K| \leq 3^{N-2} C_N - N(N-1). \tag{98}$$

This finishes the proof of Step 2.1.

Step 2.2 (Compile the network and establish its widths). The terms not assigned a retained neuron form the explicitly known remainder

$$R := \sum_{a \in \mathcal{A}_{\mathrm{aux}}} \widetilde{c}_a v(a) + \sum_{k \in \mathcal{K}_{\mathrm{act}} \cap \mathcal{P}} d_k U_k. \tag{99}$$

By (33) and (95)–(97), $\max_N = \sum_{k \in K} d_k U_k + R$. Expand the absorbed functions $x \mapsto \sigma(x_i - x_j)$ using (10) to write R as a linear combination of the auxiliary functions (20). The first hidden layer outputs the vector

$$\Phi(x) := \big(\sigma(x_0),\ \sigma(-x_0),\ (\sigma(x_i - x_0))_{i>0},\ (\sigma(x_i - x_j))_{i<j}\big). \tag{100}$$

Thus W_1 has rows $e_0^\top$ and $-e_0^\top$, then $e_i^\top - e_0^\top$ for $i > 0$, then $e_i^\top - e_j^\top$ for $i < j$, using a fixed ordering within each block. Its row count is

$$\binom{N}{2} + (N-1) + 2 = \binom{N}{2} + N + 1 = h_1. \tag{101}$$

By (10), these features recover all coordinates,

$$\begin{aligned} x_0 &= \sigma(x_0) - \sigma(-x_0), \\ x_i &= x_0 + \sigma(x_i - x_0) - \sigma(x_0 - x_i) \qquad (i > 0). \end{aligned} \tag{102}$$

For each $k = (h, M) \in K$, substitute (102) into $h(x) + \sum_{\ell \in M} \sigma(\ell(x))$ and collect the coefficients of $\Phi(x)$ into a row vector $w_k^\top$. Every hinge in M already has the orientation $i < j$, so its coefficient is added directly to the corresponding feature.

To specify the remaining row, collect the auxiliary coefficients of R as

$$R(x) = \sum_i p_i \sigma(x_i) + \sum_i n_i \sigma(-x_i) + \sum_{i<j} s_{ij} \sigma(x_i - x_j). \tag{103}$$

Each retained unit is translation invariant by (21), so

$$R(x + t\mathbf{1}) = R(x) + t, \qquad \sum_i p_i = 1. \tag{104}$$

The coefficient sum follows by setting $x = 0$ and $t = 1$. Now choose t large enough that every $x_i + t$ is nonnegative. Substituting the auxiliary expansion into (104) cancels t and gives

$$R(x) = \sum_i p_i x_i + \sum_{i<j} s_{ij} \sigma(x_i - x_j). \tag{105}$$

Substitute (102) into this expression and collect the coefficients of $\Phi(x)$ into w_R. Thus both w_k and w_R are specified by finite sums of the previously defined coefficients.

For any fixed ordering of K, the Caterpillar Net has weight matrix W_2 for the second hidden layer and weight vector w_3 for the output layer:

$$W_2 = \begin{pmatrix} (w_k^\top)_{k\in K} \\ w_R^\top \\ -w_R^\top \end{pmatrix}, \qquad w_3 = \begin{pmatrix} (d_k)_{k\in K} \\ 1 \\ -1 \end{pmatrix}, \qquad b_1 = b_2 = b_3 = 0. \tag{106}$$

In the second hidden layer, the first $|K|$ neurons output $U_k(x)$, and the last two output $\sigma(R(x))$ and $\sigma(-R(x))$. The output layer therefore computes

$$w_3^\top \sigma(W_2\Phi(x)) = \sum_{k\in K} d_k U_k(x) + \sigma(R(x)) - \sigma(-R(x)) = \max_N(x). \tag{107}$$

By (98), the second hidden layer width is at most

$$|K| + 2 \le 3^{N-2}C_N - N(N-1) + 2 = h_2. \tag{108}$$

Zero padding realizes the stated widths exactly.

For the growth rates, count the ordered choices of m spine labels:

$$\binom{N}{m}\frac{m!}{2} = \frac{N!}{2(N-m)!} \le N^m \qquad (2 \le m \le N-1). \tag{109}$$

Using $N - m \le N$ and $m^{N-m-1} \le N^{N-m-1}$ in each summand of (6) gives

$$\begin{aligned} C_N &\le \sum_{m=2}^{N-1} N^m \, N \, N^{N-m-1} = (N-2)N^N \\ &\le N^{N+1} \le N^{4N}. \end{aligned} \tag{110}$$

Thus no Stirling estimate is needed. Since $3^{N-2} \le N^N$ for $N \ge 3$, the widths satisfy

$$h_2 \le 3^{N-2}(N-2)N^N + 2 \le 2N^{5N}, \tag{111}$$

$$h_1 \le 4N^2. \tag{112}$$

Taking logarithms gives (8). This finishes the proof of Theorem 2. □

Formalization (`Lean_16`) Linked definitions, lemmas, and theorems are audited against exact types; only `propext`, `Classical.choice`, and `Quot.sound` are allowed. The repository contains the maximum construction and its proof. The general CPWL results in the introduction are cited background results; the equation links in the Wang–Sun theorem lead to its published source. Claude and Codex assisted with formalization and manuscript preparation.

5 Conclusion

The first hidden layer width is asymptotically sharp. Building on the extremal-graph approach of Safran (2026), our separate Lean development extends the bound $h_1 \ge \binom{N}{2} - \lfloor 4N/3 \rfloor$ of Lu (2026, Theorem 2.13), for $N \ge 4$, to networks with arbitrary affine skip connections. When every first hidden layer weight vector is proportional to some $e_i - e_j$, we sharpen it to $h_1 \ge \binom{N}{2} - N$. We obtain no growing lower bound for h_2; finding smaller exact maximum representations and establishing such lower bounds remain open.

References

Egor Bakaev, Florestan Brunck, Christoph Hertrich, Jack Stade, and Amir Yehudayoff. Better neural network expressivity: Subdividing the simplex. In *Proceedings of the 58th Annual ACM Symposium on Theory of Computing*, pages 500–507, 2026. doi:10.1145/3798129.3800768.

Florestan Brunck, Jack Stade, and Amir Yehudayoff. *Computing* $\max_n$ *in Depth 2—Proof Overview*. Unpublished draft supplied to the author, October 2026.

Hans Freudenthal. Simplizialzerlegungen von beschränkter Flachheit. *Annals of Mathematics*, 43(3):580–582, 1942. doi:10.2307/1968813.

Moritz Grillo, Christoph Hertrich, and Georg Loho. Depth-bounds for neural networks via the braid arrangement. *Advances in Neural Information Processing Systems*, 38:127800–127818, 2025. doi:10.52202/085713-4257.

Christoph Hertrich, Amitabh Basu, Marco Di Summa, and Martin Skutella. Towards lower bounds on the depth of ReLU neural networks. *SIAM Journal on Discrete Mathematics*, 37(2):997–1029, 2023. doi:10.1137/22M1489332.

Harold W. Kuhn. Some combinatorial lemmas in topology. *IBM Journal of Research and Development*, 4(5):518–524, 1960. doi:10.1147/rd.45.0518.

Shunqi Lu. Sharp closure thresholds for two-hidden-layer ReLU networks. Preprint, version 1, September 2026. doi:10.5281/zenodo.22728502. Author's PDF.

Itay Safran. A depth hierarchy for computing the maximum in ReLU networks via extremal graph theory. In *Proceedings of the 39th Conference on Learning Theory*, volume 336 of *Proceedings of Machine Learning Research*, pages 5721–5742, 2026. PMLR 336.

Shuning Wang and Xusheng Sun. Generalization of hinging hyperplanes. *IEEE Transactions on Information Theory*, 51(12):4425–4431, 2005. doi:10.1109/TIT.2005.859246.