

Solved Problems – Baye's Theorem

Posted on August 1, 2017 / Under Probability & Statistics

Q1. There are three bags. First bag contains 1 white, 2 red and 3 green balls. Second bag contains 2 white, 3 red and 1 green balls. Third bag contains 3 white, 1 red and 2 green balls.

A bag is chosen at random and 2 balls are drawn from it. These are found to be 1 white and 1 red. Find the probability that the balls so drawn came from the second bag.

Solution:

Let A_1 be the event of choosing first bag

Let A_2 be the event of choosing second bag

Let A_3 be the event of choosing third bag

Let E be the event of drawing 1 white and 1 red ball.

Therefore, we have to find out $P(A_2/E)$

By Baye's theorem,

$$\begin{aligned} P(A_2/E) &= \frac{P(A_2)P(E/A_2)}{\sum_{i=1}^3 P(A_i)P(E/A_i)} \\ &= \frac{\frac{1}{3} \times \frac{2}{5}}{\frac{1}{3} \times \frac{2}{15} + \frac{1}{3} \times \frac{2}{5} + \frac{1}{3} \times \frac{1}{5}} \\ &= \frac{6}{11} \end{aligned}$$

Q2. Three machines A,B, and C produce respectively 60%, 30% and 10% of the total number of items of a factory. The percentages of defective output of these machines are 2%, 3% and 4% respectively. An item is selected at random and is found to be defective. Find the probability that the item was produced by machine C.

Solution:

Let A,B and C stand for the events of selection of an item from machines A,B and C.

Therefore, $P(A) = 60/100 = 0.6$, $P(B) = 0.3$, $P(C) = 0.1$

Let E be the event of selecting defective item.

$P(E/A) = 0.02$, $P(E/B) = 0.03$, $P(E/C) = 0.04$

We have to find $P(C/E)$

By Baye's theorem,

$$\begin{aligned} P(C/E) &= \frac{P(C).P(E/C)}{P(A).P(E/A) + P(B).P(E/B) + P(C).P(E/C)} \\ &= 0.16 \end{aligned}$$

Q3. In a bolt manufacturing factory, there are four machines A,B,C and D, manufacturing 20%, 15%, 25% and 40% respectively of the total production. Out of these 5%, 4%, 3% and 2% are defective. If a bolt is drawn at random, what is the probability that it is defective. If it is defective, what is the probability that it is manufactured by A or D ?

Solution:

Let A_1 , A_2 , A_3 , A_4 be the events of selecting a bolt manufactured by machines A,B,C and D.

$P(A_1) = 0.2$

$P(A_2) = 0.15$

Search Topics

To search type and hit enter

Topics

Analytics (14)
 Social Media Integration (2)
 Electronic Projects (13)
 Arduino (6)
 ESP8266, NodeMCU (1)
 Internet Of Things (7)
 Machine Learning-AI (2)
 Mathematics (21)
 Numerical Methods (10)
 Probability & Statistics (10)
 Transforms (1)
 Python Programming (4)

Latest Notes

Covid 19 morbidity counts follow Benford's Law ?
 Laplace Transforms for B.Tech. and Engineering – KTU Syllabus
 Numerical Methods for B.Tech. and Engineering – KTU Syllabus
 Using Composite Entities in Dialogflow
 Robot remote control using NodeMCU and WiFi
 Pandas DataFrame Load Data in Chunks
 Local Maxima and Minima to classify a Bi-modal Dataset
 Pandas DataFrame – multi-column aggregation and custom aggregation functions
 Building a Robotic Arm using Arduino
 Gravity and Motion Simulator in Python – Physics Engine

$$P(A3) = 0.25$$

$$P(A4) = 0.4$$

Let E be the event of selecting a defective bolt.

$$\text{Given } P(E/A1) = 0.05, P(E/A2) = 0.04, P(E/A3) = 0.03, P(E/A4) = 0.02$$

$$P(E) = 0.2(0.05) + 0.15(0.04) + 0.25(0.03) + 0.4(0.02) = 0.0315$$

We have to find $P(A1 \cup A4/E)$

$$\begin{aligned} P((A1 \cup A4)/E) &= \frac{P(A1/E)}{P(E)} + \frac{P(A4/E)}{P(E)} \\ &= \frac{0.2(0.05)}{0.0315} + \frac{0.4(0.02)}{0.0315} \\ &= 0.3175 + 0.254 \\ &= 0.5715 \end{aligned}$$

Q4. In a certain college, 4% of the boys and 1% of girls are taller than 1.8m. Further, 60% of the students are girls. If a student is selected at random and is found to be taller than 1.8m, what is the probability that the student is a girl ?

Solution:

Let A1 be the event of selecting a girl and A2 be the event of selecting a boy. Let E be the event where a student is taller than 1.8m.

$$P(A1) = 0.60; P(A2) = 0.40$$

$$P(E/A1) = 0.01; P(E/A2) = 0.04$$

$$\begin{aligned} P(A1/E) &= \frac{P(A1) \cdot P(E/A1)}{P(A1) \cdot P(E/A1) + P(A2) \cdot P(E/A2)} \\ &= \frac{0.6 \times 0.01}{0.6 \times 0.01 + 0.4 \times 0.04} \\ &= \frac{0.006}{0.006 + 0.016} \\ &= 0.2727 \end{aligned}$$

Q5: The chance that a doctor will diagnose a disease correctly is 60%. The chance that a patient will die after correct diagnosis is 40% and the chance of death by wrong diagnosis is 70%. If a patient dies, what is the chance that his disease was correctly diagnosed?

Solution:

Let A1 be the event of correct diagnosis

Let A2 be the event of wrong diagnosis

Let E be the event that a patient dies.

$$\text{Given, } P(A1) = 0.6; P(A2) = 0.4$$

$$P(E/A1) = 0.4; P(E/A2) = 0.7$$

$$P(A1/E) = \frac{P(A1)P(E/A1)}{P(A1)P(E/A1) + P(A2)P(E/A2)} = 0.4615$$

Q5. A bag contains 3 coins, out of which, one has both faces heads. Other two coins are normal and fair. A coin is chosen at random from the bag and tossed 4 times in succession. If head turns up each time, what is the probability that it is the two headed coin.

Solution:

Let A1 be the event of selecting 2 headed coin

Let A2 and A3 be the events of selecting remaining fair coins

Let E be the event of getting 4 heads in succession.

$$P(A1) = P(A2) = P(A3) = 1/3$$

$$P(E/A1) = 1; P(E/A2) = P(E/A3) = 1/2 \cdot 1/2 \cdot 1/2 \cdot 1/2 = 1/16$$

$$P(A1/E) = \frac{1/3 \cdot 1}{1/3 \cdot (1 + 1/16 + 1/16)} = \frac{1/18}{1/3 \cdot (1 + 1/8)} = \frac{1/18}{1/3 \cdot 9/8} = \frac{1/18}{3/8} = \frac{8}{27}$$

Q6. Three typists A,B,C typed 50%, 30% and 20% pages of a book. The percentage of wrongly typed pages by them is 3,4,5 respectively. If a page is selected from the book at random, what is the probability that it is wrongly typed and is typed by A ?

Solution:

$$P(A1) = 0.5, P(A2) = 0.3, P(A3) = 0.2$$

$$P(E/A1) = 0.03, P(E/A2) = 0.04, P(E/A3) = 0.05$$

$$P(E) = 0.5 \times 0.03 + 0.3 \times 0.04 + 0.2 \times 0.05$$

$$= 0.015 + 0.012 + 0.010$$

$$= 0.037$$

$$P(A1/E) = \frac{P(A1) \cdot P(E/A1)}{P(E)} = \frac{0.015}{0.037}$$

Q7. A tea set has 4 sets of cups and saucers. Two of these sets are of one colour and the other two sets are of different colours (A total of 3 colours). If the cups are placed randomly on saucers, what is the probability that no cup is on a saucer of same colour ?

Solution:

Let 2 sets (having same colour) be of colour C1 and the other 2 sets be of colours C2 and C3.

Let saucers be kept in C1C1C2C3 order.

Then the different possibilities in which cups can be kept are:

C1C1C2C3

C1C1C3C2

C1C2C1C3

C1C3C1C2

C1C2C3C1

C1C3C2C1

C2C1C1C3

C3C1C1C2

C2C1C3C1

C3C1C2C1

C2C3C1C1

C3C2C1C1

There are a total of 12 possibilities, of which, only last 2 are favourable to the event "no cup is on a saucer of same colour".

Therefore, the required probability = $2/12 = 1/6$

« Baye's Theorem

Covid 19 morbidity counts follow Benford's Law ? »

WRITTEN BY TESSY C



Lecturer and Research Scholar in Mathematics.

Topics

Analytics (12)

Arduino (6)

Electronic Projects (1)

ESP8266, NodeMCU (1)

Internet Of Things (7)

Search Topics

To search type and hit enter

Comments

Sinjo Alex on Local Maxima and Minima to classify a Bi-modal Dataset

Zederex Review on Pandas DataFrame – multi-column aggregation and custom aggregation functions

Machine Learning-AI (2)
Mathematics (1)
Numerical Methods (10)
Probability & Statistics (10)
Python Programming (4)
Social Media Integration (2)
Transforms (1)

Opal Wesolick on Building a Robotic Arm using Arduino
Justin Jose on Mosquitto MQTT Publish – Subscribe from PHP
mohan on Mosquitto MQTT Publish – Subscribe from PHP