

Challenge Problem 3

Jayati Dutta

1 PROBLEM

Prove that - Normal matrices are unitarily diagonalizable.

2 EXPLANATION

Let A be a normal matrix, then we have to prove A is unitary diagonalizable.

Definition:

A is normal if $AA^* = A^*A$

Definition:

A is unitary diagonalizable if there is a unitary matrix U and diagonal matrix D such that $UAU^* = D$.

Proof:

As A is normal, so $AA^* = A^*A$. Now, by mathematical induction first we will consider the orthonormal vectors for $n=2$.

Consider an eigen vector U of A corresponds to the eigen value λ and U is unit vector. Now V is considered in such a way that U, V forms an orthonormal basis in C^2 .

As U is an eigen vector, so $AU = \lambda U$. Now,

$$AU = U \begin{pmatrix} \lambda & U^{-1}AV \\ 0 & \end{pmatrix} \quad (2.0.1)$$

$$AU = U \begin{pmatrix} \lambda & \alpha \\ 0 & \beta \end{pmatrix} \quad (2.0.2)$$

$$A = U \begin{pmatrix} \lambda & \alpha \\ 0 & \beta \end{pmatrix} U^* \quad (2.0.3)$$

where $U^{-1}AV = \begin{pmatrix} \alpha \\ \beta \end{pmatrix}$ Since A is normal,

$$A^*A = U \begin{pmatrix} \bar{\lambda} & 0 \\ \bar{\alpha} & \bar{\beta} \end{pmatrix} U^* U \begin{pmatrix} \lambda & \alpha \\ 0 & \beta \end{pmatrix} U^* \quad (2.0.4)$$

$$\Rightarrow A^*A = U \begin{pmatrix} \bar{\lambda}\lambda & \alpha\bar{\lambda} \\ \lambda\bar{\alpha} & \alpha\bar{\alpha} + \bar{\beta}\beta \end{pmatrix} U^* \quad (2.0.5)$$

Similarly,

$$AA^* = U \begin{pmatrix} \bar{\lambda}\lambda & \alpha\bar{\beta} \\ \beta\bar{\alpha} & \bar{\beta}\beta \end{pmatrix} U^* \quad (2.0.6)$$

$$\text{As } AA^* = A^*A$$

$$\alpha\bar{\alpha} = 0 \quad (2.0.7)$$

$$\Rightarrow \|\alpha\| = 0 \quad (2.0.8)$$

$$\Rightarrow \alpha = 0 \quad (2.0.9)$$

$$\Rightarrow A = U \begin{pmatrix} \lambda & 0 \\ 0 & \beta \end{pmatrix} U^* \quad (2.0.10)$$

$$\Rightarrow A = UDU^* \quad (2.0.11)$$

So, A is unitary diagonalizable when D is a 2×2 matrix. Now, assume that the result holds for $(n-1)$. we can claim that there is y^* such that

$$AU = U \begin{pmatrix} \lambda & y^* \\ 0 & B \end{pmatrix} \quad (2.0.12)$$

where $B \in C^{(n-1) \times (n-1)}$ Similarly,

$$AA^* = U \begin{pmatrix} \bar{\lambda}\lambda + y^*\bar{y}^* & y^*\bar{B} \\ B\bar{y}^* & \bar{B}B \end{pmatrix} U^* \quad (2.0.13)$$

Now,

$$A^*A = U \begin{pmatrix} \bar{\lambda}\lambda & y^*\bar{\lambda} \\ \lambda\bar{y}^* & B\bar{B} + y^*\bar{y}^* \end{pmatrix} U^* \quad (2.0.14)$$

$$\text{As } AA^* = A^*A,$$

$$y^*\bar{y}^* = 0 \quad (2.0.15)$$

$$\Rightarrow \|y^*\| = 0 \quad (2.0.16)$$

$$\Rightarrow y^* = 0 \quad (2.0.17)$$

$$\Rightarrow A = U \begin{pmatrix} \lambda & 0 \\ 0 & B \end{pmatrix} U^* \quad (2.0.18)$$

$$\Rightarrow A = UDU^* \quad (2.0.19)$$

As $\bar{B} = B^*$ and $B^*B = BB^*$, so B is also normal. Let,

$$B = MD_1M^* \quad (2.0.20)$$

where M is unitary matrix and D_1 is diagonal matrix and both are in $C^{(n-1) \times (n-1)}$. Now,

$$AU = U \begin{pmatrix} \lambda & 0 \\ 0 & MD_1M^* \end{pmatrix} \quad (2.0.21)$$

$$AU = U \begin{pmatrix} 1 & 0 \\ 0 & M \end{pmatrix} \begin{pmatrix} \lambda & 0 \\ 0 & D_1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & M^* \end{pmatrix} \quad (2.0.22)$$

$$AU \begin{pmatrix} 1 & 0 \\ 0 & M \end{pmatrix} = U \begin{pmatrix} 1 & 0 \\ 0 & M \end{pmatrix} \begin{pmatrix} \lambda & 0 \\ 0 & D_1 \end{pmatrix} \quad (2.0.23)$$

$$AW = W \begin{pmatrix} \lambda & 0 \\ 0 & D_1 \end{pmatrix} \quad (2.0.24)$$

$$A = W \begin{pmatrix} \lambda & 0 \\ 0 & D_1 \end{pmatrix} W^* \quad (2.0.25)$$

where W is also a unitary matrix and $W = U \begin{pmatrix} 1 & 0 \\ 0 & M \end{pmatrix}$

This implies that if A is normal then A is unitary diagonalizable.