

Challenge Problem 2

Jayati Dutta

Download all and latex-tikz codes from

svn co <https://github.com/gadepall/school/trunk/ncert/geometry/figs>

1 PROBLEM

The center of the conic section:

$$y^2 - 2\sqrt{3}xy + 3x^2 + 6x - 4y + 5 = 0 \quad (1.0.1)$$

2 EXPLANATION

$$\mathbf{V} = \begin{pmatrix} 3 & -\sqrt{3} \\ -\sqrt{3} & 1 \end{pmatrix} \quad (2.0.1)$$

and

$$\mathbf{u} = \begin{pmatrix} 3 \\ -2 \end{pmatrix} \quad (2.0.2)$$

$$f = 5 \quad (2.0.3)$$

Now, from the eigen values of $\lambda = 0, 4$ we got the eigen vectors. For $\lambda_1 = 0$, the eigen vector of \mathbf{V} is \mathbf{p}_1 and for $\lambda_2 = 4$, the eigen vector of \mathbf{V} is \mathbf{p}_2 and those are:

$$\mathbf{p}_1 = \begin{pmatrix} 1 \\ \sqrt{3} \end{pmatrix} \quad (2.0.4)$$

and

$$\mathbf{p}_2 = \begin{pmatrix} -\sqrt{3} \\ 1 \end{pmatrix} \quad (2.0.5)$$

Now, \mathbf{P} should be chosen in such a way that $\mathbf{P}^T = \mathbf{P}^{-1}$ and $\mathbf{P}^T \mathbf{V} \mathbf{P} = \mathbf{D}$. but here it is not satisfying these equations. Here, $\mathbf{P}^{-1} = \frac{1}{4} \mathbf{P}^T$. Now, in this case we also observed that,

$$4(\mathbf{V}\mathbf{c} + \mathbf{u}) = \eta \mathbf{p}_1 \quad (2.0.6)$$

But this can be written as:

$$\mathbf{V}\mathbf{c} + \mathbf{u} = \frac{\eta}{2} \begin{pmatrix} 1 \\ \sqrt{3} \end{pmatrix} \quad (2.0.7)$$

So, \mathbf{P} should be selected as :

$$\mathbf{P} = \begin{pmatrix} \frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{1}{2} \end{pmatrix} \quad (2.0.8)$$

If I did it so, then $|\mathbf{P}|=1$ and $\mathbf{P}^{-1}=\mathbf{P}^T$ and $\mathbf{P}^T \mathbf{V} \mathbf{P}=\mathbf{D}$. So,

$$\mathbf{p}_1 = \begin{pmatrix} \frac{1}{2} \\ \frac{\sqrt{3}}{2} \end{pmatrix} \quad (2.0.9)$$

and

$$\mathbf{p}_2 = \begin{pmatrix} -\frac{\sqrt{3}}{2} \\ \frac{1}{2} \end{pmatrix} \quad (2.0.10)$$

Now, η can be calculated as :

$$\eta = 2\mathbf{p}_1^T \mathbf{u} \quad (2.0.11)$$

$$\eta = 2 \begin{pmatrix} \frac{1}{2} & \frac{\sqrt{3}}{2} \end{pmatrix} \begin{pmatrix} 3 \\ -2 \end{pmatrix} \quad (2.0.12)$$

$$\eta = 3 - 2\sqrt{3} \quad (2.0.13)$$

Now,

$$\mathbf{u}^T + \frac{\eta}{2} \mathbf{p}_1^T = \begin{pmatrix} \frac{15-2\sqrt{3}}{4} & \frac{-14+3\sqrt{3}}{4} \end{pmatrix} \quad (2.0.14)$$

$$\frac{\eta}{2} \mathbf{p}_1 - \mathbf{u} = \frac{3-2\sqrt{3}}{2} \begin{pmatrix} \frac{1}{2} \\ \frac{\sqrt{3}}{2} \end{pmatrix} - \begin{pmatrix} 3 \\ -2 \end{pmatrix} \quad (2.0.15)$$

$$\frac{\eta}{2} \mathbf{p}_1 - \mathbf{u} = \begin{pmatrix} \frac{-9-2\sqrt{3}}{4} \\ \frac{2+3\sqrt{3}}{4} \end{pmatrix} \quad (2.0.16)$$

Now considering the augmented matrix,

$$\begin{pmatrix} \frac{15-2\sqrt{3}}{4} & \frac{-14+3\sqrt{3}}{4} & -5 \\ 3 & -\sqrt{3} & \frac{-9-2\sqrt{3}}{4} \\ -\sqrt{3} & 1 & \frac{2+3\sqrt{3}}{4} \end{pmatrix} \xrightarrow{R_2 \leftarrow R_2 / \sqrt{3}} \quad (2.0.17)$$

$$\begin{pmatrix} \frac{15-2\sqrt{3}}{4} & \frac{-14+3\sqrt{3}}{4} & -5 \\ \sqrt{3} & -1 & \frac{-(2+3\sqrt{3})}{4} \\ -\sqrt{3} & 1 & \frac{2+3\sqrt{3}}{4} \end{pmatrix} \quad (2.0.18)$$

$$\xrightarrow{\begin{matrix} R_3 \leftarrow R_2 + R_3 \\ R_2 \leftarrow R_2 / \sqrt{3} \end{matrix}} \quad (2.0.19)$$

$$\begin{pmatrix} \frac{15-2\sqrt{3}}{4} & \frac{-14+3\sqrt{3}}{4} & -5 \\ 1 & -\frac{1}{\sqrt{3}} & -\frac{-(2+3\sqrt{3})}{4\sqrt{3}} \\ 0 & 0 & 0 \end{pmatrix} \quad (2.0.20)$$

$$\xrightarrow{R_1 \leftarrow R_1 / \frac{15-2\sqrt{3}}{4}} \begin{pmatrix} 1 & \frac{-14+3\sqrt{3}}{15-2\sqrt{3}} & -\frac{5}{15-2\sqrt{3}} \\ 1 & -\frac{1}{\sqrt{3}} & -\frac{-(2+3\sqrt{3})}{4\sqrt{3}} \\ 0 & 0 & 0 \end{pmatrix} \quad (2.0.21)$$

$$\xrightarrow{R_2 \leftarrow R_2 - R_1} \begin{pmatrix} 1 & \frac{-14+3\sqrt{3}}{15-2\sqrt{3}} & -\frac{5}{15-2\sqrt{3}} \\ 0 & -\frac{24-16\sqrt{3}}{15\sqrt{3}-6} & \frac{-12+39\sqrt{3}}{4\sqrt{3}(15-2\sqrt{3})} \\ 0 & 0 & 0 \end{pmatrix} \quad (2.0.22)$$

So,

$$\begin{pmatrix} 1 & \frac{-14+3\sqrt{3}}{15-2\sqrt{3}} \\ 0 & -\frac{24-16\sqrt{3}}{15\sqrt{3}-6} \end{pmatrix} \mathbf{c} = \begin{pmatrix} -\frac{5}{15-2\sqrt{3}} \\ \frac{-12+39\sqrt{3}}{4\sqrt{3}(15-2\sqrt{3})} \end{pmatrix} \quad (2.0.23)$$

$$\Rightarrow c_2 = -\frac{-12+39\sqrt{3}}{4(24-16\sqrt{3})} \quad (2.0.24)$$

$$\Rightarrow c_2 = 3.74 \quad (2.0.25)$$

$$c_1 = 1.12 \quad (2.0.26)$$

From there we can get $\mathbf{c} = \begin{pmatrix} 1.12 \\ 3.74 \end{pmatrix}$. The vertex of parabola will be at (1.12, 3.74).

2.1. Verification of the above problem using python code.

Solution: The following Python code generates Fig. 0

```
codes/check_parab.py
```

So the solution is matching with the given plot, hence it is verified.




Fig. 0: Parabola