	Model Solution	Scoring	
A	Find $g'(8)$. Give a reason for your answer.		
	$g'(x) = f(x)$	Considers $g'(x) = f(x)$	**Point 1 (P1)**
	$g'(8) = f(8) = 1$	Answer	**Point 2 (P2)**
	Scoring Notes for Part A		

- **P1** is earned for $g' = f$, $g'(x) = f(x)$, or $g'(8) = f(8)$ in part A.
- A response of $g'(8) = f(8) = 1$ earns both **P1** and **P2**.
- A response that does not earn **P1** can earn **P2** with an implied application of the Fundamental Theorem of Calculus (e.g., $g'(8) = 1$ or $f(8) = 1$).
- A response of $g'(8) = f(8) - f(6) = 1$ earns **P2** but not **P1**.

© 2025 College Board

B Find all values of x in the open interval $-6 < x < 12$ at which the graph of g has a point of inflection. Give a reason for your answer.

The graph of g has a point of inflection where $g'' = f'$ changes sign, which is where $g' = f$ changes from decreasing to increasing or vice versa. The graph of g has points of inflection at $x = -3$ and $x = 6$ because f changes from decreasing to increasing there. The graph of g also has a point of inflection at $x = 3$ because f changes from increasing to decreasing there.	Answer	**Point 3 (P3)**
	Reason	**Point 4 (P4)**

Scoring Notes for Part B

- **P3** is earned only for an answer of $x = -3$, $x = 3$, and $x = 6$. If any other/additional values of x in $-6 < x < 12$ are declared to be points of inflection, the response does not earn either **P3** or **P4**. Consideration of $x = -6$ or of $x = 12$ does not impact scoring.
- To earn **P4**, a response must tie the reason to the given graph of f.
  - A response of "g has a point of inflection at $x = -3$, $x = 3$, and $x = 6$ because f changes from increasing to decreasing or decreasing to increasing there" earns both **P3** and **P4**.
  - A response of "g has a point of inflection at $x = -3$, $x = 3$, and $x = 6$ because the slope of f changes sign there" earns both **P3** and **P4**.
  - A response of "g has a point of inflection at $x = -3$, $x = 3$, and $x = 6$ because f attains relative extrema there" earns both **P3** and **P4**.
  - A response of "g has a point of inflection at $x = -3$, $x = 3$, and $x = 6$ because g changes concavity there" earns **P3** but not **P4**.
  - A response of "g has a point of inflection at $x = -3$, $x = 3$, and $x = 6$ because $g'' = f'$ changes sign there" earns **P3** but not **P4**.
  - A response that relies upon an ambiguous term such as "the function" or "the graph" does not earn **P4**.
- **Special case:** A response with two of the three correct x-values with correct reasoning and no other/additional values of x declared to be points of inflection earns **P4** but not **P3**.

© 2025 College Board

C Find $g(12)$ and $g(0)$. Label your answers.

$g(12) = \int_6^{12} f(t)\,dt = \frac{1}{2} \cdot 6 \cdot 3 = 9$	$g(12)$	**Point 5 (P5)**
$g(0) = \int_6^{0} f(x)\,dx = -\int_0^{6} f(x)\,dx = -\frac{\pi}{2}3^2 = -\frac{9\pi}{2}$	$g(0)$	**Point 6 (P6)**

Scoring Notes for Part C

- Unlabeled values do not earn either **P5** or **P6**.
- **P5** is earned for a response of $g(12) = 9$, with or without supporting work.
- **P6** is earned for a response of $g(0) = -\frac{9\pi}{2}$, with or without supporting work.

  Note: Incorrect communication between the label "$g(0)$" and the answer will be treated as scratch work and will not impact scoring. For example, $g(0) = \int_0^{6} f(x)\,dx = -\frac{9\pi}{2}$ earns **P6**.

© 2025 College Board

D Find the value of x at which g attains an absolute minimum on the closed interval $-6 \le x \le 12$. Justify your answer.

For $-6 \le x \le 12$, g attains a minimum either when $g'(x) = f(x) = 0$ or at an endpoint. $g'(x) = f(x) = 0$	Considers $g'(x) = 0$	**Point 7 (P7)**
$\Rightarrow \quad x = 0,\ x = 6$ $\begin{array}{c\|c} x & g(x) \\ \hline -6 & 0 \\ 0 & -\frac{9\pi}{2} \\ 6 & 0 \\ 12 & 9 \end{array}$	Justification	**Point 8 (P8)**
Therefore, on the closed interval $-6 \le x \le 12$, g attains an absolute minimum value at $x = 0$.	Answer	**Point 9 (P9)**

Scoring Notes for Part D

- **P7** is earned for considering $g'(x) = 0$ or $f(x) = 0$. **P7** is not earned by just presenting $x = 0$ and $x = 6$.

  A response that discusses the sign of $g'(x)$ or $f(x)$ changing OR uses the phrase "critical points of g" also earns **P7**.
- To earn **P8** using a candidates test, a response must make a global argument by providing evaluations or reasoning for each of $g(-6)$, $g(0)$, $g(6)$, and $g(12)$ (and no other x-values).
- Alternate justification and answer:

  Because $g'(x) \le 0$ (or $f(x) \le 0$) for $-6 \le x < 0$ and $g'(x) \ge 0$ (or $f(x) \ge 0$) for $0 < x \le 12$, the absolute minimum of g occurs at $x = 0$.
- A response that presents a local argument (such as a First Derivative Test or a Second Derivative Test) or an incorrect global argument does not earn **P8** but is eligible for **P9** with the correct answer of $x = 0$.
- For **P8**, values of $g(0)$ and $g(12)$ can be imported from part C. A response can earn **P9** with an answer that is consistent with the imported values.

© 2025 College Board

Part B (BC): Graphing calculator not allowed
Question 5 — 9 points

General Scoring Notes

- The model solution is presented using standard mathematical notation.
- Answers (numeric or algebraic) need not be simplified. Answers given as a decimal approximation should be accurate to three places after the decimal point. Within each individual free-response question, at most one point is not earned for inappropriate rounding.

Let $y = f(x)$ be the particular solution to the differential equation $\frac{dy}{dx} = (3 - x)y^2$ with initial condition $f(1) = -1$.

	Model Solution	Scoring	
A	Find $f''(1)$, the value of $\frac{d^2y}{dx^2}$ at the point $(1, -1)$. Show the work that leads to your answer.		
	$\frac{d^2y}{dx^2} = -y^2 + (3 - x)2y\frac{dy}{dx}$	Product rule	**Point 1 (P1)**
		Chain rule	**Point 2 (P2)**
	$f'(1) = \left.\frac{dy}{dx}\right\|_{(x,y)=(1,-1)} = (3-1)(-1)^2 = 2$ $f''(1) = \left.\frac{d^2y}{dx^2}\right\|_{(x,y)=(1,-1)} = -(-1)^2 + (3-1)(2)(-1)(2) = -9$	$f''(1)$	**Point 3 (P3)**

© 2025 College Board