

01

Introduction to Biology**Nature scope and importance of biology with reference to challenges faced by the mankind**

Biology is a Science which is focused on studying of living organisms. (Bios- Life, logos "study of")

The concept of "LIFE" is not easy to define. Still scientists are unable to provide an acceptable definition for life.

"Life" is something special and unique which cannot be explained using laws of chemistry and physics.

Biology is a subject which is very complex and vast. Hence for the convenience of studying, it has been divided into three primary branches: Zoology (the study of animals), Botany (study of plants) and Microbiology (study of microorganisms). Some areas of study in these branches:

- Cell Biology (studying cells)
- Histology (studying tissues)
- Anatomy (studying about gross structure of the body)
- Physiology (studying function)
- Biochemistry (studying biological molecules)
- Genetics (studying inheritance)
- Ecology (studying environment)

Issues pertaining to Biology

Understanding biological Diversity

At present our planet is rich in diversity. Life on earth formed around 3.8 billion years ago. The first formed organisms are believed to be heterotrophic, anaerobic prokaryotes. Since then the evolutionary process resulted in the extensive biodiversity which exists in the biosphere now.

Scientists assume, based on their studies that there are about 10 to over 100 million species in the world. There is a dynamic relationship between living world and the inanimate world and each and every organism has a specific role in the environment for existence of the Biosphere.

The variety of life on earth, the number of species of plants, animals and microorganisms, the diversity of genes in these species, the different ecosystems on the earth such as deserts, rainforests and coral reefs are all part of a biologically diverse earth.

Understanding the human Body and its functions

When studying biology, especially by studying histology, anatomy of the human body, one can gain the knowledge about the structure of the organs. This results in understanding and appreciation of the organization of the human body and understanding the functions of different organ systems and the relationship between structure and functions.

Sustainable use and Management of natural resources and Environment

Natural resources are sources of materials and energy found naturally which are used in everyday life and for economic development.

These natural resources are limited on earth. Due to the increase of growth rate of human population, overuse of natural resources is taking place. It causes threat of depletion of natural resources.

Due to over exploitation of natural resources, various environmental problems arise such as;

- Environmental pollution
- Loss of Biodiversity
- Desertification

Hence to overcome the above problems management of natural resources and Environment should be practiced. Knowledge of Biology is useful to bring about remedies for the above problems.

Sustainable Food production

Sustainable food production is the production of sufficient amounts of food for the

human population using environmentally safe methods.

The current human population is about 7 billion and expected to be double in less than 40 years. Therefore, for the survival of human beings sustainable food production is necessary.

To maintain sustainable food production following methods can be applied, which are based on knowledge in biology.

- Production of high yielding varieties of plants and animals.
- Production of disease resistant plants and animal varieties.
- Improve the post harvest technological methods.

Understanding plant life

Plants are the primary producers in the world. All the animals depend directly or indirectly on plants. Therefore understanding plant life is important. As the time human population is increasing we need to increase the productivity. Therefore understanding plant function and biology is important to produce high yielding plants, disease resistant plants, etc.

Understanding diseases and causes

To maintain healthy human body one should have the knowledge of causes of the diseases and their effects.

Some dangerous diseases which exist in current world are non communicable diseases such as cancers, heart diseases, diabetes, chronic renal diseases and communicable diseases such as dengue, AIDS, etc.

Cancers- causes for this is not fully understood yet. Cancers are one of the leading causes of death .

AIDS- is a viral disease which is a serious and growing health problem worldwide.

Heart diseases- This is also a serious and growing health problem worldwide. Causes are not fully understood yet.

Chronic renal diseases- In Sri Lanka, recently CKDu has become a serious health problem.

Currently scientists are working on prevention, remedial measures and cures for such diseases.

Solving some legal and ethical issues

Knowledge and application of biological concepts is important in solving some legal issues, such as parentage testing, in criminal investigations and to solve immigration disputes.

DNA fingerprinting is used in above circumstances.

The nature and the organizational patterns of the living world

In accordance with different criteria we can see a diversity among living organisms. Organisms are diverse based on size, shape, form and habitats.

- Living organisms show a wide range of variation in size, shape, form and habitat.
- Size – Bacteria – $0.25\ \mu\text{m}$ – $2\ \mu\text{m}$ to Giant Sequoia (Giant Red Wood)– 100m
- Shape – Organisms are diverse in shape, Ex: Cylindrical (earth worm), streamline shape(birds, fish)
- Form – Unicellular (Amoeba), multicellular (any plant or animal)
- Habitat – Terrestrial (Rat), aquatic (Fish), arboreal (Loris), aerial(Birds)

Characteristics of organisms

In order to survive, each organism whether simple or complex must be able to perform certain functions. Following features are the characteristics of organisms.

(i) Order and organization

From molecular level to biosphere there is an order and organization in organisms to perform their biological activities efficiently.

Lower level components are organized in a methodical pattern in upper level to make it most efficient.e.g: plant leaf and human eyes.

(ii) Metabolism

The sum of all chemical activities taking place in an organism is its metabolism. It includes catabolic reactions and anabolic reactions.

(iii) Growth and development

All organisms begin their life as a single cell. During growth an irreversible increase in dry mass occurs, which is characterized only by the living. Irreversible changes that occur during the life span of an organism is development. Growth and development are two consequent processes that happen in the life span of organisms.

(iv) Irritability and coordination

Irritability is the ability to respond to stimuli from both internal and external environment. Movement of organisms occurs as a result of irritability and coordination. In animals

this happens as a result of coordinated efforts of nervous, hormonal, muscular and skeletal systems

(v) Adaptation

Adaptation is a peculiarity of structure, physiology or behavior that promotes the likelihood of an organism's survival and reproduction in a particular environment. E.g: Sunken stomata in Xerophytes, Viviparity in some mangroves, Splayed out foot in camels.

(vi) Reproduction

Ability to produce offspring for continuous existence of species

(vii) Heredity & Evolution

Organisms have genes that pass from one generation to the next and control specific physiological, morphological and behavioral characters of organisms.

Ability of organisms to change over time as a result of genetic modification is evolution

Many non living entities may have one or more of these characteristics but not all of them e.g., crystals grow, waves move but only living organisms display all these characteristics simultaneously or at some point during their life cycle.

By considering this it can be said that these are occurring in single celled organisms as well as highly complex organisms such as humans and Anthophytes (flowering plants).

Hierarchical levels of organization of living things

The cell is the basic structural and functional unit of life. Some organisms are unicellular while others are multicellular. Cell is composed of several organelles which are formed by different organic molecules. Then hierarchical levels of organization of living things can be constructed by using relevant examples at each level.

Molecules, Organelles, Cells, Tissues, Organs, Organ systems, Organisms, Populations, Communities, Ecosystems, Biosphere

02

Chemical and cellular basis of life**Elemental composition of living matter**

There are about ninety two elements naturally occur in earth's crust. Of which, about 20-25% elements are essential to continue healthy life and reproduction. (about 25- elements are essential for humans and about 17 for plants).

Oxygen (O), Carbon (C), Hydrogen (H), and Nitrogen (N) make up 96% of living matter.

Calcium (Ca), Phosphorous (P), potassium (K) and sulphur (S)- make up most of the remaining 4% of the mass of the organism.

In humans, C, H, O, N- accounts for 96.3% of the body mass and Ca, P, K, S, Na, Cl, Mg and trace elements accounts for the remaining 3.7%. (e.g. B (Boron), Co (Cobalt), Cu (Copper), Cr (Chromium), F (Fluorine), I (Iodine), Fe (Iron),


Mo (Molybdenum), Mn (Manganese), Se (Selenium), Si (Silicon), Sn (Tin), V (Vanadium), Zn (Zinc)

Physical and chemical properties of water important for life

Water is a vital inorganic molecule; life could not exist on this planet without water. It is important due to following reasons,

1. Vital chemical constituent of living cell
2. Provides a biological medium for all organisms

Most of above properties are based on the chemical structure of water molecule. Physical and chemical properties of water molecule provide the ability to render the vitality. Water molecule is a small, polar and angular molecule.



δ+ partial positive

δ- partial negative

Fig 2.1: Chemical structure of the water molecule

Polarity is an uneven charge distribution within a molecule. In water molecule, oxygen atom is slightly negative and hydrogen atom is slightly positive. Weak attractions between the slightly polar hydrogen atom of one water molecule and the slightly polar oxygen atom of adjacent water molecule are known as hydrogen bonds. These hydrogen bonds play a major role in maintaining all the properties of water.




Fig 2.2: Hydrogen bonding in water

The properties of water arise due to attractions of different water molecules. When the water is in liquid form its H bonds are very fragile. H bonds form, break and reform with great frequency.

Four major properties of water to maintain life on earth

- Cohesive behavior
- Ability to moderate temperature
- Expansion upon freezing
- Versatility as a solvent

Properties of water related to functions

1. Cohesive behavior

Attraction between water molecules due to hydrogen bonding is known as cohesion. Attraction between water molecules and other substances are known as adhesion. Both of the above properties of water allow it to act as a transport medium.

Due to cohesion between water molecules, water and dissolved substances such as minerals and nutrients transport through vascular tissues, xylem and phloem against gravity.

Adhesion between water molecules and cell walls also helps in conduction of water and dissolved substances.

Water has a high surface tension. This ability is given to water molecules, due to cohesion between the water molecules. Therefore, in an aquatic system, upper surface water molecules are attracted by lower surface molecules and it forms a water film. Small insects e.g. water skaters can walk on the surface of a pond.

2. Ability to moderate temperature

Water can absorb or release a relatively high amount of heat energy by a slight change in its own temperature.

Due to the high specific heat, water will function as thermal buffer in living system and aquatic bodies during the temperature fluctuations on earth.

Due to the high heat of vaporization, with the minimum loss of water an organism can release much heat energy. Therefore, body surface of an organism maintained as cool surface.

e.g. Prevent from overheating.

Evaporation of sweat from human skin helps to maintain the body temperature at constant level.

Transpiration in plants keeps the plant body surface as a cool surface and prevent from becoming too warm in the sunlight.

3. Expansion upon freezing

Generally, in an increase in temperature of any substances, reduces their density and on the other hand, in a decrease in temperature increases their density. When the temperature of water falls below 4 °C, it begins to freeze and forms a crystalline lattice called ice cubes. Therefore water has the maximum density at 4°C. Hence, ice floats on the surface of water bodies. It is an important property of water in polar regions, where, organisms in aquatic bodies can survive during the winter.

4. Versatility as a solvent

This ability is given to water due to their polarity. Polar molecules (e.g. Glucose), non polar ionic (e.g. NaCl), both polar and ionic (e.g. lysozymes) can dissolve in water, because water molecules surround each of the solute molecules and form hydrogen bonds with them. Solubility depends on polarity and not in their ionic nature.

Chemical Nature and Functions of Main Organic Compounds of Organisms

Carbohydrates

Most abundant group of organic compound on earth is carbohydrates. Major elemental composition is C, H, and O. Hydrates of carbon contain the same proportion of H: O which equals to 2:1 as in water. General formula is $C_x(H_2O)_y$. Three major groups of carbohydrates are monosaccharides, disaccharides and polysaccharides.

Generally carbohydrates include sugars (monosaccharides and disaccharides) and polysaccharides.

Monosaccharides

The simplest form of carbohydrates having general molecular formula as $(CH_2O)_n$ are monosaccharide. Where C varies from 3-7. All monosaccharide are reducing sugars, water soluble and occur in crystalline form.



According to the number of carbon atoms, they are named as;

- 3C- Triose e.g. Glyceraldehydes (Phosphoglyceraldehyde is a derivative of Triose)
- 4C- Tetroses.g. Erythrose (rare in nature)
- 5C- Pentoses.g. Ribose, Deoxyribose, Ribulose (RUBP is a derivative of ribulose)
- 6C- Hexoses e.g. Glucose, Fructose, Galactose



According to the type of carbonyl (Keto, aldo)group, they are classified as;

- a. Aldoses-glucose, galactose
- b. Ketoses-fructose

Aldose

*Fig 2.3: Solid form of glucose**Fig 2.4: Aqueous form of Glucose molecule*

Ketose

*Fig 2.5: Solid form of fructose**Fig 2.6: Aqueous form of fructose*

In aqueous media some monosaccharides are in ring form (No need to memorize the chemical structures)

Disaccharides

They are sugars formed by joining two monosaccharides by a glycosidic bond.




Fig 2.7: Formation of sucrose





Fig 2.8: Formation of maltose

(no need to memorize the chemical structures)

Glycosidic bond is formed by removal of a water molecule from two adjacent monosaccharides by a condensation reaction. Water molecule is formed from OH group of one monosaccharide molecule and H from adjoining monosaccharide molecule.



Maltose and lactose are reducing sugars and sucrose is a non reducing sugar.

Polysaccharides

They are macromolecules and biopolymers. Polysaccharides are made up of few hundred to a few thousand monosaccharide subunits

They are non crystalline, water insoluble, and not considered as sugars.

Some polysaccharides act as storage components where others contribute to the structure of living organisms. Based on their function they are categorized as storage polysaccharides and structural polysaccharides.

- Storage- Starch, Glycogen
- Structural- Cellulose, Hemicellulose, Pectin

Based on their architecture they are categorized as

- Linear forms- Cellulose, Amylose
- Branched forms- Glycogen, Amylopectin, Hemicellulose

Table 2.1: Major polysaccharides, their monomers and functions

Polysaccharide	Monomer	Functions
Starch	Glucose	Stored in plants
Glycogen	Glucose	Stored in animals and fungi
Cellulose	Glucose	Component of Cell wall
Inuline	Fructose	Stored in tubers of Dahlia
Pectin	Galacturonic acid	Component of Middle lamella of plant cell wall
Hemicellulose	Pentose	Component of Plant cell walls
Chitin (nitrogen containing polysaccharide)	Glucosamine	Component of Fungal cell walls and exoskeleton of Arthropods

Functions of carbohydrates

Monosaccharides

- Energy source
- Building blocks of disaccharides and polysaccharides (disaccharides such as maltose, sucrose and polysaccharides such as starch, glycogen)
- Components of nucleotides (DNA, RNA)

Disaccharides

- Storage sugar in milk- Lactose
- Translocation in phloem –Sucrose
- Storage sugar in sugarcane- Sucrose

Polysaccharides

a.) Storage polysaccharides-

- starch stores glucose as energy source in plants and chlorophytes
- glycogen stores glucose as energy source in animals and fungi
- inulin stores fructose as energy source in Dahlia tubers

b.) structural polysaccharides-

- Cellulose in the cell walls of plants and chlorophytes
- Pectin in the middle lamella of plant tissues.
- Hemicellulose in cell walls of plants.
- Peptidoglycan in the cell walls of prokaryotes.
- Chitin in the cell walls of fungi and in exoskeleton in Arthropods.

Lipids

- Diverse group of hydrophobic molecules
- Large biological molecules but not considered as polymers or macromolecules.
- Consist of C, H, O and H:O ratio is not 2:1. Comparatively more H are present.
- Biologically important types of lipids: Fats, Phospholipids and Steroids.

Fats

Fats are made up of glycerol and fatty acids; Glycerol belongs to alcohol group having 3 carbons where each of them bear single hydroxyl group. Fatty acids are hydrocarbon chains with long (16-18) carbon skeleton with a carboxyl group at its one terminal.

Fatty acid molecules bind to each hydroxyl group of glycerol by ester bond. Resulting fat molecules are called as triacylglycerol.




Fig 2.9: Formation of Triacylglycerol

Hydrocarbon chains of fatty acids contribute to the hydrophobic nature of the fats. Based on the nature of hydrocarbon chains of fatty acids, they are categorized as

- Saturated fats- fats are made up of saturated fatty acids: fatty acids with hydrocarbons having no any double bonds. Usually animal fats come under this category. They are mostly solid at room temperature. e.g: butter
- Unsaturated fats- fats are made up of unsaturated fatty acids- fatty acids with hydrocarbons having one or more double bonds. Usually plant fats come under


this category. They are mostly liquid in room temperature. e.g: vegetable oils. Unsaturated fats may classify based on the nature of their double bonds. a) *Cis* Unsaturated fat b) *Trans* Unsaturated fat

Consumption of excess saturated fats and trans unsaturated fats contribute atherosclerosis.

Phospholipids

Phospholipids are major components of the cell membranes. They are composed of two fatty acids and one phosphate group attached to one glycerol molecule. The phosphate group gives the negative electrical charge to the phospholipid molecule. Typically an additional polar molecule or small charged molecule is also linked to the phosphate group e.g. choline.

The two ends of the phospholipids show different behavior. The hydrocarbon tails are hydrophobic while phosphate group and its attachment (head) are hydrophilic.



(no need to memorize the structure)

Fig 2.10: structure of the phospholipid molecule

Functions of Lipids

- food reserve as energy source (triglycerides such as fats and oils)
- maintain the fluidity of plasma membrane (phospholipids, cholesterol)
- act as signaling molecules (eg. Hormones) that travel through the body
- found as components of animal cell membrane (cholesterol)

Protein

Proteins are made up of amino acids. Twenty different amino acids are involved in the formation of proteins. Elemental composition is C, H, O, N and S. At the centre of the amino acid is an asymmetric carbon atom except in glycine. Each amino acid is composed of an amino group, a carboxyl group, a hydrogen atom and a variable group symbolized by R, which is an alkyl group. In the case of glycine R is replaced by H atom. The R group also called the 'side chain' differs with each amino acid where as the other groups are in the 'back bone' (including the H atom).




Fig 2.11: Structure of an Amino acid molecule

Amino acids may have one or more carboxyl groups and amino groups. Amino group has alkaline nature and carboxyl group has acidic nature. When both characteristics are found in one molecule they are known as amphoteric molecules. Therefore, amino acids are amphoteric.

Two Amino acids undergo condensation reaction by removing a water molecule from both and result a bond known as peptide bond;




Fig 2.12: Formation of peptide bond

Protein is composed of one or more polypeptide chains which are composed of amino acids.

Levels of protein structures

There are four levels of structure which play important roles in their functions;

- a) Primary
- b) Secondary
- c) Tertiary
- d) Quaternary

a) Primary structure

The unique sequence of linearly arranged amino acids linked by peptide bonds is the primary structure of proteins.

b). Secondary structure

The primary structure of a single polypeptide chain coils and folds, as a result of intra molecular hydrogen bonds between the oxygen atoms and the hydrogen atoms attached to the nitrogen atoms, of the same poly peptide chain backbone, to form the secondary structure, which is either β pleated or alpha helical.

- Alpha helix- e.g.Keratin.
- β pleated sheet e.g.spider's silk fiber





Fig 2.13: beta pleated sheet and alpha helix of secondary structures of proteins

b) Tertiary structure

Usually the secondary polypeptide chain bends and folds extensively forming a precise compact unique, functional and three-dimensional shape resulting from following interactions between the side chain/ R-group of amino acids;

- H bonds
 - Disulphide bonds
 - Ionic bonds
 - Van der Waals interactions/ Hydrophobic interactions
- e.g. most of the enzymes, myoglobin, albumin

c) Quaternary structure

Aggregation of two or more polypeptide chains involve in the formation of one functional protein. Separate chains are called protein subunits which were held together by inter and intra-molecular interactions.

e.g. Haemoglobin, Collagen




Fig 2.14: structure of the hemoglobin molecule

Denaturation of proteins

Denaturation of protein is the loss of specific chemical three dimensional shape due to the alteration of weak chemical bonds and interactions.

Agents affecting the denaturation

1. High temperature and high energy radiation
2. Strong acids, alkaline and high concentrations of salts
3. Heavy metals
4. Organic solvents and detergents

Functions of the proteins

Table 2.2 Functions of Proteins

Type of protein	Example	Functions
Catalytic protein	Pepsin, Amylase	Catalyze biochemical reaction
Structural protein	Keratin,	Prevent desiccation
	Collagen	Provide strength and support
Storage	Ovalbumin	Storage protein in egg
	Casein	Storage protein in milk
Transport	Haemoglobin	Transport O ₂ and CO ₂
	Serum albumin	Transport fatty acids
Hormones	Insulin	Regulate blood glucose level
	Glucagon	
Contractile/ Motor	Actin/Myosin	Contraction of muscle fibres
Defensive	Immunoglobins	Eliminate foreign bodies

Nucleic acids

Nucleic acids are Polymers exist as polynucleotides made up of monomers called nucleotides. They contain C, H, O, N and P. Nucleic acids are macromolecules, biopolymers. There are two types of Nucleic acids: DNA (Deoxyribo nucleic acids) and RNA (Ribonucleic acids).

Structure of nucleotides

Nucleotides have 3 components; namely pentose sugar, nitrogenous base and a phosphate group




Fig 2.15: Structure of nucleotides (no need to memorize chemical structures)

A nucleotide without a phosphate group is called a nucleoside.e.g. Adenosine, Guanosine

Pentose sugar

Pentose sugars are two types; namely Deoxy ribose and ribose (in deoxyribose one oxygen atom is less than in ribose)

Nitrogenous bases

There are two major groups of nitrogenous bases:

1. Purines- larger in size with two rings
2. Pyrimidines- smaller in size with a single ring

In purines there are two types; namely Adenine, Guanine. In pyrimidines there are three types, Thymine, Uracil and Cytosine. Bases are commonly represented by letters A, G, T, U and C respectively.

Phosphate group

It gives the nucleic acids the acidic nature.

Formation of nucleic acids

Millions of nucleotides join by phospho-di-ester bond to form polynucleotide chains by condensation between the –OH group of the phosphate of one nucleotide with the –OH attached to 3rd carbon of pentose sugar of the other. These bonds result in a backbone with a repeating pattern of sugar-phosphate units. Nucleic acids are linear polymers of nucleotides. There are two kinds of nucleic acids depending on the type of the sugar molecules involved. If the sugar molecule in the nucleotide is deoxyribose, the nucleic acid is (DNA). If the pentose sugar is ribose, then the nucleic acid is RNA. DNA contains Adenine, Thymine, Guanine and Cytosine and RNA contains Adenine, Guanine, Cytosine and Uracil.

Structure of DNA molecule (Watson and Crick model)

DNA molecules have two anti-parallel polynucleotide chains that spiral around an imaginary axis, forming a double helix. The two sugar-phosphate backbones run in opposite directions from each other, and the arrangement is referred to as anti-parallel. The sugar phosphate backbones are on the outside of the helix, and the nitrogenous bases are paired in the interior of the helix. The two strands are held together by hydrogen bonds between the paired nitrogen bases.

Base pair rule

Always a purine base, pairs with a specific pyrimidine base,

A=T (2 hydrogen bonds)

G≡C (3 hydrogen bonds)

Hence two chains (strands) are said to be complementary to each other. These pairs are known as complementary base pairs. In this original double helical structure, one complete turn consists of ten base pairs as shown in the diagram.




Fig2.16: The structure of the DNA and RNA molecules (no need to memorize chemical structures)

Functions of DNA

- Store and transmit genetic information from one generation to the next generation
- Store the genetic information for protein synthesis

Structure of RNA

This is normally a single stranded nucleic acid composed of ribo-nucleotides containing bases, Uracil (U), Cytosine (C), Guanine (G), Adenine (A). Complementary base pairing between two RNA molecules or within the same molecule may occur in some. Complementary base pairing facilitates three dimensional shapes essential for their functioning. Adenine binds with Uracil with two hydrogen bonds and Guanine binds with Cytosine with three hydrogen bonds. There are three types of RNA present in cells,

1. Messenger RNA (mRNA)
2. Transfer RNA (tRNA)
3. Ribosomal RNA (rRNA)

1. Messenger RNA

Messenger RNA is a linear molecule and is the least abundant type of RNA in a cells comparatively. There are two functions;

- Copies the genetic information stored in DNA molecule as a sequence of nitrogenous bases
- Transports genetic information from nucleoplasm to the site of protein synthesis (ribosome) through nucleopores

2. Transfer RNA (tRNA)

Smallest RNA molecule. Linear, but forms three- looped structure as shown in the diagram.




Fig 2.17: Structure of the tRNA molecule

Function - transportation of amino acids to the site of protein synthesis.

3. Ribosomal RNA

It is the most abundant type of RNA. rRNA has a complex irregular structure. It provides the site where polypeptide chains are assembled.

Differences between DNA and RNA

1. DNA is double stranded molecule while RNA is a single stranded molecule.
2. DNA consists of A, T, G and C and absence of U, while RNA consists of A, U, G and C and absence of T
3. Sugar molecule in RNA is ribose, while in DNA it is deoxyribose.

Nucleotides other than those found in nucleic acids

ATP, NAD⁺, NADP⁺, FAD and their functions

Functions of ATP

- Universal energy carrier

Functions of NAD⁺

- Act as a coenzyme
- Act as an electron carrier
- Function as an oxidizing agent during respiration

Functions of NADP⁺

- Act as coenzymes
- Act as an electron carrier
- NADP⁺ act as a reducing agent in photosynthesis

Functions of FAD

- Act as a coenzyme
- Act as an electron carrier

Contribution of microscope to the expansion of knowledge on cells and cellular organization

Advancement of the cytology is mostly based on the microscopy. The discovery and early study of cells progressed with the invention of microscope.

Light microscope

Visible light is passed through the specimen and then through glass lenses. The lenses refract the light in such a way that the image of the specimen is magnified as it is projected into the eye. The simplest microscope is a single lens.

The compound light microscope

Compound light microscopes are commonly used in school laboratories and it is used in medical laboratories as a diagnostic tool.

Resolution power and magnification are important parameters which can be seen in a microscope.

Magnification is ratio of an object's image size to its actual size. Usually the maximum magnification of light microscope is 1000 times the actual size of the specimen)

Resolution power is minimum distance between two points that can be distinguished as separate points (resolution power of light microscope is 0.2µm). It is a measure of

the clarity of the image.

Magnification is limited due to the resolution.

Light from an object (specimen on the slide) passes first through objective lens. Then produce a magnified image.

Above image then acts as an object for the second lens (the eye piece lens) which further magnifies it.

The total magnification is hence the product of the magnification of each lens.

$$\text{Total magnification} = \text{Magnification of objective lens} \times \text{Magnification of eyepiece lens}$$

e.g- If magnification of Objective lens = $\times 40$, eyepiece = $\times 15$

Total is $= 15 \times 40 = \times 600$ time magnified

The Electron Microscope

The limitation imposed upon the resolution power of the light microscope by the wavelength of light. The resolution power is inversely proportional to the wavelength. Due to this, scientists considered the use of other forms of radiations with comparatively shorter wavelengths.

As a result, electron microscopes were developed. In electron microscopy, a beam of electrons is focused through the specimen or on to its surface.

This means, that in theory, the electron microscope should be able to magnify objects up to 1×10^8 times. In practice, it magnifies just over 5×10^5 times.

Electron microscopes have revealed many organelles and other sub cellular structures those were impossible to resolve with the light microscopes.

There are two types of electron microscopes.

1. Transmission electron microscopes (TEM)
2. Scanning electron microscopes (SEM)

Transmission electron microscopes

It is used to study the internal structures of cells. In this microscope, a beam of electrons is passed through a thin, especially prepared slice of material. A very thin specimen is used. Specimens stained with heavy metals which attach more to certain cellular structures than other areas. Image reflects the pattern of electrons passed through the specimen, displays on a screen. While electrons pass through the specimen, more electrons may get displayed in regions where structures were densely stained.

Scanning electron microscopes

In this instrument, a fine beam of electrons is reflected from the surface of specimen. Specimen is mostly coated with gold prior to observation. Here the specimen scatters many electrons whereas others are absorbed. This instrument is ideal to observe the surface view in three dimensional appearances.

Table 2.3: Differences between light and electron microscope

Light Microscope	Electron microscope
Glass lenses are used to focus the light rays	Powerful magnets are used to focus beam of electrons
Image is directly detected by naked eye	Not directly detected by naked eye, micrographs are used
Living and non living objects can be observed	Only non-living objects are observed
Actual color of the object can be observed	Actual color cannot be observed. Images are developed
Dyes used to stain the object	Heavy metals are used to stain the object

Historical background of the cell and analyses the structure and functions of the sub cellular units**Cell theory**

All organisms are composed of cells.

Recall the hierarchy of life, the levels of organization mentioned earlier. The basic unit which can be called “living” is the cell, which may form a single celled organism (e.g. *Chlamydomonas*, Yeast) or a multi-cellular plant or animal. The cell is the basic structural and functional unit of life.

The level of organization of matter represented by a cell shows all the characteristics of life. Any stage below level of a cell cannot be considered living, whether it is a single celled organism or multi-cellular plant or an animal.

Robert Hooke (1665) examined a cork using simple microscope and gave the term “CELL” to describe the basic units.

Anton Van Leeuwenhook (1650), a contemporary of Robert Hooke, was the first to describe and record living single celled organisms, *Euglena* & bacteria

Matthias Schleiden (1831), a botanist, studying plant tissues concluded that all plants are made up of cells.

Theodore Schwann a zoologist (1839) concluded that animal tissues are also made up of cells.

Rudolf Virchow (1855) showed that all cells arise from pre-existing cells by cell division,

Schleiden, Schwann and Virchow presented the 'Cell Theory' which included the following.

1. All organisms are composed of one or more cells.
2. The basic structural and functional unit of organisms is the cell.
3. All cells arise from pre-existing cells.

Organization of cells

There are two kinds of cellular organization - Prokaryotic and Eukaryotic

All cells share certain basic features. They are;

- All cells are bounded by a plasma membrane which is a selective barrier
- Within the cell have, a semifluid, jelly like substance which is called cytosol. Subcellular components are suspended within the cytosol.
- They carry DNA as genetic materials.
- Ribosomes are found in all cells

Table 2.4: The differences between Prokaryotic cells and Eukaryotic cells

Feature	Prokaryote	Eukaryote
organism	Bacteria, Archaeobacteria	Protists, Fungi, plants, animals
Cell size	Average diameter 1-5µm	10µm-100µm diameter
Form	Mainly unicellular	Mainly multicellular (except most of protista and some fungi are unicellular)
Evolutionary origin	3.5 billion years ago	1.8 billion years ago, evolved from prokaryotes
Cell division	Binary fission, no mitosis and meiosis	Mitosis, meiosis, or both;
Genetic material	DNA is circular and lies free in the cytoplasm. This region is called nucleoid, DNA is naked and not associated with proteins	DNA is linear and contained in a nucleus. DNA is associated with proteins
Type of ribosomes	70s ribosome (smaller)	Both 70s (Mitochondria and Chloroplast) and 80s ribosomes (larger) present (may attach to endoplasmic reticulum)

Organelles	Few organelles, none are surrounded by membrane Internal membranes scarces; if present usually associated with respiration, photosynthesis and N ₂ fixation.	Many organelles, membrane bounded organelles present. Great diversity of organelles. e.g. nucleus, mitochondria, chloroplasts bounded by two membranes. e.g. Lysosomes, Vacuole, bounded by single membrane.
Cell walls	Peptidoglycan present in Bacteria and cyanobacteria, polysaccharide and protein present in Archae bacteria	Cell walls of green plants and fungi are rigid and contain polysaccharides; cellulose in plant cell walls and chitin in fungal walls (none in animal cells)
Flagella	Simple, lacking microtubules; extracellular (not enclosed by cell surface membrane) 20nm diameter	Complex, with '9+2' arrangement of microtubules; intracellular (surrounded by cell surface membrane) 200nm diameter
Respiration	Mostly by mesosomes	Mitochondria for aerobic respiration
Photosynthesis	No chloroplasts; takes place on membranes which show no stacking	Chloroplasts containing membranes which are usually stacked into lamellae or grana
Nitrogen fixation	Some have the ability	None have the ability

Bacteria, Cyanobacteria and Achaea are prokaryotic cells. All the other organisms have eukaryotic cells.




Fig 2.18: Structure of an animal cell




Fig 2.19: Structure of plant cell

Structures and functions of organelles and other subcellular components

Plasma membrane: Plasmamembrane is the outer limit of cytoplasm. All cellular membranes resemble the ultra structure of plasma membrane. In 1972, Singer and Nicolson put forward the fluid mosaic model of cell membrane. It is mainly composed of;

- Phospholipids (most abundant type of lipid in plasma membrane)
- Protein

The Plasma membrane has the following features. It is about 7nm thick. It is mainly made up of a phospholipid bilayer. Phospholipids are amphipathic molecules. The hydrophilic heads of the phospholipids face outwards into the aqueous environment of both inside and outside of the cell.

The hydrophobic hydrocarbon tails face inwards and create a hydrophobic interior.

Plasmamembrane is compared to the fluid mosaic model. Since phospholipid molecules are moveable, they provide the fluid nature to the membrane.

Protein molecules embedded randomly contribute to its mosaic nature.

Some of the protein molecules penetrate all the way through the membrane, called transmembrane proteins and some others penetrate only part of the way into the membrane. These are called **integral proteins**.

Most of the integral proteins are transmembrane proteins which have hydrophilic channels. These act as pores through which ions and certain polar molecules can pass.

Some proteins are not embedded in the lipid bilayer at all, and are loosely bound to the inner surface of the membrane, called **peripheral proteins**.

Some proteins and lipids have short branching carbohydrate chains like antennae, forming **glycoprotein** and **glycolipids**, respectively.

Animal's cell membrane may contain few **cholesterol molecules** randomly integrated into the lipid bilayer.

These cholesterol molecules provide flexibility and stability to the membrane by reducing membrane fluidity at moderate temperatures and prevent membrane solidification at low temperatures.

The two sides of the membrane may differ in composition and function.




Fig 2.20: Structure of the plasma membrane

Functions

- The plasma membrane surrounds the cytoplasm of living cell physically separating the intracellular components from the extracellular environment.
- Plasma membrane is selectively permeable and able to regulate the exchange of material needed for survival.
- Proteins embedded in the plasma membrane identify the cell, enabling nearby cells to communicate with each other (involved in cell recognition).
- Some protein molecules act as receptor molecules for interacting with specific biochemical, such as hormones, neurotransmitters and immune proteins.
- Some proteins in the cell membrane attach to some cytoskeletal fibers and help

to maintain the shape of the cell.

- Some proteins in the membrane act as enzymes. (e.g. Microvillus on epithelial cell lining of some parts of the gut contains digestive enzymes in their cell surface membrane.)

Subcellular components

There are many sub-cellular components in the cell. Some of them are organelles, which are bound by membranes and suspended in the cytosol of eukaryotic cell to perform specialized functions.

Nucleus

Most prominent organelle, consist most of the genes, having an average diameter of $5\mu\text{m}$ and enclosed by a double membrane cover called nuclear envelope.

- Nuclear envelope- composed of two membranes, inner and outer membranes, separated by a space of 20-40 nm. Nuclear envelope is perforated by nuclear pores which has pore complex to regulate the entry and exit of substances. It has nuclear lamina, made up of protein filaments which line the interior side of the nuclear envelope.
- Nuclear matrix is made up of protein filaments and extended throughout the interior of the nucleus. Chromatin and nucleolus are embedded in the nuclear matrix.
- Nucleolus- appears as darkly stained granules with fibers adjoining part of the chromatin.
- Chromatin –appears as a diffused mass in electron micrographs of non dividing cells. It is a complex of DNA and proteins. During nuclear divisions, chromatin condenses, tightly coils and form threads, called chromosomes. Each species has a constant number of chromosomes. (e.g. typical human cell has 46 chromosomes).

Functions

- Control all cellular activities.
- Synthesize DNA to produce new nuclei for cell divisions.
- Synthesize rRNAs and ribosomal subunits required for protein synthesis, through nucleolus.
- Synthesize mRNA and tRNA according to the information present on the DNA.
- Store and transport genetic information.

Ribosomes

These are subcellular components which carry out protein synthesis. They consist of two subunits; larger subunit and smaller subunit. They are composed of rRNA and protein. Ribosomes are found in two types; 70S and 80S. 70S ribosomes are found freely in the cytoplasm of prokaryotes, matrix of mitochondria and stroma of chloroplasts. 80S ribosomes are found only in eukaryotes. Based on the nature of presence, 80S ribosomes are categorized as two types; free ribosomes and bound ribosomes.

Free ribosomes: freely available as group in cytoplasm. Bound ribosomes are attached to the membrane surface of rough endoplasmic reticulum.

Functions

Protein synthesis

Endoplasmic reticulum

It is a network of internal membranes forming flattened or tubular sacs separating cytosol from ER lumen. It is continuous with the outer membrane of nuclear envelope. There are two types of ER; Rough ER and Smooth ER

Rough ER

Rough ER consists of flattened sacs, and ribosomes bound to surface. Proteins synthesized by ribosomes move into lumen of ER.

Functions

- Transport protein synthesized by ribosomes
- Synthesizing glycoproteins
- Produce transport vesicles
- Facilitate the growth of own membrane by adding phospholipids proteins and carbohydrates. Therefore called as membrane factory

Smooth ER

Smooth ER is a network of tubular sacs without ribosomes. Membrane bound enzymes are present.

Functions

- It synthesizes lipids including oils, steroids and phospholipids.
- Metabolism of carbohydrates.
- Produce transport vesicles to transport within cell.
- Involves in detoxification.
- Stores Ca^{2+} ions.




Fig 2.21: structure of the endoplasmic reticulum

Golgi apparatus




Fig 2.22: Structure of the golgi apparatus

Golgi apparatus is a stacks of flattened sacs or Cisternae. Inner and outer surfaces can be identified as cis face and transface respectively. Cis face is located near the E.R to receive vessicles from E.R. Trans face give rise to secretory vessicles which budded off and travel other side. Golgi complex is abundant in secretory cells.

Functions

- Collecting, packaging and distribution of materials
- Manufacturing cellulose and non cellulose cell wall components such as pectin
- Produce lysosomes

Lysosomes

They are single membrane bounded vesicles contributing to digestive activity. They contain hydrolytic enzymes which catalyze breakdown of carbohydrates, proteins, lipids and nucleic acids.

Functions

- Digest food particles received by phagocytosis
- Transport residue material out of cell by exocytosis.
- Digest worn out organelles
- Autolysis causing cell death.

Peroxisome

They are single membrane bounded vesicles with oxidizing enzymes. They are present in both plants and animals. Enzymes in peroxysome catalyze the breakdown of H_2O_2 .

Functions

- Detoxification of peroxides
- Photorespiration in plants

Specialized peroxysomes called glyoxysomes are found in fat storing tissues in plants. Glyoxysomes converts fatty acids into sugar.

Mitochondria

It is one of the most common organelles in eukaryotic cells. It is an elongated organelle with two enclosing membranes. Outer membrane is smooth but the inner membrane is convoluted to form cristae. Cristae increase the surface area and they contain stalk particles. The gap/space in between inner and outer membranes of the mitochondrion is called intermembrane space. The inner most part of the organelle is known as mitochondrial matrix, which consists of 70 s ribosomes circular DNA

molecule (mitochondrial DNA), phosphate granules and enzymes. The matrix carries enzymes for the reactions in Krebs cycle (in cellular respiration). Further, cristae composed of proteins and enzymes essential for electron transport chain and oxidative phosphorylation.

Functions

- Synthesize ATP in aerobic respiration
- Involve in Photorespiration




Fig 2.23: Structure of the mitochondria

Chloroplast

It is a biconvex lens shaped organelle with two membranes which is found in plants and some protists. The outer and inner membranes are smooth and are separated by a very narrow intermembrane space. Inside the chloroplast there is another membrane system. This membrane produces flattened and interconnected sacks called thylakoids. Thylakoids contain complexes called photosystems which are made up of photosynthetic pigments. Thylakoids stacked to form a granum. The grana are interconnected by inter granal lamellae. The fluid outside the thylakoid is stroma which contain circular DNA (chloroplast DNA), 70s ribosomes, many enzymes, starch granules and lipid droplets.




Fig 2.24: Structure of the chloroplast

Functions

- Photosynthesis

Cytoskeleton

Cytoskeleton is the supporting structure of the cell and maintains its shape. It is more important for animal cells which lack cell walls. Cytoskeleton is made out of microtubules and protein filaments. Additionally, it is Dynamic hence, has the ability to break and reform as needed.

There are three types of components in the Cytoskeleton as follows;

- Microtubules
- Actin filaments or Microfilaments,
- Intermediate filaments

Table 2.5: Differences between Microtubules, Microfilaments and intermediate filaments

Property	Microtubules (Tubulin polymers)	Microfilaments (Actin filaments)	Intermediate filaments
Structure	Hollow tubes; wall consists of 13 columns of tubulin molecules	Two intertwined strands of actin, each strand is a polymer of actin subunits	Fibrous proteins supercoiled into thicker cables
Protein subunits	Tubulin	Actin	One of several different proteins (e.g. Keratin), depending on the cell type.
Main functions	Maintenance of cell shape Cell motility (as in cilia or flagella) Chromosome movements in cell division Organelle movements	Maintenance of cell shape (tension-bearing elements) Changes in cell shape Muscle contraction Cytoplasmic streaming in plant cells Cell motility (as in pseudopodia) Cell division in animal cells (cleavage furrow formation)	Maintaining of cell shape (tension-bearing elements) Anchorage of nucleus and certain other organelles. Formation of nuclear lamina

Functions

- Provide strength to the cytoplasm
- Anchorage organelles and cytosolic enzymes of the cell
- Movement of cytoplasm, cytoplasmic streaming, positioned organelles and move chromosomes when necessary.
- Maintain the shape of the cell (mainly in animal cells)

Cilia and Flagella

Cilia and flagella share a common structure. Flagella are long elongated structures and Cilia are short cellular projections that are often organized in rows. Cilia are more numerous than flagella on the cell surface. They are made of microtubules, with a 9+2 structure (Nine doublets of microtubules are arranged in a ring, with two single microtubules in its center). They are covered by plasma membrane and bound to a basal body which anchors the cilium or flagellum to the cell. The Basal body has 9 + 0 arrangement (no microtubules in its center)

Functions

- Act as locomotor appendages
- Can move fluid over the surface of the tissue
- Cilia lining in oviducts help move an egg toward the uterus




Fig 2.25: Structure of the Cilium

Centrioles

Centriole is made up of cylindrically arranged microtubules which are non membrane bounded subcellular component present only in animal cells. Each centriole composed of nine sets of triplet microtubules arranged in a ring (9+0). A pair of centrioles which arranged perpendicular to each other are located in a region called centrosome near the nucleus.

Functions

- Produce aster and spindle in cell division

Central Vacuole

Central vacuole is a large structure, bound by tonoplast, filled with liquid called cell sap found in plant cells. The composition of sap differs from cytosol and it contains water, ions such as Potassium and Chloride and sometimes water soluble colored pigments such as anthocyanin.

Functions

- Stores water and other materials such as sugars, ions and pigments.
- Maintains water balance of the cell
- Gives turgidity and support to cell.
- Produce colours in some plants with sap pigments
- Stores soluble substances needed for cellular activities.

Extracellular components**1. Cell wall**

Cell wall is an extracellular structure of plant cells. Animal cells do not have cell walls. However, prokaryotes, fungi and some protists also have a thin and flexible cell wall. The chemical composition of the wall greatly varies from species to species and even from one cell type to another even in the same plant. Nevertheless in Plants, cell wall is generally made up of cellulose, pectin, hemicellulose, lignin and suberin (in some plant cells only).

Plants generate two types of cell walls: primary and secondary walls. Young cells first secrete primary cell wall: it is the wall laid down during plant cell division.

Just outside the primary wall there is a thin layer (middle lamella) which is rich in sticky polysaccharides called pectins (magnesium and calcium pectate). Middle lamella glues adjacent cells together. Due to the deposition of hardening substances on the primary wall a secondary cell wall is generated secondarily.

Primary cell wall is permeable, relatively thin, flexible, composed mainly of cellulose fibers which are laid unevenly running through the extracellular matrix (middle lamella). Water can move freely through the free spaces of cell wall.

Secondary cell wall lies between plasma membrane and primary cell wall. It contains several layers of hard materials, forming a rigid structure. In addition to cellulose, impermeable substances such as lignin and suberin are also incorporated into the secondary wall. Lignin cement anchors cellulose fibers together providing hard and rigid matrix, giving the cell wall an extra support.

Cell wall has pits through which cytoplasm of adjoining cells join through plasmodesmata.

Functions

- Protection and support
- Allows development of turgidity when water enters the cell
- Prevents bursting during turgidity
- Limits and control cell growth
- Component of appoplast pathway
- Maintaining cell shape
- hold the plant up against the force of gravity

2. Cell junctions

Cell junctions are structures at which neighbouring plasma membranes are joined. They are also interact and communicate via sites of direct physical contacts.

Functions

- Connects the internal chemical environment of adjacent cells.
- Cell junctions are structures at which cytoplasm of adjoining cells are joined. There are three types of cell junctions in animal cells
- Tight junctions – connect the plasma membranes of adjacent cells tightly bound by specific proteins forming continuous seals around the cells. Prevent leakages of extracellular fluids through intercellular space. e.g. skin epithelium
- Desmosomes/Anchor junctions – mechanically attach the cytoskeletons of adjoining cells by intermediate filaments for strong binding.e.g. muscle tissue
- Gap junctions /Communicating junctions – provide cytoplasmic channels from one cell to an adjacent cell. Gap junctions consists of special membrane proteins that surround the pore through which ions, sugars amino acids may pass. They allow signal and material exchange between adjacent cells through direct connections. e.g.heart muscles, animal embryo.

Plasmodesmata

- Microscopic channels which runs through plant cell walls. They are cytoplasmic living connections between cytoplasm of adjoining cells. These are membrane lined channels filled with cytoplasm.

Extracellular matrix of animal cells

Although animal cells lack cell walls they do have elaborate extracellular matrix (ECM). Main components of the ECM are glycoproteins and other carbohydrates containing molecules secreted by the cells. Most abundant glycoprotein in the ECM of most animal cell is collagen which forms strong fibres outside the cell. The collagen fibres are embedded in a network woven out of proteoglycan secreted by cells.

Functions

- Forms a protective layer over the cell surface
- Linking extra cellular matrix and cytoskeleton.
- Influences the cell behavior by Involving in the mechanical and chemical signaling.

The cell cycle and the process of cell division

The sequence of events that takes place in the cell from the end of one cell division to the end of the next cell division is referred to as cell cycle. At the end of the cell division, two genetically identical daughter cells resembling the parent cell are produced in mitosis.




Fig 2.26: The cell cycle

Eukaryotic cell cycle

Mitosis

Eukaryotic cell cycle may divided into two major phases.

- Interphase
- Mitotic phase/ M-phase

Interphase is the longer phase of cell division. It covers about 90% of the cell cycle. Interphase could be divided into three phases;

- G₁ phase (first gap phase)
- S phase (synthetic phase)
- G₂ phase (second gap phase)

G₁ phase

In this phase synthesis of proteins and production of cellular organelles leading to cell growth occur. Proteins essential for S phase are produced during this phase.

S phase

DNA replication occurs and synthesis of histone proteins takes place. DNA wind around histone beads and form chromatin.

G₂ phase

Cells continue to grow through protein synthesis as well as cellular organelles. Proteins essential for mitotic phase will be synthesized. Duplication of centrosomes takes place.

There are cell cycle-controlling checkpoints available at G₁, G₂ and M phases to ensure that the cell is ready for moving into upcoming phases of cell division. Some cells receive a go-head signal at the G₁ check point, it will usually complete the G₁, S, G₂ and M phases and divide. If it does not receive a go head signal at that point it may exit the cycle, entering into a non dividing stage called the Go phase. The most cells of the human body are actually in the Go phase. e.g. nerve cells and muscle cells.

Mitotic phase/ M phase

M phase covers only about 10% of cell cycle. This includes mitosis and cytokinesis.

Mitosis

Mitosis is referred to the nuclear division which gives rise to two genetically identical daughter nuclei from a mother nucleus. This may get divided into five stages; prophase, prometaphase, metaphase, anaphase and telophase in order to ease the learning of activities of cell cycle.

1. Prophase

Chromatin fibers get condensed by shortening and thickening and transformed into chromosomes. As a result chromosomes will be visible through light microscope. Nucleoli get disappeared and chromosomes appear with two sister chromatids attached at the centromere. Chromosomal arms of sister chromatids attached by special proteins called cohesion. The formation of mitotic spindles begins. Spindle includes the centrosomes, the spindle microtubules and the aster.

Centrosomes move toward opposite poles of the cell due to the lengthening of microtubules between them.

2. Prometaphase

The nuclear envelope fragments. Chromosomes get even more condensed. A special protein called kinetochore attaches the sister chromatids of each chromosome at their centromere. Some of the microtubules that attach to the kinetochore of the chromosomes move the chromosomes back and forth. Microtubules which are not attached to the kinetochore interact with those from the opposite poles.

3. Metaphase

Centrosomes reach the opposite poles. The chromosomes have arrived to a place called metaphase plate which is located in equal distance from each pole. The centromeres of all chromosomes are located in the metaphase plate. At the end of this phase, each chromosome of the cell get attached to the kinetochore microtubule at their centromere and aligned at the metaphase plate.

4. Anaphase

Sister chromatids are separated at the centromere. Microtubules attached to kinetochore get shorten and pull sister chromatids towards the opposite poles. Cell elongates as the non kinetochore microtubules are lengthen. By the end of anaphase equal and complete set of chromosomes found at each pole of the cell.

5. Telophase

Nuclear envelope reforms around each set of chromosomes at opposite poles. Nucleoli reappears. Spindle microtubules get depolymerized. Chromosomes unwind and become less condense to form chromatin. Two genetically identical daughter nuclei are formed.

Cytokinesis

The division of the cytoplasm starts at the end of the telophase. Therefore at the end of the mitosis two genetically identical daughter cells are produced.

In animal cells- a cleavage furrow forms. This produces two genetically identical daughter cells.

In plant cells- cell plate forms as a result of vesicle produced by golgi apparatus. This divides the cytoplasm in to two and generates two genetically identical daughter cells to the parent cell.

Significances of mitosis

1. Maintains the genetic stability
2. Growth and development
3. Cell repair, replacement and regeneration
4. Asexual reproduction




Fig 2.27 : The phases of mitotic cell cycle

Meiosis

Sexually reproducing organisms undergo different type of cell division called meiosis.

Meiosis

Meiosis is a type of nuclear division which gives rise to four haploid, genetically non identical daughter nuclei, from a diploid mother nucleus.

Meiosis involves two consecutive nuclear divisions, Meiosis I and Meiosis II.

Meiosis I is a reduction division and Meiosis II is similar to mitosis, each stage consists of four sub-phases: prophase, metaphase, anaphase, and telophase.

Before meiosis one cell is in interphase, during S phase of the interphase DNA replication occur.

Meiosis I

1. Prophase I

Cell enters to the prophase from interphase. Chromosomes begin to condense. Nucleolus begins to disappear. Next the formation of zipper like structure called the synaptonemal complex by a specific proteins holds two homolog tightly together. The pairing and physical connection of homologous chromosomes is called synapsis.

During synapsis part of the DNA molecule of non-sister chromatids paired homologous chromosomes break, exchange and rejoin at corresponding point. This process is called crossing over. These points of crossing over become visible as chiasmata after the synaptonemal complex disassembles and the homologous chromosomes slightly apart from each other

Nuclear envelope breaks. Centrosomes move towards opposite poles forming spindle in animal cells.

The kinetochore of each homologue attach to microtubule from one pole or the other. The homologous pair then moves toward the metaphase plate.

2. Metaphase I

The pair of homologous chromosomes get arranged on the metaphase plate with one chromosome of each pair faces each pole. Both chromatids of a homologue are attached to kinetochore microtubules from one pole and those of the other homolog are attached to kinetochore microtubules from the opposite pole. Homologous chromosome arrange randomly at metaphase plate.

3. Anaphase I

Kinetochore microtubules of the spindle get shorten. Homologous pair separates and one chromosome of each pair moves towards the opposite pole. Sister chromatids of each chromosome remain attached at the centromere and move as a single unit towards the same pole.

4. Telophase I

One complete haploid set of chromosomes accumulate at each pole. Nuclear envelope reforms around each set of chromosomes. Nucleoli reappear. Spindle disintegrates. Chromosomes decondensed into chromatin. Genetically non identical, haploid, two daughter nuclei are formed within one cell.

Cytokinesis

Usually occurs simultaneously with telophase I. Genetically non identical, haploid, two daughter cells are formed. In animal cells, cleavage furrow is formed. In plant cells a cell plate is formed.

No DNA replication occurs between meiosis I and meiosis II

Meiosis II**1. Prophase II**

Centrosomes start producing spindle apparatus (spindle fibers, aster centrosome). Chromatin fibers condense and produce chromosomes with two sister chromatids. Nuclear envelope breaks down into fragments. Nucleolus disappears. During the late prophase II centromere of the chromosomes are moved to the metaphase II plate.

2. Metaphase II

All Chromosomes get attached to the microtubules at their centromere and aligned on the metaphase plate. Kinetochores of sister chromatids are attached to microtubules extending from both poles.

Due to the crossing over in meiosis I, the two sister chromatids of each chromosome are not genetically identical.

Meiosis II usually takes place in the perpendicular direction of Meiosis I. Therefore, metaphase plate of meiosis II is perpendicular to the metaphase plate of meiosis I.

3. Anaphase II

Due to the breakdown of proteins attaching sister chromatids, they are separated at centromere. As a result of shortening of microtubules, sister chromatids of each chromosome move towards opposite poles.


4. Telophase II

Nuclear envelope and nucleolus reform. Chromosomes decondense into chromatin. Spindle disassembles. Genetically non identical, haploid, four daughter nuclei are formed from one parent cell.

Cytokinesis


Cytokinesis occurs as in mitosis. Genetically non identical, haploid, four daughter cells are formed. These four daughter cells are not even identical to their parent cell.

Centrosomes or centrioles are not available in plant cells. However, spindle is formed during cell division from accumulated microtubule complex.




**Meiosis I
Prophase I**

**Meiosis I
Metaphase I**



Meiosis I
Anaphase I

Meiosis I
Telophase I
and cytokinesis



Meiosis II
Prophase II

Meiosis II
Metaphase II




Fig 2.28: The phases of meiosis

Significance of meiosis

- Maintains the constant number of chromosomes through generations in sexually reproducing species.
- Produce new genetic variations leading to evolution.
- Genetic variation occurs due to crossing over ,recombination and independent assortment.

Tumor, cancer and galls

- Cell division is driven by external and internal factors. They may be chemical or physical factors
- Cancer cells do not respond to normally to the body's control mechanism
- They divide excessively and invade other tissues. If unchecked they can kill the organism.

- Cancer cells do not consider the normal signals that regulate the cell cycle.
- They do not need growth factors. They may make required growth factors themselves or giving signals to continue cell cycle without growth factors. Another possibly is an abnormal cell cycle control system.
- The problem begins when a single cell in a tissue undergoes transformation, the process converts a normal cell to abnormal cell.
- If the body immune system can not recognize and destroy it, it may leads to proliferation of cells and formation of a tumor.
- If the abnormal cells remain at the original site, the lump is called benign tumor. Most benign tumors do not cause serious problems and can be completely removed by a surgery.
- A malignant tumor becomes invasive and attack one or more organs. An individual with a malignant tumor is said to have a cancer.
- A few tumor cells may separate from the original tumor, enter blood vessels or lymph vessels and travel to other parts of the body. They may proliferate and form a new tumor.
- This spread of cancer cells to locations distant from their original site is called metastasis.

Galls in plants

- This occurs due to uncontrolled mitotic division of plant cell.
- The plant cell division is controlled by maintaining a proper balance between plant growth regulators such as auxins and cytokinins. When this balance is lost plant cells produce undifferentiated mass of cells.
- Galls are the bumps and growths that develop on different parts of plants after being invaded by some very unique organisms.
- Galls have range of causes, including viruses, fungi, bacteria, insects and mites.
- Usually the gall causers in some way attack or penetrate the plants growing tissues and causes the host to reorganize its cells and to develop an abnormal growth.

The energy relationships in metabolic processes

Sum of all biochemical reactions of living being is known as the metabolism and it consists of all catabolic and anabolic reactions.

Catabolism is breaking down of complex molecules into simple molecules by releasing free energy. Therefore it is an exergonic reaction. Anabolism is making complex molecules from the simple molecules by absorbing free energy. Hence it is an endergonic reaction.

Biochemical reactions involved in usage of energy released by catabolic reactions in

living system are called as anabolic reactions. ATP acts as the energy carrier in all living organism including the simplest bacteria. Therefore the ATP is known as the universal currency of energy transactions.

Energy can be defined as the capacity to do work. All living organisms require energy for their living process in many ways. Such processes are;

- Synthesis of substances
- Active transport across plasma membrane
- Transmission of nerve impulses
- Muscle contraction
- Beating of cilia and flagella
- Bioluminescence
- Electrical discharges.

Overall idea of the energy relations of living system on biosphere is composed of following steps.

- Energy flows into biological systems from the environment through solar radiation. (Primary energy source is the Sun)
- Light energy is captured in the cells having photosynthetic pigments (chlorophyll) by the process of photosynthesis and stored as chemical energy in the organic compounds such as carbohydrates
- Captured energy in organic food is transformed into chemical energy in ATP by a process called cellular respiration.
- The energy stored in ATP is utilized in various energy requiring processes.

ATP (Adenosine Tri Phosphate)

ATP is a nucleotide, consisting of,

- Ribose- sugar
- Adenine - nitrogenous base
- A chain of three phosphate groups.

During the hydrolysis of ATP, ADP and Pi are produced. As a result, a very high energy is released. This is because the reactants (ATP and water) contain more energy in comparison to products (ADP and Pi). Therefore it yields energy and is an exergonic reaction.

When ATP is hydrolyzed, the free energy yield of each of the two end phosphate groups is -30.5kJ/mol.




Fig 2.29: Chemical structure of ATP molecule (need not be memorized)

Most biological reactions use the energy released during breaking of the terminal phosphate bond. ATP is mobile. Therefore it can carry energy to anywhere in the cell, for any energy consuming reaction.

ATP can be produced within living cells within a short period of time, using ADP, inorganic phosphate (Pi) and energy. Production of ATP within cells is called phosphorylation. According to the energy source phosphorylation is divided as;

- | | | |
|--------------------------------|--|---------------------------|
| i. Photophosphorylation | – synthesis of ATP using solar energy in photosynthesis | |
| ii. Substrate phosphorylation | – synthesis of ATP using energy released by the breaking down of complex molecules into simple ones. | } In cellular respiration |
| iii. Oxidative phosphorylation | – synthesis of ATP using energy released as a result of oxidation of molecules. | |

In living cells energy in ATP is transformed into various energy forms which are used for different functions.