

68

Total Training Days

Live Edit semantic models in
Direct Lake mode

Select a semantic model in Direct Lake mode
and begin editing it using Power BI Desktop.

Consistency & Progress: A Data-Driven Retrospective

An analysis of 2.5+ years of personal training data
(842 logged sets across 92 exercises), covering
training consistency, muscle group balance, and lift-
by-lift progress.

842

Total Sets Logged

123

Longest Gap

90

Exercises Tracked


Sets by

Live Edit semantic models in
Direct Lake mode

Select a semantic model in Direct Lake mode
and begin editing it using Power BI Desktop.

Volume by Muscle Group (side by side)


Training Consistency Over Time



65% of months had 0-1 training days. A 123-day gap in late 2025 preceded a sustained comeback in early 2026, with July 2026 nearly matching the peak of the initial 2024 training block.

Sets by Muscle Group

Training Volume by Muscle Group




Live Edit semantic models in Direct Lake mode

Select a semantic model in Direct Lake mode and begin editing it using Power BI Desktop.

Sets by Muscle Group (side by side)

Sets by Muscle Group




Shoulders led in training frequency (129 sets, #1) but ranked only 4th in total training volume. Lats trained less often (68 sets, 4th) but carried the heaviest real load by a wide margin, revealing that frequency alone doesn't capture true training intensity.

Live Edit semantic models in Direct Lake mode

Select a semantic model in Direct Lake mode and begin editing it using Power BI Desktop.

Methodology

Exercise Progress Summary



Of 10 exercises with sufficient recent data, 2 are actively progressing and 2 show a genuine decline, not just a plateau, distinguished by tracking both weight and rep changes over a rolling 120-day window.

Exercise	Status	Peak Weight (Before)	Peak Weight (Recent)
Lat Pulldown (Cable)	progressing	55	55
Shoulder Press (Dumbbell)	plateaued	15	15
Shrug (Dumbbell)	progressing	10	15
Bulgarian Split Squat (Dumbbell)	plateaued	10	10
Dumbbell Row	plateaued	10	10
Hip Thrust (Barbell)	regressing	50	10
Reverse Curl (Dumbbell)	plateaued	10	10
Triceps Extension (Dumbbell)	plateaued	10	10
Bicep Curl (Dumbbell)	regressing	10	7
Lateral Raise (Dumbbell)	plateaued	7	5

Methodology Notes: Data cleaned and analyzed in Python (Pandas), with muscle group classifications sourced from Hevy's official Exercise Template API. Two data quality issues were identified and excluded: a 5-minute session with 84 logged sets (confirmed as app exploration, not real training) and a single-session weight outlier inconsistent with surrounding data (confirmed data entry error). Progress classification requires a minimum of 2 sessions within the recent 120-day window to draw a conclusion; exercises with insufficient recent data are excluded rather than forced to a verdict.

Overall Analysis & Recommendation

Across 2.5+ years and 842 logged sets (roughly April 2024 through January 2026 beginning March 2026 that has held for comeback reflects a rebuilt habit rather

Live Edit semantic models in Direct Lake mode

Select a semantic model in Direct Lake mode and begin editing it using Power BI Desktop.

strong, structured start (Jan-Mar 2024), a long, inconsistent middle stretch (Apr-Sep 2024), a sharp recovery (Oct-Dec 2024), and a genuine, sustained comeback (Jan-Mar 2025). The volume nearly matches the original 2024 peak, suggesting this

What's working:

- Lats and quadriceps are the best-trained muscle groups by real volume, not just frequency, with Lat Pulldown showing sustained progressive overload
- Shrug is also actively progressing, with a clean weight increase in recent sessions
- The current March-July 2026 stretch is the strongest sustained consistency in the entire dataset
-

What's worth addressing:

- Shoulders lead in total sets (129, #1) but rank only 4th in actual training volume, indicating lighter relative loading despite high frequency
- Hip Thrust (Barbell) and Bicep Curl show genuine declines rather than plateaus; Hip Thrust's drop (50 lbs to 10-15 lbs) may reflect a deliberate technique correction, while Bicep Curl's decline has no clear cause in the data
- Hamstrings and forearms are the most under-trained muscle groups by both frequency and volume
- 6 of 10 evaluable exercises are currently plateaued (no new weight or rep record in the last 120 days), suggesting a program-level adjustment (rep range, tempo, or exercise variation) may be more effective than isolated tweaks

Recommendations:

- Protect the current streak with planned lighter weeks every 4-6 weeks rather than waiting for a forced break
- Add a hamstring-focused and forearm-focused movement to close the clearest muscle-balance gap
- Introduce a deliberate variable (grip, tempo, rep range) across plateaued lifts rather than repeating the same numbers
- Confirm whether the Hip Thrust weight drop reflects a technique reset, so it isn't misread as regression going forward

Note: these observations are drawn directly from the logged training data and are not professional training or medical advice.