

68

Sum of training_days

Training Consistency & Progress: A Data-Driven Retrospective

123

LongestGap


842

Count of exercise_title

An analysis of 2.5+ years of personal training data
(842 logged sets across 92 exercises), covering
training consistency, muscle group balance, and lift-
by-lift progress.


90

Count of exercise_title




65% of months had 0-1 training days. A 123-day gap in late 2025 preceded a sustained comeback in early 2026, with July 2026 nearly matching the peak of the initial 2024 training block.

Training Volume by Muscle Group




Sets by Muscle Group



Shoulders led in training frequency (129 sets, #1) but ranked only 4th in total training volume. Lats trained less often (68 sets, 4th) but carried the heaviest real load by a wide margin, revealing that frequency alone doesn't capture true training intensity.

Exercise Progress Summary



Shoulders led in training frequency (129 sets, #1) but ranked only 4th in total training volume. Lats trained less often (68 sets, 4th) but carried the heaviest real load by a wide margin, revealing that frequency alone doesn't capture true training intensity.

Exercise	Status	Peak Weight (Before)	Peak Weight (Recent)
Lat Pulldown (Cable)	progressing	55	55
Shoulder Press (Dumbbell)	plateaued	15	15
Shrug (Dumbbell)	progressing	10	15
Bulgarian Split Squat (Dumbbell)	plateaued	10	10
Dumbbell Row	plateaued	10	10
Hip Thrust (Barbell)	regressing	50	10
Reverse Curl (Dumbbell)	plateaued	10	10
Triceps Extension (Dumbbell)	plateaued	10	10
Bicep Curl (Dumbbell)	regressing	10	7
Lateral Raise (Dumbbell)	plateaued	7	5
Total		187	147

Methodology Notes: Data cleaned and analyzed in Python (Pandas), with muscle group classifications sourced from Hevy's official Exercise Template API. Two data quality issues were identified and excluded: a 5-minute session with 84 logged sets (confirmed as app exploration, not real training) and a single-session weight outlier inconsistent with surrounding data (confirmed data entry error). Progress classification requires a minimum of 2 sessions within the recent 120-day window to draw a conclusion; exercises with insufficient recent data are excluded rather than forced to a verdict.

Overall Analysis & Recommendations

Across 2.5+ years and 842 logged sets, this data tells a clear three-act story: a strong, structured start (Jan-Mar 2024), a long, inconsistent middle stretch (roughly April 2024 through January 2026, with 65% of months at 0-1 training days and one 123-day gap), and a genuine, sustained comeback beginning March 2026 that has held for five consecutive months. July 2026's training volume nearly matches the original 2024 peak, suggesting this comeback reflects a rebuilt habit rather than a short-lived burst of motivation.

What's working:

- Lats and quadriceps are the best-trained muscle groups by real volume, not just frequency, with Lat Pulldown showing sustained progressive overload
- Shrug and Seated Row are also actively progressing, with clean weight increases in recent sessions
- The current March-July 2026 stretch is the strongest sustained consistency in the entire dataset

What's worth addressing:

- Shoulders lead in total sets (129, #1) but rank only 4th in actual training volume, indicating lighter relative loading despite high frequency
- Hip Thrust (Barbell) and Bicep Curl show genuine declines rather than plateaus; Hip Thrust's drop (50 lbs to 10-15 lbs) may reflect a deliberate technique correction, while Bicep Curl's decline has no clear cause in the data
- Hamstrings and forearms are the most under-trained muscle groups by both frequency and volume
- 6 of 10 evaluable exercises are currently plateaued (no new weight or rep record in the last 120 days), suggesting a program-level adjustment (rep range, tempo, or exercise variation) may be more effective than isolated tweaks

Recommendations:

- Protect the current streak with planned lighter weeks every 4-6 weeks rather than waiting for a forced break
- Add a hamstring-focused and forearm-focused movement to close the clearest muscle-balance gap
- Introduce a deliberate variable (grip, tempo, rep range) across plateaued lifts rather than repeating the same numbers
- Confirm whether the Hip Thrust weight drop reflects a technique reset, so it isn't misread as regression going forward

Note: these observations are drawn directly from the logged training data and are not professional training or medical advice.