

Calorimetry

Lab: 01

Jaden Moore

Orange Coast College
Physics A285L
September 18th, 2021

1 Introduction

In this lab, we measure the final temperature of a system consisting of a hot solid submerged in cold water after reaching thermal equilibrium. We then compare the experimental equilibrium temperature with the theoretical temperature to obtain a percent error between the two.

2 Calorimetry

Consider the experiment provided by Physlet® Physics Exploration 19.3: Calorimetry. Recall that when two systems at different temperatures come in thermal contact, they will eventually become equal in temperature. That is, the heat lost by the solid block is equal to the heat gained by the water since thermal energy travels from hot to cold. Thus, the theoretical temperature can be calculated such that,

$$T_{f-theory} = \frac{m_w c_w T_{iw} + m_b c_b T_{ib}}{m_w c_w + m_b c_b} \quad (1)$$

Where m_w , m_b is the mass of the water and block respectively; c_w , c_b is the specific heat capacity of water and block respectively; and T_{iw} , T_{ib} is the initial temperature of the water and block respectively.

From this we can calculate the theoretical equilibrium temperature T_f . For example, for a system consisting of $m_b = 1$ kg, $m_w = 10$ kg, $T_{ib} = 500$ K, and $T_{iw} = 300$ K. We get that,

$$T_{f-theory} = \frac{(10 \text{ kg})(4186 \text{ J/(kg } ^\circ\text{C)})(300 \text{ K}) + (1 \text{ kg})(390 \text{ J/(kg } ^\circ\text{C)})(500 \text{ K})}{(10 \text{ kg})(4186 \text{ J/(kg } ^\circ\text{C)}) + (1 \text{ kg})(390 \text{ J/(kg } ^\circ\text{C)})} \quad (2)$$

$$T_{f-theory} = 301.8 \text{ K}$$

We then calculate the percent error between the experimental temperature gathered from the experiment and the theoretical value gathered from equation (1) above, such that

$$\text{percent error} = \left(\frac{T_{f-exp} - T_{f-theory}}{T_{f-theory}} \right) 100 \quad (3)$$

$$\text{percent error} = -0.0153 \%$$

Using the approach above, we put into a table below, the experimental equilibrium temperatures from multiple experiments along with the corresponding theoretical temperatures and percent error between the two.

Table 1: Equilibrium Temperatures at different masses of the solid block				
$m_b[kg]$	$T_{ib}[K]$	$T_{f-exp}[K]$	$T_{f-theory}[K]$	%error
1	500	301.8	301.85	-0.0153
1	750	304.2	304.15	0.0152
1	1000	306.5	306.46	0.0126
2	500	303.7	303.66	0.0137
2	750	308.2	308.23	-0.0103
2	1000	312.8	312.80	-0.0016
3	500	305.4	305.44	-0.0125
3	750	312.2	312.24	-0.0114
3	1000	319.0	319.03	-0.0104

From the table, we notice that as the mass and initial temperature of the solid block increases, the final equilibrium temperature of the system increases. This makes intuitive sense based on the equation of heat energy $Q = mc\Delta T$ which shows that the heat energy produced by an object is directly proportional to the mass of the object times the change in temperature.

3 Conclusion

From this lab, we get that two systems that come in thermal contact with each other will eventually become in thermal equilibrium with each other. Furthermore, we get that the final temperature will be smaller or greater depending on the initial temperatures of the two systems. During this experiment we did not encounter any difficulties, however we gained a greater understanding of how to apply the rules of thermal energy and the zeroth law of thermal dynamics to predict the equilibrium temperature of a system.