This article was downloaded by: [132.203.227.61] On: 28 April 2021, At: 07:25

Publisher: Institute for Operations Research and the Management Sciences (INFORMS)

INFORMS is located in Maryland, USA



### **Operations Research**

Publication details, including instructions for authors and subscription information: <a href="http://pubsonline.informs.org">http://pubsonline.informs.org</a>

# Random Walks, Fire Damage Amount and Other Paretian Risk Phenomena

Benoit Mandelbrot,

#### To cite this article:

Benoit Mandelbrot, (1964) Random Walks, Fire Damage Amount and Other Paretian Risk Phenomena. Operations Research 12(4):582-585. <a href="https://doi.org/10.1287/opre.12.4.582">https://doi.org/10.1287/opre.12.4.582</a>

Full terms and conditions of use: <a href="https://pubsonline.informs.org/Publications/Librarians-Portal/PubsOnLine-Terms-and-Conditions">https://pubsonline.informs.org/Publications/Librarians-Portal/PubsOnLine-Terms-and-Conditions</a>

This article may be used only for the purposes of research, teaching, and/or private study. Commercial use or systematic downloading (by robots or other automatic processes) is prohibited without explicit Publisher approval, unless otherwise noted. For more information, contact permissions@informs.org.

The Publisher does not warrant or guarantee the article's accuracy, completeness, merchantability, fitness for a particular purpose, or non-infringement. Descriptions of, or references to, products or publications, or inclusion of an advertisement in this article, neither constitutes nor implies a guarantee, endorsement, or support of claims made of that product, publication, or service.

© 1964 INFORMS

Please scroll down for article—it is on subsequent pages



With 12,500 members from nearly 90 countries, INFORMS is the largest international association of operations research (O.R.) and analytics professionals and students. INFORMS provides unique networking and learning opportunities for individual professionals, and organizations of all types and sizes, to better understand and use O.R. and analytics tools and methods to transform strategic visions and achieve better outcomes.

For more information on INFORMS, its publications, membership, or meetings visit <a href="http://www.informs.org">http://www.informs.org</a>

## RANDOM WALKS, FIRE DAMAGE AMOUNT AND OTHER PARETIAN RISK PHENOMENA

#### by Benoit Mandelbrot

Harvard University and I. B. M. Corporation (Received September 3, 1963)

Being one of the oldest branches of operations research, actuarial science has accumulated a substantial store of knowledge about the risks associated with living. The present paper will discuss one such question. Although it is relative to a specific problem of fire casualty, it illustrates more generally why the Paretian distribution of incomes and fortunes should constitute "a source of anxiety for the risk theory of insurance." Very similar mechanisms apply in many other problems.

IN A somewhat simplified form, the following statement summarizes an empirical law established by Lars-G. Benckert and Ingvar Sternberg:[1]

The damage to a house due to fire reaches different amounts with probabilities that obey a frequency function déduced as follows from Pareto's hyperbolic law: If the damage is greater than a minimum threshold m = 20 and smaller than the maximum destroyable amount M of the building, the probability that the damage be equal to x has the density  $x(x) = \alpha x^{-\alpha-1}m^{\alpha}$ . The probability that the maximum destroyable amount x0 be actually destroyed is equal to x1, the integral from x2 to infinity of the law of Pareto applicable to damages smaller than x3.

This law applies to all classes of Swedish houses outside of Stockholm. The values of  $\alpha$  were found to oscillate between 0.45 and 0.56; it seems fair therefore to begin by investigating the consequences of an assumed value  $\alpha = \frac{1}{2}$ .

#### A MODEL OF FIRE DAMAGE AMOUNT

THE LAW of Pareto of exponent 0.5 plays classically a central role as the distribution of the returns to equilibrium in the game of tossing a fair coin. This theory is developed in most textbooks of probability (such as reference 3), and it can be translated into insurance terms as follows.

Suppose that the intensity of a fire is characterized by a single number, designated by U, which can only take integral values; there is no fire when U=0; a fire starts when U becomes equal to 1, and it will end either when U becomes equal to 0 again, or when all that can possibly burn has already burned out.

Suppose now that, at any instant of time, there is a probability  $p=\frac{1}{2}$  that the fire encounter new material so that its intensity increases by 1,

and the probability  $q=\frac{1}{2}$  that the absence of new materials or the action of fire-fighters decrease the intensity by 1. In the preceding statement, 'time' is to be measured by the extent of damage. If there is no finite maximum extent of damage, and no lower bound to recorded damages, the duration of a fire will be an even number given by a classical result concerning the return to equilibrium in cointossing:

Pr(duration of fire = 
$$x$$
) =  $2 \binom{1/2}{x/2} (-1)^{x/2-1}$ .

Except for the first few values of x, this expression is proportional  $x^{-3/2}$ . Hence, if one assumes that very small damages, smaller than m, are not even properly recorded, the duration of a fire (i.e., the extent of damage) will be given by the law of Pareto with exponent  $\frac{1}{2}$ :

Pr(duration of fire>
$$x>m$$
) =  $(x/m)^{-1/2}$ .

Suppose finally that one takes account of the fact that the fire must end if and when all has burnt out. One then obtains precisely the interpretation of the Benckert-Sternberg findings that we gave above.

### RELATIONS INVOLVING THE SIZE OF THE PROPERTY AND THE EXPECTED AMOUNT OF THE DAMAGE DUE TO FIRE

It is easy to compute the expected value of the random variable considered in the opening section.

Expected amount of damage = 
$$\int_{m}^{M} x(\frac{1}{2}) x^{-(1/2+1)} m^{1/2} dx + M(M/m)^{-1/2}$$
  
=  $2\sqrt{Mm} - m$ .

This mean value tends to infinity with  $M.\dagger$ 

On the other hand, according to von Savitsch and G. Benktander, <sup>[2]</sup> the expected number of fires per house in the course of a year is a linear function of M. If this is indeed so, it would imply that, for large values of M, the rate of isnurance should be proportional to  $\sqrt{M}$ .

Note also that, when the distribution of property sizes M is known, one has:

Pr(amount of damage>x)=Pr(
$$M>x$$
) $(x/m)^{-1/2}$ .

If moreover the von Savitsch inference is correct, one has

d Pr(amount of damage per year>x) = 
$$Cxd$$
Pr(amount of damage>x)  
=  $Cxd$ [Pr( $M$ >x)( $x/m$ )<sup>-1/2</sup>].

<sup>†</sup> Random variables with infinite population moments are often considered to be mathematical devices with no practical application. Such is certainly not the case; see reference 4.

Let the distribution of M be itself Paretian with the exponent  $\alpha^*$ , as is the case of all kinds of liability amounts. The distribution of damage in a single fire will then be Paretian with exponent  $\alpha^* + \frac{1}{2}$ , and the distribution of damage per year will be Paretian with exponent  $\alpha^* - \frac{1}{2}$ . This demonstrates that mathematical manipulations based on Pareto's law are especially convenient.

#### GENERALIZATION

The random walk with  $p=q=\frac{1}{2}$  represents a kind of equilibrium state between the fire and the fire-fighters: if indeed the quantity of combustible property were unbounded, a fire following the rules given above will surely die out, although its expected duration would be infinite. On the contrary, if p>q, the fire-fighting efforts would be inadequate, and there would be a nonvanishing probability that the fire continue forever.

If p < q, the probability of running forever would again be zero and the expected duration of the fire would be finite. The law giving the duration of the fire would then take the form

d Pr(duration of fire 
$$\geq x > m$$
)  $\sim c^* \exp(-cx) x^{-3/2} dx$ ,

where  $c^*$  and c are two constants depending upon m, p, and q. If x is actually bounded, and if q-p (and hence c) is small, the above formula will be indistinguishable in practice from a law of Pareto with an exponent slightly greater than  $\frac{1}{2}$ ; this is perhaps an explanation of the more precise experimental results of Benckert and Sternberg.

We shall not stop to rephrase the generalization of the random walk, provided by any other of the many classical models of diffusion. Nor shall we go into the details of a random walk model in which there is the possibility that the intensity of the fire remain unchanged from one instant of time to the next. This would not add anything beyond a change in time scale.

#### ANOTHER APPLICATION

The results of Benckert and Sternberg strongly resemble those of Lewis F. Richardson,<sup>[7] [8]</sup> and the model that has been sketched above is easy to translate in the terms of Richardson's problem. It would be fascinating to ponder how relevant that translation may be.

#### REFERENCES

- LARS-G. BENCKERT AND INGVAR STERNBERG, "An Attempt to Find an Expression for the Distribution of Fire Damage Amount," Trans. 15th Internat. Congress of Actuaries, vol. II, pp. 288-294, New York, 1957.
- 2. Gunnar Benktander, "On the Variation of the Risk Premium with the Di-

- mension of the House within Fire Insurance," Skandinavisk Aktuarietidskrift, 26, 203-214 (1953).
- 3. WILLIAM FELLER, An Introduction to Probability Theory and Its Applications, Wiley, New York, 1957.
- K.-G. HAGSTROEM, "Remarks on Pareto Distributions," Skandinavisk Aktuarietidskrift, 33, 57-71 (1960).
- 5. Benoit Mandelbrot, "The Pareto-Levy Law and the Distribution of Income," International Economic Rev. 1, 79-106 (1960).
  - ——, "The Stable Paretian Income Distribution, When the Apparent Exponent Is Near Two," International Economic Rev. 4, 111-115 (1963).
  - -----, "Stable Paretian Random Processes and the Multiplicative Variation of Income," *Econometrica* 29, 517-543 (1961).
  - ——, "Paretian Distributions and Income Maximization," Quart. J. Economics 76, 57-85 (1962).
  - , "The Variation of Certain Speculative Prices," J. Business 36, 394-419 (1963).
  - , "New Methods in Statistical Economies," J. Political Economy 71, 421-440 (1963).
  - JAY M. BERGER AND —— "A New Model of Error Clustering in Telephone Circuits," I. B. M. J. Research and Development, 7, 224-236 (1963.)
- BIRGER MEIDELL, "Zur Theorie des Maximums" (with thorough English and French summaries) Septième Congrès d'Actuaires, Amsterdam, vol. I, pp. 85-99 (1912).
  - -----, "Zur Theorie und Praxis des Maximums in Lebensversicherung," Comptes-Rendus du XI-ième Congrès d'Actuaires, Vol. I, p. 469, Paris (1937).
- 7. Lewis F. Richardson, Statistics of Deadly Quarrels and Arms and Insecurity, Quadrangle Books, Chicago, 1960.
- HERBERT K. WEISS, "Stochastic Model for the Duration and Magnitude of a Deadly Quarrel", Opns. Res. 11, 101-121 (1963).

Copyright 1964, by INFORMS, all rights reserved. Copyright of Operations Research is the property of INFORMS: Institute for Operations Research and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.