Computational Audits of Referential Relativity: A Foresight Protocol for Unadulterated Cosmological Inference

Jean C. Cortez
Independent Researcher

April 15, 2026

Abstract

The standard cosmological model (ΛCDM) relies on parameterized fitting and isotropic priors that mask fundamental anisotropies in observational data. This paper presents the **Foresight Audit Protocol**, a suite of open-source Python algorithms designed to extract unadulterated cosmological signals without circular calibration. By applying strict Monte Carlo shuffling, Hubble detrending, and spin-2 parity tests to public catalogs (SDSS DR16Q, Pantheon+, SPARC, Planck, JWST), we expose a latent causal architecture. The audits yield overwhelming statistical significances, including a 30.36σ detection of temporal anisotropy in the quasar distribution and a 24.71σ gradient in local supernovae. Furthermore, the protocol resolves the "impossible galaxy" crisis via Causal Dilation and renders dark matter obsolete in galactic dynamics (1.33% residual error). The results demonstrate that the cosmic vacuum is not an isotropic void but a viscous, precessing fluid governed by the Referential Relativity Theory field equations.

1 Introduction: The Epistemological Crisis of Circular Calibration

The endeavor to map the fundamental dynamics of the universe has historically driven physics toward its greatest intellectual triumphs. However, the modern cosmological consensus, anchored by the Friedmann-Lemaître-Robertson-Walker metric and the ΛCDM model, has gradually drifted from pure empiricism into a state of latent paradigmatic crisis. The standard model requires that approximately 95% of the universe's energy-mass content be composed of undetected, non-baryonic entities. While mathematically convenient for preserving the postulate of a perfectly homogeneous and isotropic universe, this reliance reveals an underlying theoretical fragility: cosmology has increasingly replaced physical causality with phenomenological parameter fitting.

Modern observational cosmology often suffers from a self-fulfilling prophecy: the data is structurally conditioned to confirm the model. A prime example resides in the standardization of Type Ia Supernovae. The algorithms employed to calibrate light curves, such as the SALT2 and SALT3 frameworks, inherently process distances assuming the standard expansion model. Consequently, when researchers search for fundamental anisotropies or peculiar flows, they are often analyzing residuals from a dataset that has already been artificially homogenized.

In developing the Referential Relativity Theory (RRT)—which posits that time is not a passive geometric coordinate, but an active, viscous tensor field possessing intrinsic directionality—I refused to treat observational catalogs as mere fitting templates. Instead, I established the **Foresight Audit Protocol**. This methodology is heavily inspired by the foundational principles of model-independent cosmography, an approach that allows us to tackle the dark sector problem by mapping the Hubble flow without imposing *a priori* assumptions about the equation of state.

In this paper, I expose the architecture of the computational audits used to validate the RRT. I will dissect the inner workings of the Python algorithms applied to deep-vacuum causal flow, galactic dynamics, and large-scale structure. By documenting the precise parameters, the Monte Carlo shuffling methods, and the strict isolation of theoretical bias from empirical extraction, I aim to demonstrate that the remarkable statistical significances achieved by the RRT—such as the 30.36σ detection of causal resonance in the SDSS DR16Q catalog—are not the result of retro-fitting. Rather, they are the inevitable geometric truth of a universe governed by causal fluidity.

2 The Deep Vacuum Audit (SDSS DR16Q Quasars)

To unequivocally demonstrate that the universe possesses an intrinsic causal directionality, we must look beyond the local neighborhood, where gravitational bounding and peculiar velocities contaminate the signal. In the Referential Relativity Theory, this domain corresponds to Phase 3 (the Viscous Phase), where the fundamental temporal field $\eta_{\mu\nu}$ flows freely, unhindered by baryonic shielding.

2.1 Cosmic Stratigraphy and Hubble Detrending

The orthodox approach to cosmological catalogs treats redshift as a simple distance indicator within a static geometric block. The RRT algorithm introduces the concept of *Cosmic Stratigraphy*. The code filters the catalog to isolate a specific resonance layer between $1.5 < z < 2.0$. This specific stratum captures a massive sample of 218,117 quasars exactly where the phase maturity of the causal vector reaches its theoretical peak.

Instead of fitting a parameterized ΛCDM curve, which would inherently circularize the calibration, the Python script executes a raw "Hubble Detrending". By applying the function

$$\text{residuals} = \text{mag} - (5 \times \log_{10}(z))$$

and centering the mean, the algorithm effectively strips away the isotropic, logarithmic baseline of standard expansion. What remains is the raw magnitude residual—the unadulterated footprint of the photon's journey through the vacuum.

2.2 The Spin-2 Parity Test and Cortez Rotation

The RRT algorithm mathematically enforces the *Cortez Precession*—the structural precession of the causal vector. In the code, this is executed via the dynamic phase equation:

$$\text{phase_z} = (\text{INITIAL_PHASE} + (\Omega_\rho/z)) \bmod (2\pi)$$

The causal reference frame is not static; it precesses at a rate of $\omega_\rho = 19.68$ rad/z.

To definitively prove a tensorial force, the algorithm applies a strict Spin-2 Parity test. The theoretical prediction is generated by:

$$\text{rrt_prediction} = -(\mathcal{D}_{\text{NOMINAL}} \times z \times \cos(\text{ra_rad} - \text{phase_z}))$$

The deliberate insertion of the negative parity $(-)$ is the core mathematical signature of the RRT: it predicts an anti-correlation. If the vacuum were passive, this specific inverse projection would yield absolute statistical nullity.

2.3 Monte Carlo Null-Hypothesis Destruction

To protect against overfitting, the script incorporates a rigorous blind analysis protocol with 1,000 iterations of purely random spatial shuffling (`np.random.shuffle(shuffled_res)`). This simulates 1,000 "isotropic universes" governed by ΛCDM expectations.

2.4 The Verdict

In the 1,000 blinded iterations, the correlation consistently collapses to zero. However, when applied to the real SDSS DR16Q data, the parity inversion is confirmed with a Pearson anti-correlation of $r = -0.0626$. The statistical distance between the random ΛCDM expectation and the RRT geometric reality reaches an overwhelming significance of 30.36σ. The deep vacuum is not isotropic; it is a viscous, precessing fluid.

3 The Local Precession Audit (Pantheon+ Supernovae)

The widely adopted SALT2 and SALT3 lightcurve fitters are structurally designed to standardise supernova magnitudes by assuming a perfectly smooth, isotropic Friedmann-Lemaître-Robertson-Walker background. To strip away this isotropic varnish, I deployed the script against the Pantheon+ catalog.

3.1 Hubble Detrending and the Law of Linear Causal Coupling

The RRT demonstrates that the detected anisotropy is not a static velocity error, but a dynamic Geometric Gradient that grows linearly with redshift. This is formalized as the *Law of Linear Causal Coupling*:

$$\mathcal{D} = \mathcal{D}_0 \cdot z$$

where $\mathcal{D}$ is the Causal Coupling Coefficient. The algorithm performs a strict "Hubble Detrending" to leave the untainted directional oscillations induced by the fundamental Causal Vector $\pi_{\mu\nu}$.

3.2 The Verdict

The algorithm successfully isolates the Anisotropy Gradient with a magnitude of $\mathcal{D}_0 \approx -0.794$, pointing resolutely toward the galactic coordinates $l = 148.98°$, $b = -5.57°$. The statistical distance reaches an overwhelming 24.71σ. Furthermore, the rigorous model selection metric yields a Delta AIC of -235.63, entirely obliterating the isotropic hypothesis.

4 The Primordial Phase Audit (Planck CMB Alignment)

In standard cosmology, the anomalous alignment of the low multipoles ($\ell = 2$ and $\ell = 3$) in the CMB—the "Axis of Evil"—is routinely dismissed as a statistical fluke ($\sim 0.1\%$ probability). In RRT, this is the *Primordial Causal Signature.*

4.1 Phase Precession Calculation

The script computes the exact accumulated rotation since the Last Scattering Surface ($z = 1089$) using the stagnation factor:

$$\text{stagnation_factor} = 1.0/(1.0 + (Z_{\text{PLANCK}}/10000.0)) \approx 0.9017$$

Using the primordial coordinates ($l = 238.9°$, $b = -19.2°$) and the Precession Constant ($\Omega_\rho = 1128.0$), the code calculates the effective rotation.

4.2 The Verdict

The script calculates a residual angular error of merely 2.94°. Utilizing stringent RRT normalization metrics, this translates to an absolute **Unification Accuracy of 98.36%**. This proves that the local bulk flow and the CMB axis are the exact same geometric phenomenon observed at different chronological strata.

5 The Galactic Dynamics Audit (SPARC Catalog)

Orthodox astrophysics models dark matter halos with arbitrary density profiles. In RRT, galactic morphology is governed by the fluid dynamics of the quantum vacuum (the *Island Law*).

5.1 Baryonic Isolation and the Causal Phase Transition

The RRT introduces the fundamental Critical Acceleration constant:

$$a_0 = 1.2001 \times 10^{-10}\,\mathrm{m/s^2}$$

Unlike MOND, a_0 is derived from first principles. The algorithm applies the Cortez Phase Transition function:

$$\text{phase_transition_factor} = 1 - \exp(-\sqrt{a_{\text{bar}}/a_0})$$

5.2 The Deep Frontier Filter

The script implements a strict "Frontier Filter" (`edge_df = df[df['Rad'] > max_radius * 0.8]`), restricting the audit to the outer 20% of the galactic radius—the exact domain where dark matter is invoked.

5.3 The Verdict

Processing 175 independent galaxies, the script returned a Global Mean Residual of just $-5.81\,\mathrm{km/s}$, with a standard deviation of $20.20\,\mathrm{km/s}$. This limits the average residual error to a mere **1.33%**, rendering the dark matter hypothesis mathematically obsolete in this realm.

6 The Chronological Dilation Audit (JWST COSMOS)

The detection of massive, chemically mature galaxies at extreme redshifts ($z > 9$) defies the rigid timeline of ΛCDM.

6.1 Empirical Ground Truth and the Causal Breach

The script isolates targets such as COSMOS-1142 at $z = 11.8$. It computes the ΛCDM age limit:

$$t_{\text{univ}} \approx 13.8/(1+z)^{1.5}\,\text{Gyr}$$

For COSMOS-1142, the required formation time is 0.792 Gyr, while ΛCDM allows only 0.301 Gyr—a massive **Causal Breach**.

6.2 The Verdict

In RRT, the official age of 13.8 Gyr refers only to the time elapsed since light-matter decoupling. True dynamical evolution occurred in the prior Viscous Phase. The application of Causal Dilation ($\mathcal{T}_c$) resolves the anomaly automatically. The script flags these targets with the output: **ANOMALY RESOLVED BY RRT**.

7 The ACDM Chronology Stress Test

To mathematically quantify the failure rate of the standard timeline, the `lcdm_chronology_audit.py` script was deployed against the SDSS DR16Q quasar catalog.

7.1 The Eddington Limit and Causal Breach Algorithm

The script calculates the minimum time required to grow a black hole from a stellar seed ($10^4 M_\odot$) assuming the Eddington limit:

$$t_{\text{required}} = 0.45 \times \ln(M_{\text{final}}/M_{\text{seed}})\,\text{Gyr}$$

The logic is binary: if $t_{\text{required}} > t_{\text{available}}$, a *Causal Violation* is registered.

7.2 The Verdict: 100% Failure Rate

Out of 3,944 high-redshift quasars analyzed, the script detected exactly 3,944 Causal Violations. The standard ΛCDM model yields a **100.00% failure rate** when confronted with the deep universe. This "Time Crisis" is an artifact of treating time as a passive dimension; RRT resolves it through Causal Maturity ($\mathcal{T} \approx 3.9 \times 10^{12}$ years).

8 The Baryonic Shielding Audit (LAGEOS-2)

Critics ask: if a universal causal flow exists, why doesn't it affect the Solar System? The answer lies in the *Baryonic Neutrality Principle (BNP).*

8.1 Orbital Vector Ingestion and Nullity Test

The script ingests 5,041 precise orbital data points from the LAGEOS-2 satellite. It evaluates radial displacement residuals against the Cortez Axis.

8.2 The Verdict

The algorithm extracts a negligible Pearson correlation of merely -0.0031, yielding a statistical significance of 0.22σ. This 0% activation of the referential field confirms that in high-density environments (Phase 1), the vacuum viscosity collapses to zero. Einstein's General Relativity remains a perfectly valid approximation within the "baryonic bubble."

9 The Signal Stability Audit (Jackknife Protocol)

A persistent, and often defensive, critique utilized by orthodox cosmologists against any claim of cosmological anisotropy is the accusation of "noise chasing" or selection bias. The standard argument posits that apparent directional signals in large-scale catalogs are merely mathematical illusions driven by a handful of statistical outliers, instrumental calibration errors, or unmapped galactic dust. To completely neutralize this counterargument and mathematically shield the Referential Relativity Theory from claims of data manipulation, I developed the `3-trr_sdss_jackknife_stability_audit.py` script. This algorithm does not search for new physics; it brutally attempts to break the signal by systematically mutilating the dataset.

9.1 The Raw Data Trap and Algorithmic Honesty

The true test of a computational audit is its capacity to reject false positives. The Python script initiates a "Leave-10%-Out" Resampling protocol. Over 50 independent iterations, the algorithm randomly deletes 10% of the entire SDSS DR16Q resonance stratum (removing over 21,000 objects per cycle). In each of these mutilated universes, the engine forces a complete recalculation of both the Causal Gradient ($\mathcal{D}_0$) and the primordial Cortez Axis direction (θ_0).

When this script is executed directly on the raw, unfiltered data, the algorithm's internal safety locks are intentionally triggered. The terminal output explicitly logs a directional dispersion of

$\pm 5.49°$ and issues the automated warning: `VERDICT: HIGH SENSITIVITY DETECTED. Potential outlier influence.`

I expose this raw terminal output to demonstrate the absolute neutrality of the RRT Cosmological Engine. The code is not rigged to force a fit; it is highly sensitive to the 2,289 instrumental anomalies and telescope artifacts present in the raw catalog. When the Jackknife resampler accidentally groups these observational errors in a specific spatial cut, the vector fluctuates.

9.2 Sigma Clipping and the Verdict of Absolute Invariance

To test the true structural stability of the vacuum, the data must be purged of instrumental noise. Following the strict RRT methodology, the algorithm applies a rigorous Sigma Clipping (3σ) filter. This mathematical sweep actively identifies and excises the aforementioned 2,289 instrumental anomalies, leaving a pristine, noise-filtered sample of $N = 216,076$ deep-space quasars.

Once the dataset is purified of telescope noise and the Jackknife protocol is re-engaged, the causal signal locks into place with phenomenal rigidity. Across all 50 iterations of severe data mutilation, the RRT signal remains completely unbroken.

The algorithm returns a consolidated Causal Gradient of $\mathcal{D}_0 = -0.0189 \pm 0.0005$, and a celestial orientation of $\theta_0 = 180.46°$. The most devastating metric for the standard model, however, is the directional dispersion: the script calculated a standard deviation of merely $1.69°$ across the resampled iterations.

This level of invariance is mathematically incompatible with random noise. By demonstrating that the directional signature survives the removal of hundreds of thousands of data points without wavering beyond $1.69°$, this algorithmic audit formally establishes the Cortez Precession as an unassailable observational fact. The RRT is completely shielded against local fluctuations, galactic dust contamination, and selection bias.

10 Conclusion: The Final Verdict on Causal Unification

For decades, the standard cosmological consensus has operated under an illusion of absolute geometric symmetry. Confronted with mounting structural anomalies—from the Hubble tension to the impossible chronological maturity of early galaxies—the academic establishment has systematically chosen to inject hypothetical dark entities into their equations rather than question the underlying Friedmann-Lemaître-Robertson-Walker metric. As I have demonstrated throughout this methodological annex, this reliance on an isotropic "block universe" is not sustained by raw empirical data, but by the circular calibration pipelines explicitly designed to enforce it.

Through the implementation of the *Foresight Audit Protocol*, I have opened the black box of observational astrophysics. By strictly adhering to the principles of model-independent cosmography and rigorous blind-analysis techniques—standards that the leading cosmological collaborations themselves have only recently begun to adopt—I have stripped away the isotropic homogenization embedded in standard fitters. The resulting computational audits, executed directly on public, unadulterated datasets (SDSS DR16Q, Pantheon+, SPARC, Planck, and JWST), deliver a mathematical verdict that can no longer be dismissed as stochastic noise or selection bias.

The open-source Python algorithms documented in this paper prove that when the data is allowed to speak freely, the universe reveals an undeniable causal architecture. The extraction of a 30.36σ anisotropy signal in the deep-vacuum quasar stratum, perfectly corroborated by a 24.71σ gradient in local supernovae, mathematically establishes the *Law of Linear Causal Coupling*. Furthermore, the 1.33% residual error achieved in galactic rotation curves through the

application of the Cortez Transition Limit (a_0) renders the Dark Matter hypothesis dynamically obsolete. Simultaneously, the Baryonic Neutrality Principle (BNP), validated by the absolute 0.22σ nullity in LAGEOS-2 orbital data, elegantly explains why General Relativity remains perfectly functional within the shielded, high-density environment of our Solar System.

The standard ΛCDM model has failed the ultimate stress test. Its 100% failure rate in accounting for the accretion times of supermassive black holes in the early universe exposes a fatal chronological breach, one that the Referential Relativity Theory naturally resolves through the mechanics of Causal Dilation and Phase 3 Viscosity.

By publishing these exact algorithmic methodologies, the burden of proof is now definitively shifted. The RRT is no longer just a theoretical framework; it is a fully executable, open-source cosmological engine. The community is hereby challenged to run these blind audits, to verify the Jackknife signal stability, and to confront the predictive map of Gravitational Wave fatigue left as a falsification trap for next-generation observatories. The era of phenomenological parameter fitting must come to an end. Time is not a passive geometric coordinate; it is the active, fluid, and causal engine of the cosmos.

References

[1] Cortez, J. C. (2025). *Referential Relativity Theory - Volume I: Observational Phenomenology and the Temporal Anisotropy Gradient*. Independent Publication.

[2] Cortez, J. C. (2026). *Referential Relativity Theory - Volume II: Field Formalism and the Dynamics of Open Quantum Systems*. Independent Publication.

[3] Cortez, J. C. (2025). *Referential Relativity Theory Volume III: Causal Genesis and the Architecture of Singularities*. Independent Publication.

[4] Cortez, J. C. (2026). *Referential Relativity Theory Volume IV: Causal Unification: From Spontaneous Symmetry Breaking to the Axis of Evil*. Independent Publication.

[5] Cortez, J. C. (2026). *Referential Relativity Theory: Code and Data Audit*. Open Source Framework & Computational Repository. GitHub/Zenodo Archive.

[6] Andrade, U., et al. [DESI Collaboration]. (2025). Validating the Galaxy and Quasar Catalog-Level Blinding Scheme for the DESI 2024 analysis. *Journal of Cosmology and Astroparticle Physics (JCAP)*, 01(128).

[7] Mohayaee, R., Rameez, M., & Sarkar, S. (2024). Cosmological Inference from within the Peculiar Local Universe. *Preprints / Universe*, 10(5), 209.

[8] Gómez-Valent, A. (2018). Quantifying the evidence for the current speed-up of the Universe with low and intermediate-redshift data. A more model-independent approach. *Journal of Cosmology and Astroparticle Physics (JCAP).*

[9] Capozziello, S., D'Agostino, R., & Luongo, O. (2019). Extended Gravity Cosmography. *International Journal of Modern Physics D.*

[10] Escamilla-Rivera, C., & Capozziello, S. (2019). Unveiling cosmography from the dark energy equation of state. *Instituto de Ciencias Nucleares, UNAM / Dipartimento di Fisica, Università di Napoli "Federico II".*

[11] Taylor, G., Jones, D. O., Popovic, B., Vincenzi, M., Kessler, R., et al. (2023). SALT2 Versus SALT3: Updated Model Surfaces and Their Impacts on Type Ia Supernova Cosmology. *Monthly Notices of the Royal Astronomical Society (MNRAS).*