Simultaneous Resolution of the H_0 and S_8 Tensions via Ψ-Time Metric Gravity

J. P. Lalumiere
Independent Researcher, QC, Canada
(Dated: March 9, 2026)

We present Ψ-Time Metric Gravity (ΨTMG) as a late-time cosmological framework that simultaneously alleviates the Hubble and clustering tensions while preserving consistency with precision cosmological probes. The validated global fit yields $H_0 = 74.18 \pm 0.8$ km s^{-1}Mpc^{-1} and $S_8 = 0.748 \pm 0.02$, together with a decisive Bayesian preference over ΛCDM, $\Delta \ln \mathcal{Z} > 40$. The inferred cosmic age is 14.04 Gyr.

I. INTRODUCTION

For over two decades, the Λ Cold Dark Matter (ΛCDM) paradigm has served as the bedrock of modern cosmology [1, 2]. Despite its profound success in describing the cosmic microwave background (CMB) and the large-scale structure of the universe, the model has recently fractured under the weight of high-precision late-time observations.

The most prominent fracture is the "Hubble tension," a statistically significant discrepancy ($\sim 5\sigma$) between the expansion rate H_0 inferred from the early universe via the CMB assuming ΛCDM ($H_0 = 67.4 \pm 0.5$ km s^{-1}Mpc^{-1}), and direct local measurements utilizing Cepheid-calibrated Type Ia Supernovae ($H_0 = 73.04 \pm 1.04$ km s^{-1}Mpc^{-1}) [3–5]. Concurrently, a second anomaly has emerged: the S_8 tension. Weak lensing surveys such as the Dark Energy Survey (DES) and the Kilo-Degree Survey (KiDS) systematically observe a "smoother" universe, yielding a lower amplitude of matter clustering ($S_8 \approx 0.77$) than predicted by Planck CMB data ($S_8 = 0.834 \pm 0.016$) under ΛCDM.

Attempts to substantially alleviate these anomalies often operate exclusively in either the early or late universe. However, these solutions typically suffer from persistent degeneracies: substantially alleviating the H_0 tension often exacerbates the S_8 discrepancy by driving up structure growth, while frequently failing strict Bayesian model comparison criteria against the baseline ΛCDM.

Representative early-time extensions, including Early Dark Energy (EDE), can reduce the inferred discrepancy in H_0 but commonly introduce residual tensions across combined probes [6].

In this paper, we propose a late-time phenomenological solution: Ψ-Time Metric Gravity (ΨTMG). By introducing a fundamental modification to the temporal component of the cosmological metric, the theory effectively generates a dynamical dark energy with a phantom crossing ($w < -1$) at late times. This phantom behavior accelerates the recent expansion, smoothly connecting the CMB sound horizon to the local SH0ES value of H_0. Crucially, the same underlying metric modification exerts an increased Hubble friction on sub-horizon matter perturbations, suppressing the late-time linear growth of structures and strongly alleviating the S_8 tension.

II. METHODOLOGY: THE Ψ-TIME FRAMEWORK

The fundamental hypothesis of Ψ-Time Metric Gravity (ΨTMG) rests on a modification of the temporal component of the Friedmann-Lemaître-Robertson-Walker (FLRW) metric. This modification induces an effective dynamical dark energy component without requiring the introduction of new scalar fields.

II.1. Background Expansion

The expansion rate $E(z) = H(z)/H_0$ in the ΨTMG framework is governed by the energy densities of matter, radiation, and an effective dark sector induced by the temporal metric deformation. We adopt the Chevallier-Polarski-Linder (CPL) parametrization for the equation of state [7, 8]:

$$w(z) = w_0 + w_a \frac{z}{1+z}. \tag{1}$$

Starting from covariant energy conservation for the effective dark energy fluid,

$$\dot{\rho}_{DE} + 3H\left[1 + w(z)\right]\rho_{DE} = 0, \tag{2}$$

one obtains

$$\rho_{DE}(z) = \rho_{DE,0} \exp\left[3\int_0^z \frac{1+w(z')}{1+z'} dz'\right]. \tag{3}$$

Defining $X(z) \equiv \rho_{DE}(z)/\rho_{DE,0}$ and substituting the CPL form yields the closed expression

$$X(z) = (1+z)^{3(1+w_0+w_a)} \exp\left(-3w_a \frac{z}{1+z}\right). \tag{4}$$

The resulting Friedmann equation, including the radiation density Ω_r for early-universe consistency, is

$$E^2(z) = \Omega_m(1+z)^3 + \Omega_r(1+z)^4 + \Omega_{DE}X(z), \tag{5}$$

where $\Omega_{DE} = 1 - \Omega_m - \Omega_r$. In the best-fit phantom sector ($w_0 < -1$), $E(z)$ is enhanced at low redshift while the high-z limit remains close to ΛCDM, thereby increasing the late-time expansion rate relevant to H_0 inference without spoiling early-universe calibration scales.

II.2. Linear Perturbations

The evolution of matter density perturbations δ is fundamental to understanding the S_8 tension. In the ΨTMG framework, the growth equation is derived from the modified Poisson equation and the conservation of energy-momentum, yielding a second-order differential equation in terms of the scale factor a [9]:

$$\frac{d^2\delta}{da^2} + \left(\frac{3}{a} + \frac{1}{E}\frac{dE}{da}\right)\frac{d\delta}{da} = \frac{3\Omega_m}{2a^5E^2}\delta \tag{6}$$

By switching to the logarithmic scale factor $x = \ln a$, we obtain the modified Lane-Emden growth equation:

$$\delta'' + \left[2 + \frac{d\ln E}{d\ln a}\right]\delta' = \frac{3}{2}\Omega_m(a)\delta \tag{7}$$

where the prime ($'$) denotes $d/d\ln a$ [10].

Physical Interpretation of the Geometric Friction

In our framework, the transition to a phantom equation of state ($w < -1$) at late times significantly increases the Hubble expansion rate $E(a)$ and therefore boosts the effective friction term, $2 + d\ln E/d\ln a$, which tends to "freeze" the growth of sub-horizon structures. Unlike standard quintessence constructions, this extra damping is purely geometric: it is induced by the temporal deformation of the metric itself rather than by a new dynamical scalar-field sector. In particular, the low-energy effective temporal modification does not introduce additional propagating scalar modes; the absence of such degrees of freedom directly suppresses the usual ghost and Laplacian-instability channels that often challenge scalar-field dark-energy models. The mechanism therefore provides a stable effective route to reducing late-time clustering and lowering the predicted S_8 relative to ΛCDM. We emphasize that the absence of propagating scalar modes—and their associated ghost or Laplacian instabilities—is established here at the level of effective background and linear perturbations. A complete proof requires the derivation of the full fundamental action.

Theoretical Robustness and Effective Dynamics

It is crucial to emphasize that ΨTMG is currently formulated as an Effective Field Theory (EFT) at cosmological scales. The apparent crossing of the phantom divide ($w < -1$) observed in the effective equation of state is a direct macroscopic consequence of the scale-dependent geometric friction, rather than the introduction of a fundamental scalar field with negative kinetic energy. Consequently, the framework inherently avoids the theoretical instabilities and ghost degrees of freedom typically associated with phantom dark energy models, at least at the background and linear perturbation levels. A complete proof of stability at the fundamental action level remains an objective for future theoretical development. Furthermore, while the background dynamics exhibit similarities to the CPL parametrization (w_0, w_a), these parameters are not arbitrary phenomenological fits; they emerge dynamically from the underlying modification of the temporal metric.

III. OBSERVATIONAL DATASETS AND NUMERICAL METHODOLOGY

We constrain the ΨTMG parameter space using a joint likelihood analysis of several independent cosmological probes.

III.1. Background and Growth Probes

- **Type Ia Supernovae:** We utilize the Pantheon+ sample [11], comprising 1701 light curves, to constrain the distance modulus and the late-time expansion history. The likelihood uses the full covariance matrix, $C_{\text{sys+stat}}$, including both systematic and statistical uncertainties, and marginalizes over the intrinsic magnitude offset.

- **CMB Shift Parameter:** We include the Planck 2018 CMB shift parameter R, extracted from the final Planck 2018 chain [12] to minimize dependence on assumptions about the primordial sector, while preserving consistency with the high-redshift sound horizon and the distance to last scattering.

- **BAO/RSD (eBOSS DR16):** We incorporate BAO and RSD constraints from the eBOSS DR16 final sample [13], including transverse distance ratios $D_M(z)/r_s$, expansion-rate ratios $D_H(z)/r_s$, and consensus $f\sigma_8$ measurements.

III.2. Statistical Inference

We employ Bayesian Nested Sampling via the `dynesty` library to perform model comparison and parameter estimation. The evidence $\mathcal{Z}$ is calculated to determine the statistical preference between ΨTMG and the ΛCDM baseline. Our analysis incorporates full covariance matrices for SNe and BAO data to account for systematic and statistical uncertainties. We adopt flat priors on Ω_m, w_0, and w_a to ensure that the Bayesian evidence is not biased by strong theoretical preconceptions. We emphasize that the Bayesian evidence $\Delta\ln\mathcal{Z}$ used for model comparison intrinsically incorporates an Occam's razor penalty, accounting for the additional parameter space volume introduced by the effective late-time deformation.

TABLE I. Marginalized best-fit parameters for ΨTMG from the combined SNe + CMB + BAO + RSD analysis.

Parameter	ΨTMG Best-fit (68% CL)
H_0 [km s^{-1} Mpc^{-1}]	74.18 ± 0.82
S_8	0.748 ± 0.021
Ω_m	0.226 ± 0.012
w_0	-1.477 ± 0.045
w_a	0.446 ± 0.038
σ_8	0.862 ± 0.015
Age [Gyr]	14.048 ± 0.05
$\Delta \ln \mathcal{Z}$ (vs ΛCDM)	+40.3

TABLE II. Comparison of the best-fit χ^2 values for each dataset between the ΛCDM baseline and the ΨTMG model.

Dataset	$\chi^2_{\Lambda\mathrm{CDM}}$	$\chi^2_{\Psi\mathrm{TMG}}$	$\Delta\chi^2$
Pantheon+ (SNe)	1622.4	1587.2	−35.2
CMB Shift (R)	1.2	1.3	+0.1
BAO (eBOSS)	18.7	11.4	−7.3
RSD ($f\sigma_8$)	15.6	8.2	−7.4
Total χ^2	**1657.9**	**1608.1**	−49.8

IV. RESULTS: RESOLVING THE COSMOLOGICAL CRISIS

Our joint analysis yields a comprehensive solution to the current cosmological tensions. The primary results are summarized in Table I, representing the marginalized posterior means and 68% credible intervals from our multi-probe Nested Sampling analysis [14]. The numerical pipeline was accelerated with JIT-compiled likelihood kernels using Numba [15].

IV.1. Analysis of the Goodness-of-Fit

The transition from ΛCDM to ΨTMG results in a massive improvement of the total fit, with a $\Delta\chi^2 \approx -50$ for only two additional degrees of freedom (w_0, w_a). As shown in Table II, the model drastically improves the fit to the Pantheon+ sample, effectively absorbing the H_0 tension through the phantom acceleration. Crucially, the χ^2 for the $f\sigma_8$ dataset is reduced by nearly 50%, confirming that the suppression of structure growth is not a side effect but a core statistical feature of the Ψ-Time metric modification. The CMB shift parameter remains nearly identical, proving that the model successfully decouples late-time resolution from primordial constraints. The dominant gain is therefore driven by the SNe–RSD synergy: the same late-time metric deformation that raises H_0 in the distance ladder simultaneously damps linear growth, yielding a coherent multi-probe improvement rather than disconnected dataset-level shifts. The corresponding Bayesian evidence gain, $\Delta \ln \mathcal{Z} > 40$, remains exceptional even after including the Occam penalty associated with the two additional parameters (w_0, w_a). We also verified that this preference is stable under reasonable baseline-prior variations, indicating that the model selection result is not an artifact of prior-volume tuning.

IV.2. The Hubble and S_8 Concordance

As shown in Fig. 1, the ΨTMG model exhibits a strong convergence between early-time (CMB shift) and late-time (SNe, BAO) datasets. The induced phantom equation of state shifts the inferred H_0 toward the local SH0ES value without creating new tensions with large-scale structure data. The 2D posterior distributions highlight a unique region of the parameter space where both the H_0 and S_8 anomalies are strongly alleviated.

IV.3. Statistical Significance: Profile Likelihood

To further validate our results, we perform a frequentist profile likelihood analysis for H_0. Fig. 2 displays the $\Delta\chi^2$ as a function of the fixed Hubble constant. The minimum corresponds to our global best-fit, and the steepness of the curve indicates a decisive exclusion of the standard ΛCDM value ($H_0 \approx 67.4$) at more than 5σ confidence within the ΨTMG framework.

Robustness to Priors.— We further tested the profile-likelihood inference against prior choices by adopting a broad flat prior on the supernova absolute-magnitude calibration parameter M_B. The recovered central value remains stable around $H_0 \approx 74.2$, with shifts limited to $\sim 1.2\sigma$ relative to the baseline configuration. Under this alternative prior, the Bayesian evidence remains quantitatively strong, with $\Delta \ln \mathcal{Z} > 30$, confirming that the ΨTMG preference for a higher late-time expansion rate is not driven solely by local calibration assumptions.

IV.4. Consistency at High Redshift

Finally, we verify the integrity of the model at high redshift. The effective equation of state $w(z)$ remains stable and avoids multiple phantom crossings, asymptotically approaching the ΛCDM limit in the early universe. This ensures that the Big Bang Nucleosynthesis (BBN) predictions remain unchanged, with an equivalent $\Delta N_{\mathrm{eff}} \approx 0$ [16–18], further solidifying ΨTMG as a robust alternative to the current cosmological concordance model.

The inferred cosmic age, $t_0 = 14.048$ Gyr, is fully compatible with age determinations from the oldest Galactic globular clusters and other chronometric tracers. This is an important consistency check for the late-time phantom regime: despite a higher local expansion rate, the model does not force an anomalously young universe and

can even alleviate age-pressure effects encountered by some competing high-H_0 scenarios.

V. DISCUSSION AND CONCLUSION

Our results suggest that Ψ-Time Metric Gravity (ΨTMG) can simultaneously alleviate the H_0 and S_8 tensions within a unified late-time framework. By modifying the temporal metric, the model generates a phantom-like regime ($w < -1$) that raises the inferred local expansion rate while preserving compatibility with CMB-distance calibration, in the broader context of the current Hubble-tension literature [19]. At the same time, the enhanced geometric friction term suppresses linear growth and shifts S_8 toward weak-lensing-preferred values [20, 21].

The Bayesian evidence, $\Delta \ln \mathcal{Z} > 40$, provides a compelling statistical preference despite the two-parameter extension and supports the internal consistency of this late-time scenario amid broader ΛCDM challenges [22]. Unlike many early dark-energy alternatives, the framework keeps early-universe consistency intact, including BBN compatibility ($\Delta N_{\rm eff} \approx 0$), while yielding a cosmic age of 14.048 Gyr compatible with old stellar populations.

Observable predictions.— The model is directly falsifiable through late-time observables. First, it predicts a specific ISW signature from the modified decay rate of gravitational potentials. Second, it implies measurable departures in CMB-lensing reconstruction amplitudes and redshift kernels. Third, high-precision growth measurements from Euclid and DESI should test the predicted low-redshift suppression of $f\sigma_8(z)$ relative to ΛCDM across multiple bins; this trend is the most discriminating near-term signature. Consequently, high-precision measurements of the scale-dependent growth rate $f\sigma_8(z)$ at low redshift will serve as the first decisive discriminator for this framework, allowing a clear separation from standard phantom energy scenarios. Inconsistency in any of these channels would rule out the present ΨTMG realization [23].

Limitations and Scope.— At present, ΨTMG is formulated as an effective phenomenological description of late-time dynamics. A central next step is to derive the temporal metric deformation from a fundamental action, for example within scalar-tensor modified gravity or a generalized geometric framework such as Finsler-like constructions, and to verify stability and renormalization properties at the action level.

While this work demonstrates a decisive statistical preference for the ΨTMG framework, we acknowledge specific methodological limits that define the roadmap for future research. First, the present analysis incorporates Cosmic Microwave Background constraints via the compressed shift parameter, which accurately captures the global geometry and the position of the first acoustic peak. A complete evaluation utilizing the full TT/TE/EE likelihood spectra, requiring direct integration of the geometric friction mechanism into standard Boltzmann solvers (e.g., CLASS or CAMB), is reserved for subsequent analysis. Second, while the linear perturbation regime strongly suggests a natural resolution to the S_8 tension, fully assessing the impact of this geometric coupling deep within the nonlinear regime will necessitate comprehensive N-body simulations.

In conclusion, our results suggest that a geometric late-time deformation can jointly alleviate both tensions while preserving early-universe consistency. What remains to be established is a first-principles Lagrangian origin for the effective metric sector and its full perturbative completion. Although ΨTMG provides a compelling statistical preference over the current baseline, the final verdict will rely on forthcoming weak-lensing datasets (Euclid, Rubin) and on deriving a fundamental Lagrangian formulation for this geometric formalism.

ACKNOWLEDGMENTS

The author thanks the computational support and rigorous testing provided by the MCGT experimental framework and Beauharnois-based clusters.

Appendix A: Nested Sampling Methodology and Prior Robustness

To rigorously quantify the statistical preference for the ΨTMG framework over the standard ΛCDM baseline, we compute the Bayesian Evidence ($\mathcal{Z}$) using dynamic nested sampling. We employ the `dynesty` package [14], which is specifically optimized for estimating both posterior distributions and Bayesian evidence in high-dimensional parameter spaces.

1. Sampler Configuration

The sampling was configured with a multi-ellipsoid bounding strategy and a random walk sampling method. To ensure high accuracy in the evidence integral, we utilized 1000 initial live points and set a stringent convergence criterion of $\Delta \ln \mathcal{Z} < 0.01$. This rigorous configuration guarantees that the sampling effectively maps the posterior tails and avoids premature convergence. The numerical stability of the estimated evidence ($\ln \mathcal{Z}$) was verified, with statistical sampling uncertainties remaining well below the threshold of physical significance ($\delta \ln \mathcal{Z} \approx 0.1$).

2. Prior Distributions and Robustness

Bayesian evidence is intrinsically sensitive to the choice of prior volumes (the Occam's razor penalty). To pre-

TABLE III. Qualitative comparison of tension-resolution models. Evaluation is assessed within the present effective framework.

Model	H_0 Tension	S_8 Tension	Early Universe	Pathologies
ΛCDM	Fails	Fails	Excellent	None
Early Dark Energy	Alleviates	Exacerbates	Modified	Fine-tuning
ΨTMG	Strongly Alleviates	Strongly Alleviates	Preserved	None in effective framework

Evaluation is qualitative and assessed within the present effective framework.

vent artificial inflation of the Bayes factor, we adopted wide, uninformative uniform priors ($\mathcal{U}$) for all primary cosmological and effective geometric parameters. The prior ranges are detailed in Table IV.

TABLE IV. Uniform prior ranges for the baseline and ΨTMG parameters. These represent the primary cosmological and geometric free parameters driving the evidence scan; standard instrumental and nuisance parameters are marginalized over implicitly according to their respective dataset likelihoods.

Parameter	Symbol	Prior Range ($\mathcal{U}$)
Hubble Constant	H_0	$[50, 90]$
Matter Density	Ω_m	$[0.1, 0.6]$
Supernova Calibration	M_B	$[-19.6, -19.1]$
Effective EoS Today	w_0	$[-2.5, -0.5]$
EoS Evolution	w_a	$[-1.5, 1.5]$

By employing these expansive prior bounds, we ensure that the reported decisive evidence ($\Delta \ln \mathcal{Z} = +40.3$) is driven purely by the framework's superior phenomenological alignment with the joint dataset (Pantheon+, BAO, CMB, RSD), rather than by artificially restricted or fine-tuned prior volumes.

[1] A. G. Riess *et al.*, Astron. J. **116**, 1009 (1998).
[2] S. Perlmutter *et al.*, Astrophys. J. **517**, 565 (1999).
[3] A. G. Riess *et al.*, Astrophys. J. Lett. **934**, L7 (2022).
[4] L. Verde *et al.*, Nature Astron. **3**, 891 (2019).
[5] L. Knox and M. Millea, Phys. Rev. D **101**, 043533 (2020).
[6] V. Poulin *et al.*, Phys. Rev. Lett. **122**, 221301 (2019).
[7] M. Chevallier and D. Polarski, Int. J. Mod. Phys. D **10**, 213 (2001).
[8] E. V. Linder, Phys. Rev. Lett. **90**, 091301 (2003).
[9] W. Hu and N. Sugiyama, Astrophys. J. **471**, 542 (1996).
[10] A. R. Liddle and D. H. Lyth, *Cosmological Inflation and Large-Scale Structure* (Cambridge University Press, 2000).
[11] D. Brout *et al.*, Astrophys. J. **938**, 110 (2022).
[12] Planck Collaboration, Astron. Astrophys. **641**, A6 (2020).
[13] S. Alam *et al.*, Phys. Rev. D **103**, 083533 (2021).
[14] J. S. Speagle, Mon. Not. Roy. Astron. Soc. **493**, 3132 (2020).
[15] S. K. Lam *et al.*, in *Proceedings of the Second Workshop on the LLVM Compiler Infrastructure in HPC* (2015).
[16] C. Pitrou *et al.*, Phys. Rept. **754**, 1 (2018).
[17] E. Aver, K. A. Olive, and E. D. Skillman, JCAP **2015** (07), 011.
[18] R. J. Cooke, M. Pettini, K. M. Nollett, and R. Jorgenson, Astrophys. J. **830**, 148 (2016).
[19] E. Di Valentino *et al.*, Class. Quant. Grav. **38**, 153001 (2021).
[20] C. Heymans *et al.*, Astron. Astrophys. **646**, A140 (2021).
[21] DES Collaboration, Phys. Rev. D **105**, 023520 (2022).
[22] L. Perivolaropoulos and F. Skara, New Astron. Rev. **95**, 101659 (2022).
[23] S. L. Finkelstein *et al.*, Astrophys. J. Lett. **946**, L13 (2023).



FIG. 1. Marginalized 1D and 2D posterior distributions for the ΨTMG baseline model. The convergence on $H_0 \approx 74.2$ demonstrates the reconciliation of disparate probes.



FIG. 2. Frequentist profile likelihood for H_0 within the ΨTMG framework. The blue curve represents the $\Delta\chi^2$ evolution, showing a decisive exclusion of the standard ΛCDM value (Planck18, gray dashed line) at more than 5σ confidence level. The inset provides a detailed view of the ΨTMG minimum region, with horizontal dashed lines indicating the 1σ, 2σ, and 3σ confidence thresholds for precise parameter estimation.



FIG. 3. Comprehensive tension synthesis for H_0 and S_8. The dual-panel plot highlights the ΨTMG model's ability to simultaneously reconcile local and weak-lensing measurements with CMB constraints, supported by an evidence of $\Delta\ln\mathcal{Z} = +40.3$. Numerical values correspond to the 68% confidence levels summarized in Table I.