

# AS / A Level **Physics** *Definitive*

A guide for answering questions asking “define” or “what is meant by”, as well as some other tricky questions

Contains 95 exam-style questions, each have listed correct, as well as incorrect responses.

Lynn Mao Wtm <m@maowtm.org>

Content is likely to be *somewhat* incomplete.

## Contents

<b>Preface</b>	<b>2</b>
<b>1 Dynamics and Energy</b>	<b>3</b>
<b>2 Electricity</b>	<b>9</b>
<b>3 Waves</b>	<b>15</b>
<b>4 Particle and Nuclear Physics</b>	<b>20</b>
<b>5 Measurement, Units, Estimates, etc...</b>	<b>24</b>
<b>Index</b>	<b>27</b>



This work is licensed under a Creative Commons Attribution-ShareAlike 4.0 International License.

Proudly produced with L<sup>A</sup>T<sub>E</sub>X and TikZ-source available at <https://github.com/micromaomao/A-Physics-Definitive>

This is **not** a CIE official document.

# Preface

Good to see you, physics learner!

This document is written by learners like you, and it serves a very specific purpose—to help you answer a type of exam question in CIE. I’m talking about things like:

- Define *elastic potential energy*
- State the principle of *conservation of momentum*
- State the difference between a *stationary wave* and a *progressive wave*...
- Explain the origin of *upthrust* for a body in liquid
- Explain the part played by *diffraction* and *interference* in the production of the *first order maximum* by the *diffraction grating*.
- Distinguish between an  $\alpha$ -particle and a  $\beta^+$ -particle.

You get the idea. This type of question, where it asks to *state* or *explain* something sometimes can turn out to be pretty hard, even if you do have an sound understanding of the concepts involved and nailed the calculation part. Some people hate these questions a lot. However, it doesn’t has to be this hard.

This document aims to help you get better at answering those questions by showing you how examiner wants you to answer them, and also showing you what to avoid doing. This is NOT a list for you to memorize, instead, you should seek understanding of the logic/reasoning/key ideas behind the answers shown here, which is **often** an exact copy from the relevant Mark Scheme, and think about how you could write better answers—such as making your answer more complete or concise, learning new ways and perspectives to describe/explain a things you understand, or correcting inaccuracies in your knowledge.

Answers here follows the same style as CIE Mark Scheme, only edited/rephrased to make the meaning clearer. A  $n'$  denotes that the expression before  $n'$  could gain  $n$  marks. Some common errors mentioned in examiner reports are listed here with a **NOT** prefix. This indicates that the responses could not gain full mark, and often could gain no mark at all.

It is worth reminding again that I recommend **against** anyone memorizing this or any other pre-written ‘answers’. It is a terrible waste of time and effort and does little to improve your knowledge of physics.

Happy learning and best wishes for your CIE!

Mao Wtm  
January 27, 2018

# 1 Measurement, Units, Estimates, etc...

1 For the measurement of the following items, state the name of a suitable measuring instrument.

Diameter of a wire ( $10^{-1}$  mm): micrometer (screw gauge) **or** digital calipers 1'

2 Explain the difference between accuracy and precision [2]

*Accuracy* means the closeness of the value(s)/measurement(s) to the true value 1'

*Precision* means the *range* (i.e. spread) of the values/measurements 1'

3 Describe the effects, one in each case, of systematic errors and random errors [2]

- systematic errors: the reading is larger or smaller than / varying from the true reading by a constant amount. 1'

- random errors: scatter in readings about the true reading 1'

4 Define *density* [1]

$$\frac{\text{mass}}{\text{volume}}$$
 1'

5 Make *estimates* of the following quantities:

Quantity	Lower bound	Upper bound	Unit
<i>Speed</i> of sound in air	100	900	m s <sup>-1</sup>
<i>Frequency</i> of an audible sound wave	20	20 × 10 <sup>3</sup>	Hz
<i>Density</i> of air at room temperature and pressure	0.5	1.5	kg m <sup>-3</sup>
Mass of a protractor	5	50	g
Mass of a plastic 30 cm ruler	10	100	g
Mass of a wooden metre rule	50	200	g
Mass of an apple	50	500	g
Volume of the head of an adult person	2 × 10 <sup>3</sup>	9 × 10 <sup>3</sup>	cm <sup>3</sup>
Volume of a cricket ball or a tennis ball	50	300	cm <sup>3</sup>
<i>Pressure</i> due to a depth of 10 m of water	5 × 10 <sup>4</sup>	5 × 10 <sup>5</sup>	Pa
Wavelength of red light in a vacuum	600	800	nm
Wavelength of ultraviolet radiation	10	400	nm
Diameter of an atom	10 × 10 <sup>-9</sup>	10 × 10 <sup>-11</sup>	m
Diameter of a nucleus	10 × 10 <sup>-13</sup>	10 × 10 <sup>-15</sup>	m
Thickness of a sheet of paper	0.05	0.15	mm
Time for sound to travel 100 m in air	0.25	0.5	s
Weight of 1000 cm <sup>3</sup> of water	8	12	N

6 Give examples of *SI base units*

Any *n'* from:

- kilogram
- meter
- second
- kelvin
- mole
- ampere
- candela

**NOT** *coulomb*

**NOT** gram

**NOT** temperature, *current*, etc.

7 Give examples of *SI base quantities*

Any  $n'$  from:

- mass
- length
- time
- temperature
- amount of substance
- *current*
- luminous intensity

**NOT** *charge*

**NOT** mole, kelvin, *current*, etc.

## 2 Electricity

8 *Charge...*

Quantity: *current*  $\times$  time 1'

Unit: *coulomb* = ampere second 1'

9 State what is meant by an electric *current* [1]

flow of charge carriers 1'

10 Electric *current* is a flow of charge carriers. The *charge* on the carriers is quantised. Explain what is meant by *quantised*. [1]

charge exists only in discrete amounts 1'

11 *Potential difference...*

Quantity:  $\frac{\text{work (done) or energy (transformed) (from electrical to other forms)}}{\text{charge}}$  1'

**NOT** Energy transferred *by* unit charge / 1 C.

Unit: *volt* =  $\frac{\text{joule}}{\text{coulomb}}$  1'

Not to be confused with *electromotive force*.

12	Define <i>electromotive force</i> (e.m.f.) of a cell.	[1]
	energy transformed from chemical to electrical per unit <i>charge</i>	1'
	Not to be confused with <i>potential difference</i> .	
13	<b>Resistance...</b>	
	Quantity: $\frac{\text{potential difference}}{\text{current}}$	1'
	Unit: <i>ohm</i> = $\frac{\text{volt}}{\text{ampere}}$	1'
14	Define the <i>ohm</i>	[1]
	$\frac{\text{volt}}{\text{ampere}}$	1'
	(See question 36)	
	<b>NOT</b> 'unit of <i>resistance</i> '	
	<b>NOT</b> $\frac{\text{potential difference}}{\text{current}}$ ,	
15	Explain what is meant by an <i>electric field</i>	[1]
	a region/space/area where a (stationary) <i>charge</i> experiences an (electric) <i>force</i>	1'
	<b>NOT</b> 'Force per unit charge' etc.	
16	Define <i>electric field strength</i>	[1]
	force <b>per</b> unit positive charge.	1'
	See also: <i>electric field</i>	

17

On figure 2, draw at least six field lines to represent the *electric field* between the plates. [1]

Question:






Figure 1: for question 17

Answer:

NOT




NOT



NOT

NOT

NOT



i.e. use ruler.

18	<p>Explain why the calculation of the <i>force</i> on a electron in an <i>electric field</i> does not need to include the gravitational effects on the electron. [1]</p> <p>electric force is way bigger than gravitational force (on electron)/weight 1'  <b>or</b> something about acceleration being way bigger.</p> <p><b>NOT</b> gravitational force is negligible</p>
19	<p>Describe the I-V characteristic of a metallic conductor at constant temperature [1]</p> <p>straight line <b>through the origin</b> 1'</p> <p>See figure 3</p>
20	<p>Describe the I-V characteristic of a semiconductor diode [2]</p> <p>zero current for one direction up to zero or a few tenths of volt in another direction 1'  straight line positive gradient/increasing gradient (after that) 1'</p> <p>See figure 3</p>
21	<p>Use figure 3 to describe the variation of the resistance of the diode between <math>V = -0.5\text{ V}</math> and <math>V = 0.8\text{ V}</math>. [2]</p> <p>very high/infinite <i>resistance</i> for negative <i>voltages</i> up to about <math>0.6\text{ V}</math> / some number in graph given in question. 1'  resistance decreases from <i>&lt;that voltage&gt;</i> 1'</p> <p><b>NOT</b> zero <i>current</i> means zero <i>resistance</i>  The gradient of the graph is <b>NOT</b> <i>resistance</i>.</p>
22	<p>State Kirchhoff's <i>n</i>th law...</p> <ul style="list-style-type: none"> <li>• <math>n = 1</math>: [1]  sum of currents into a junction = sum of currents out of junction 1'</li> <li>• <math>n = 2</math>: [2]  <b>total/sum</b> of electromotive forces/e.m.f.s = <b>total/sum</b> of potential differences/p.d.s 1'  around a loop/(closed) circuit 1'</li> <li>• There is no third law. (Although there is <i>a third law by Newton</i>.)</li> </ul>



////////////////////////////////////




Figure 2: *I-V graph* for question 42, 43 and 44

////////////////////////////////////

23

*Kirchhoff's first law* is linked to the conservation of a certain quantity. State this quantity. [1]

*charge*


1'

**NOT** *current*


24

Sketch the temperature characteristic of a (NTC) thermistor [2]

Answer:




**NOT**



### 3 Waves

- 25 State what is meant by the *amplitude* of a wave [1]  
the maximum *displacement* 1'



- 26 State what is meant by the *displacement* of a wave [1]  
the distance from the equilibrium position / undisturbed position / midpoint / rest position 1'

27 *Frequency...*

Def: the number of wavefronts / crests / wavelength passing a (fixed) point on the wave per unit time **or** number of oscillations of the source per unit time.

**NOT** something per *second*. See the NOTs under *velocity*

**NOT** the number of *complete* oscillations per unit time since frequency is not necessarily an integer value.

Quantity: *period*<sup>-1</sup>

Unit: Hz = s<sup>-1</sup>

Not to be confused with *period* or wavelength.

28 *Period...*

Def: time between adjacent wavefronts **or** time for one oscillation.

Quantity: *frequency*<sup>-1</sup>

Unit: s

Not to be confused with *frequency* or wavelength.

29

State the difference between a *stationary wave* and a *progressive wave* in terms of...

(i) the energy transfer along the wave: [1]

in a stationary wave energy is not transferred **or** in a progressive wave energy is transferred 1'

(ii) the phase of two adjacent vibrating particles: [1]

in a stationary wave (adjacent) particles are in phase **or** in a progressive wave (adjacent) particles are out of phase/have a phase difference/not in phase 1'

(iii) the amplitude of the particles' vibration: [1]

in a progressive wave all particles have same amplitude **or** in a stationary wave nodes have minimum / zero amplitude and antinodes have maximum amplitude (or simply 'amplitude varies for stationary wave') 1'

**Note:** 'Progressive wave being formed by one / stationary wave being formed by two waves' is **NOT** a difference and is actually not correct.

30

State what is meant by an *antinode* of the *stationary wave* [1]

Position where maximum *amplitude* 1'

31

By reference to vibrations of the points on a wave and to its direction of energy transfer, distinguish between *transverse waves* and *longitudinal waves*. [2]

Transverse waves have vibrations / displacement of particles that are perpendicular to the direction of energy travel / propagation 1'

Longitudinal waves have vibrations / displacement of particles that are parallel to the direction of energy travel / propagation 1'

**NOT** direction of motion of the wave / wave travel

32

State the conditions required for the formation of a *stationary wave* [2]

(two) waves travelling (at same speed) in opposite directions overlap 1'

waves (are same type and) have same *frequency*/wavelength 1'

33

Describe the features that are seen on the stretched string that indicate *stationary waves* have been produced. [1]

points on string have different **amplitudes** varying from maximum to zero/minimum 1'

34 Explain how *stationary waves* are formed in a tube with one end closed / with a microwave source and a metal reflector (figure 4). [2]

waves **from source** (e.g. loudspeaker) (travel down tube and) are reflected at closed end / reflector 1'

two waves (travelling) in opposite directions with same *frequency*/wavelength and *speed overlap* 1'




Figure 3: Figure for question 58

35 Explain how *D* is used to show that stationary waves are formed between reflector and wave source in figure 4 [2]

detector/*D* is moved between reflector and source 1'

maximum, minimum/zero, (maximum... etc.) observed on meter/deflections/readings /measurements/recordings 1'

**NOT** nodes and *antinodes* observed.

36 Describe the *Doppler effect* [1]

**observed frequency** is different to source *frequency* when source moves relative to observer 1'

**NOT** due to change in position of source

37 Describe what is meant by a *polarised wave* [2]

vibrations are in a single direction 1'

applies to transverse waves **or** normal to direction of wave energy travel / propagation 1'

**NOT** vibration in only one plane

38 Use the principle of *superposition* to explain <some observation> [2]

the waves (that overlap) have phase difference of  $x^\circ$  /  $y$  rad / path difference of  $z\lambda$  1'  
constructive / destructive interference **or** displacement larger / smaller (depend on question) 1'

39 State what is meant by the *diffraction* of a wave. [2]


When wave incident on/passes by/through an aperture/edge 1'  
 it spreads (into the geometrical shadow) 1'

**NOT** bending

**NOT** when the wave passes through an obstacle

40 State what is meant by *interference* / *superposition* [2]

when two (or more) waves superpose/meet/overlap 1'  
 resultant displacement is the sum of the displacement of each wave 1'



41 Explain the meaning of *coherent* [1]

constant phase difference 1'

42 Explain the part played by *diffraction* in the production of the fringes in the double slit experiment [2]

waves at (each) slit/aperture spread (into the geometric shadow) 1'  
 (the spread) wave(s) overlap/*superpose*/sum/meet/intersect 1'

43 Explain the reason why a double slit is used rather than two separate sources of light in the double slit experiment [1]

two separate light sources are not in constant phase difference/*coherence*  
**or** waves/light from the double slit are *coherent*/have a constant phase difference 1'




Figure 4: for question 67 and 68. Not to scale.

44 Explain why a bright fringe is produced at point  $X$  in figure 5. [2]

waves (from slits) overlap (at point  $X$ ) 1'

path difference (from slits to  $X$ ) is zero  
**or** phase difference (between the two waves) is zero 1'  
 (so constructive interference gives bright fringe)

**NOT** statements that applies to all bright fringes in general – e.g. path difference =  $n\lambda$  or phase difference =  $360^\circ n$ .

45 The intensity of the light passing through the two slits in figure 5 was initially the same. The intensity of the light through one of the slits is now reduced. Compare the appearance of the fringes before and after the change of intensity. [2]

Any 2' of:  
 same separation/fringe width/number of fringes  
 bright fringes/central bright fringe/(fringe at)  $X$  less bright  
 dark fringes/(fringe at)  $Y/Z$  brighter

**NOT** 'fringes' if it is not clear whether it refers to the dark fringes or the white fringes.

46 Describe the *diffraction* of light at a diffraction grating [2]

waves at the slits 1' spread (into the geometric shadow) 1'

**NOT** light spread without the word 'wave'

47 Explain the part played by *diffraction* and *interference* in the production of the first order maximum by the diffraction grating. [3]

diffraction: spreading/diverging of waves/light (takes place) at (each) slit/element /gap/aperture 1'

interference: waves (from *coherent* sources at each slit) overlap 1' with phase difference  $360^\circ$  / path difference  $\lambda$  1'

See also: question 71

48 By reference to *interference*, explain the zero and first order maximum in a diffraction grating. [3]

zero: waves (from each slit) overlap/meet/*superpose* 1' with a phase difference/path difference of zero 1'

first: phase difference is  $360^\circ$ /path difference of  $\lambda$  1'

For the first mark, explicit mentioning that the waves **meet** or otherwise interference is necessary.

See also: question 70

## 4 Dynamics and Energy

49 Define *speed* [1]

$\frac{\text{change in distance}}{\text{change in time}}$  or  $\frac{\text{distance}}{\text{time}}$  1'

*Avoid* 'distance over time'

**NOT** 'change of distance *with* time'

50 Define *velocity* [1]

rate of change of *displacement* or  $\frac{\text{change in displacement}}{\text{time (taken)}}$  1'

**NOT** rate of change of displacement per unit time

*Avoid* 'displacement over time'

**NOT** 'change of displacement *with* time'

**NOT** something with 'distance'

**NOT** displacement per second (just like **NOT** 'meter per time')

Not to be confused with *speed*.



51	Define <i>acceleration</i>	[1]
	rate of change of <i>velocity</i> <b>or</b> $\frac{\text{change in } \textit{velocity}}{\text{time (taken)}}$	1'
	<b>NOT</b> rate of change of velocity per unit time	
	<b>NOT</b> something with 'speed'	
52	Define <i>force</i>	[1]
	Rate of change of momentum	1'
	<b>NOT</b> $F = ma$ or "mass $\times$ <i>acceleration</i> "	
	Definitely <b>NOT</b> "a push or pull"—this is primary grade stuff.	
53	Define <i>power</i>	[1]
	$\frac{\textit{work} \text{ (done)}}{\text{time (taken)}}$ <b>or</b> $\frac{\text{energy transferred}}{\text{time (taken)}}$ <b>or</b> rate of <i>work</i> done	1'
	<b>NOT</b> $\frac{\text{energy}}{\text{time}}$	
	<b>Avoid</b> 'in a certain time' / 'unit of time'	
	<b>Avoid</b> 'over time'	
54	Define <i>work</i> done	[2]
	<i>force</i> $\times$ distance moved (by force)	1'
	in the direction of the force	1'
55	Explain what is meant by <i>kinetic energy</i> .	[1]
	energy/ability to do work a object/body/mass has due to its speed/velocity/motion /movement	1'
56	Define <i>potential energy</i>	[1]
	Stored energy available to do work	1'
	<b>NOT</b> description of any specific type of energy e.g. <i>gravitational</i>	
57	Define <i>gravitational potential energy</i>	[1]
	Energy due to height/position of mass <b>or</b> distance from mass <b>or</b> moving mass from one point to another.	1'
	<b>NOT</b> about 'height of a body above the Earth'	
	<b>NOT</b> about gravitational potential	

58	Define <i>elastic potential energy</i>	[1]
	Energy (stored) due to deformation/stretching/compressing/change in shape/size <b>NOT</b> 'the energy stored in an elastic body' without mentioning deformation <b>NOT</b> any formula	1'
59	State <i>Hooke's law</i> .	[1]
	force/load is proportional to extension/compression (provided proportionality limits not exceeded)	1'
60	Define the <i>Young modulus</i>	[1]
	$\frac{\text{stress}}{\text{strain}}$	1'




Figure 5: Figure for question 13

- 61 Use data from <some  $F/x$  (extension) graph> (figure 1) to show that the spring obeys *Hooke's law* for this range of extensions / compression. [2]

two values of  $F/x$  are calculated which are the same

or ratio of two forces and the ratio of the corresponding two extensions are calculated which are the same

or gradient of graph line calculated and coordinates of one point on the line used with straight line equation  $F = mx + c$  to show  $c = 0$  1'

(so) force is proportional to extension (and so *Hooke's law* obeyed) 1'

**NOT** straight line  $\Rightarrow$  Hooke's law obeyed, since line must cross origin.

- 62 Describe how to determine whether the extension of the spring is *elastic*. [1]

or

State how you would check that the spring has not exceeded its *elastic limit* [1]

remove the force/masses and if the spring returns to its original length its an elastic extension. 1'

**NOT** something about the extension being proportional to the force.

63	For the following scenario, state and explain the changes in energy that occur.	
	(a) Stuff falls through liquid:	[2]
	decrease in <i>gravitational potential energy</i> due to decrease in height (since $E_p = mgh$ )	
	increase in thermal energy due to <i>work</i> done against viscous drag	
	loss/change of (total) $E_p$ equal to gain/change in thermal energy	
	Any 2' points.	
64	State the principle of <i>conservation of momentum</i> (linear momentum)	[2]
	Sum/total momentum is constant <b>or</b> before = after	1'
	for an isolated system <b>or</b> with no (resultant) external force	1'
65	Explain what is meant by particles colliding <i>elastically</i>	[1]
	the total <i>kinetic energy</i> before (the collision) is equal to the total <i>kinetic energy</i> after (the collision)	1'
66	Define <i>strain</i>	[1]
	$\frac{\text{extension}}{\text{original length}}$	1'
67	<i>Stress</i> ...	
	Quantity: $\frac{\text{force}}{\text{cross-sectional area}}$	1'
	Unit: Pa = $\frac{\text{force}}{\text{area}}$	1'
68	State the two conditions for a system/object to be in <i>equilibrium</i>	[2]
	resultant <i>force</i> (in any direction) is zero	1'
	resultant <i>torque/moment</i> (about any point) is zero <b>or</b> sum of clockwise moment and sum of anticlockwise moment is zero	1'
	<b>NOT</b> 'no turning effect'	
	<b>NOT</b> 'the forces are balanced'/'cancel'	
	<b>NOT</b> 'no forces acting'	

69


Define the *torque* of a couple

[2]

Torque is the product of one of the *forces* ( $F$ ) and the perpendicular distance ( $d$ ) between forces.

One of the forces  $\times$  distance  $1'$

Perpendicular  $1'$




70

Define the *moment* of a *force*

[1]

*force* ( $F$ )  $\times$  **perpendicular** distance ( $d$ )  
(of line of action of force) to/from a point / pivot

 $1'$ 

71

Explain what is meant by *centre of gravity*

[1]

the point from where (all) the weight (of a body) seems to act

 $1'$ 

**NOT** weight concentrated on this point

**NOT** the point where mass acts

72

Define *pressure*

[1]

*force*

area (normal to the force)

 $1'$ 

**NOT** 'cross-sectional area'

73 Explain the origin of *upthrust* for a body in liquid [2]

Pressure / force up on bottom **greater** than pressure / force down on top 2'

1' for pressure on bottom **different** from pressure on top **or** pressure changes with depth.

**NOT** having less density than the liquid.

74 State Newton's *n*th law of motion. . .

- *n* = 1: a body/mass/object continues (at rest or) at constant/uniform *velocity* unless acted on by a **resultant force** 1'  
**NOT** 'constant speed' without mentioning straight line motion  
**NOT** 'uniform motion'
- *n* = 2: See definition of *force*
- *n* = 3: force on body A is equal in magnitude to force on body B (from A) 1', in opposite directions 1', of the same kind. 1'

75 State and explain whether momentum is conserved during the collision of <some ball or stuff> with <some floor, wall or stuff> [2]

there is a change/gain in momentum of <some floor, wall or stuff> 1'

there is an equal (and opposite) change to the momentum of <some ball or stuff> so momentum is conserved 1'

**or**

change of (total) momentum of <some floor, wall or stuff> and <some ball or stuff> is zero

**or** (total) momentum of <some ball or stuff> and <some floor, wall or stuff> before is equal to the (total) momentum after 1'  
so momentum is conserved 1'

**NOT** not conserved for any reason such as an open system.

76 Explain how the collision of two objects can support *Newton's third law* [2]

change in momentum equal (and opposite) for the two objects 1'

*force* is change in momentum over time and time (of collision) is the same hence force on the two objects are equal and opposite as for *Newton's third law*. 1'

77

In practice, air resistance is not negligible. State and explain the effect of air resistance on the time taken for a ball thrown upward to reach maximum height. [2]

deceleration is greater/resultant force (weight and friction force) is greater 1'  
take less time 1'

78

Use the *kinetic model* to explain the *pressure* exerted by gases to wall of container [3]

molecule collides with wall/container and there is a change in momentum 1'

$\frac{\text{change in momentum}}{\text{time}}$  is *force* or  $\Delta p = Ft$ . 1'

many/all/sum of molecular collisions over surface/area of container produces pressure 1'

## 5 Particle and Nuclear Physics

79

Distinguish between an  $\alpha$  *particle* and a  $\beta^+$  *particle*. [3]

Any 3' from:

- $\alpha$  is 2 protons and 2 neutrons;  $\beta^+$  is positron.
- $\alpha$  has *charge*  $+2e$ ;  $\beta^+$  has *charge*  $+e$ .
- $\alpha$  has mass  $4u$ ;  $\beta^+$  has mass  $\frac{1}{2000}u$ .
- $\alpha$  made up of *hadrons*;  $\beta^+$  made up of a *lepton*.

80

Similarity and difference between a  $\beta^+$  *particle* and a  $\beta^-$  *particle*...

Similarity: same (rest) mass, equal magnitude of *charge*.

Difference: opposite sign of *charge*, one is matter / electron and one is antimatter / antielectron / positron.

81

State the name of the force responsible for  $\beta$  decay. [1]

weak (nuclear force/interaction) 1'  
**NOT** simply 'nuclear force'

82 State the quantities that are conserved in a nuclear reaction.

Any  $n'$  from:

- mass-energy **NOT** separately ‘mass’ or ‘energy’ or ‘mass *and* energy’
- momentum
- proton number
- nucleon number
- *charge*

83 State the names of the *leptons* produced in each of the decay processes:

- $\beta^-$  decay: electron and (electron) antineutrino
- $\beta^+$  decay: positron and (electron) neutrino

84 State the name of the class (group) to which each of the following belongs. . .

electron /  $\beta$  *particle* / neutrino:

*leptons*

neutron / proton:

*hadrons* **or** baryons

**Please** avoid mis-spelling.

85 Explain why the sum of the kinetic energies of the carbon-13 nucleus and the  $\beta^+$  *particle* cannot be equal to the total energy released by the decay process  $X \longrightarrow {}^{13}_6\text{C} + \beta^+$ .

a (electron) neutrino/ $\nu_{(e)}$  is also produced (and thus has energy) 1'

86 State the composition of the proton and of the neutron in terms of quarks. [1]

- proton: up up down (no strange) /  $u u d$
- neutron: up down down (no strange) /  $u d d$

87 Give one example of . . .

- (i) *hadron*: neutron **or** proton
- (ii) *lepton*: electron **or** (electron) neutrino



88

State one difference between a *hadron* and a *lepton*

[1]

hadrons are not fundamental particle / leptons are fundamental particle

**or** hadron made of quarks/lepton not made of quarks**or** strong force/interaction acts on hadrons/does not act on leptons

1'

**NOT** comparing mass between proton and electron.**NOT** 'only leptons experience the weak force'

89

State what may be inferred from the following results in the  $\alpha$  particle scattering experiment.

- The vast majority of  *$\alpha$  particles* pass straight through the metal foil or are deviated by small angles. [1]

**most** of the atom is empty space**or** the nucleus (volume) is (very) small **compared to the atom**

1'

- A very small minority of  *$\alpha$  particles* are scattered through angles greater than  $90^\circ$ . [2]

nucleus is (positively) charged

1'


the mass/charge is concentrated / the majority of mass/charge in (very small)

nucleus / small region/volume/core

1'

When asked to state the results, avoid expressions such as 'some', 'a lot/few' or 'many' particles—use 'vast majority' or 'vast majority'.

Figure 6: Two parallel, vertical metal plates in a vacuum are connected to a power supply, and a radioactive source emitting  $\alpha$  particle is placed below the plates. This is for question 83 and 84.



90 Explain why the metal plates are placed in a vacuum in figure 6. [1]

range of  $\alpha$  particle is only few cm in air

or loss of energy of the  $\alpha$  particles due to collision with air molecules/ionisation of the air molecules 1'

91 The  $\alpha$  particle source in figure 6 is replaced by a  $\beta$  particle source. By reference to the properties of  $\alpha$  particle and  $\beta$  particle, suggest three possible differences in the deflection observed with  $\beta$  particle. [3]

$\beta$  have opposite charge to  $\alpha$  therefore deflection in opposite direction 1'

$\beta$  has a range of velocities/energies hence a number of different deflections would be seen 1'

$\beta$  have less mass or  $\frac{\text{charge}}{\text{mass}}$  is larger hence deflection is greater

or  $\beta$  with (very) high speed (may) have less deflection 1'

There must be references to properties, such as 'opposite charge'.

92 State the constituent particles of the <some element> nucleus [1]

$x$  protons and  $y$  neutrons 1'

NOT  $x$  electrons.

93	State the constituent particles of <i><math>\alpha</math> particle</i>	[1]
	2 protons and 2 neutrons	1'
	<b>NOT</b> ${}^4_2\text{He}$ / Helium / Helium nucleus	
94	Explain, using the law of <i>mass-energy conservation</i> , how energy is released in a nuclear reaction	[2]
	(total) mass on left-hand side (of equation)/reactants is greater than (total) mass on right-hand side (of equation)/products	1'
	difference in mass is (converted to) energy.	1'
95	Explain the meaning of <i>spontaneous radioactive decay</i>	[1]
	(rate of decay) not affected by any external factors or changes in temperature and pressure etc.	1'
	<b>NOT</b> decay occurred randomly / naturally	

# Index

- $\alpha$  particle, 23, 25–27
- $\alpha$  particle scattering experiment, 25
- $\beta^+$  particle, 23, 24
- $\beta^-$  particle, 23
- $\beta$  particle, 24, 26
- acceleration, 17
- accuracy, 3
- air resistance, 23
- amount of substance, 5
- ampere, 4, 6
- amplitude, 11, 12
- antimatter, 23
- antinode, 12, 13
- atom, 25
- baryon, 24
- candela, 4
- centre of gravity, 21
- charge, 5, 6, 10, 23, 24, 26
- coherent, 14, 16
- conservation of momentum, 20
- constituent particles, 26, 27
- coulomb, 4, 5
- couple, 21
- current, 4–6, 8, 10
- density, 3
- diffraction, 14–16
- diffraction grating, 15, 16
- diode, 8, 9
- displacement, 11, 16
- Doppler effect, 13
- double slit experiment, 14
- elastic, 19
- elastic collision, 20
- elastic limit, 19
- elastic potential energy, 18
- electric field, 6–8
- electric field strength, 6
- electromotive force, 5, 6
- electron, 24, 25
- energy, 5, 6, 12, 17, 27
- equilibrium, 20
- estimates, 4
- filament lamp, 9
- force, 6, 8, 17, 20–23
- forces, 21
- frequency, 11–13
- friction, 23
- gram, 4
- gravitational, 17
- gravitational potential energy, 17, 20
- hadron, 23–25
- Hooke’s law, 18, 19
- I-V graph, 9
- in phase, 12
- interference, 14, 16
- joule, 5
- kelvin, 4, 5
- kilogram, 4
- kinetic energy, 17, 20
- kinetic model, 23
- Kirchhoff’s first law, 8, 10
- Kirchhoff’s second law, 8
- length, 5
- lepton, 23–25
- longitudinal waves, 12
- luminous intensity, 5
- mass, 5, 26, 27
- mass-energy, 24
- mass-energy conservation, 27
- meter, 4
- microwave, 13
- mole, 4, 5
- moment, 20, 21
- momentum, 17, 20, 22–24
- neutron, 23, 24, 26, 27
- Newton’s first law, 22
- Newton’s second law, 22
- Newton’s third law, 22
- node, 13
- nuclear reaction, 27
- nucleon number, 24
- nucleus, 25
- ohm, 6
- out of phase, 12

period, 11  
 phase difference, 12–14  
 polarised wave, 13  
 positron, 23  
 potential difference, 5, 6  
 potential energy, 17  
 power, 17  
 precision, 3  
 pressure, 21, 23  
 progressive wave, 12  
 proton, 23–27  
 proton number, 24  
  
 quantised, 5  
 quark, 24, 25  
  
 random errors, 3  
 resistance, 6, 8, 10  
  
 second, 4  
 SI base quantities, 5  
 SI base units, 4  
 speed, 13, 16  
 spontaneous radioactive decay, 27  
 stationary wave, 12, 13  
 strain, 18, 20  
  
 stress, 18, 20  
 strong nuclear force, 25  
 superpose, 14, 16  
 superposition, 13, 14  
 systematic errors, 3  
  
 temperature, 4, 5, 10  
 thermal energy, 20  
 thermistor, 10  
 time, 5  
 torque, 20, 21  
 transverse waves, 12  
  
 upthrust, 22  
  
 vacuum, 26  
 velocity, 11, 16, 17, 22  
 viscous drag, 20  
 volt, 5, 6  
  
 wavelength, 11–13  
 weak nuclear force, 23, 25  
 Weight, 4  
 weight, 23  
 work, 5, 17, 20  
  
 Young modulus, 18