

Q. What if I vomit after taking Paracetamol?

If vomiting occurs within 30 minutes of a dose, retake the same dose. If it happens after 30 minutes, wait until the next scheduled dose.

Q. When will I feel better after taking Paracetamol?

Effects are usually felt within 30 minutes.

Q. How often can I take Paracetamol?

Limit to four doses within 24 hours, with a minimum of 4 hours between doses. Avoid taking it for more than 3 days without consulting a doctor.

Q. Does Paracetamol make babies sleepy?

No, Paracetamol does not induce sleep; it mainly reduces pain and fever in children.

Q. Is Paracetamol safe for children?

Yes, it is safe for children when used according to the doctor's instructions.

Q. Is Paracetamol an antibiotic?

No, Paracetamol is not an antibiotic; it is a pain and fever reliever.

Q. Can I take Paracetamol and ibuprofen together?

Though both are safe, they should generally not be taken together without medical advice.

Q. How long does Paracetamol take to work?

It typically begins working within 30 & 45 minutes. Consult your doctor if any side effects occur.

Q. What are the risks of Paracetamol overdose?

Overdosing can lead to severe liver damage, kidney injury, decreased platelet count, and possibly coma. Initial symptoms include nausea, vomiting, and fatigue. Seek immediate medical help in case of suspected overdose.

1 item added

[View Cart >](#)



Home



Offers



Notification



Profile