

# ML Week

## 0x03 Logistic Regression

Jeff Abrahamson

2–5 novembre 2015

# Linear regression

- Continuous output
- Normal residues

# Logistic regression

- Binary output
- Classification

# Logistic regression

- Have: continuous and discrete inputs
- Want: class (0 or 1)

# Probabilistic inspiration

$$h_{\theta}(x) = .75 \iff \text{event has 70\% of being true}$$

# Probabilistic inspiration

$$h_{\theta}(x) = \Pr(y = 1 \mid x; \theta) = 0.70$$

So this must be true:

$$\Pr(y = 0 \mid x; \theta) + \Pr(y = 1 \mid x; \theta) = 1$$

# Probabilistic inspiration

$$\text{Set } y = 1 \iff h_{\theta}(x) = \Pr(y = 1 \mid x; \theta)$$



# Probabilistic inspiration

Math review:

- $z = (\theta^T x)$
- $\theta^T x \geq 0 \iff h_\theta \geq 0.5$
- $\theta^T x \geq 0 \iff \text{predict } y = 1$

# Logistic (sigmoid) function


$$g(z) = \frac{1}{1 + e^{-z}}$$

# Logistic (sigmoid) function

$$g(z) = \frac{1}{1 + e^{-z}}$$

Exercise: plot this

# Non-linear decision boundaries



$$h_{\theta}(x) = g(\theta_0 + \theta_1 x_1 + \theta_2 x_2 + \theta_3 x_1^2 + \theta_4 x_2^2)$$

Andrew Ng

# Non-linear decision boundaries

OvA, OvR

OvO

# Cost function in logistic regression

In linear regression, we had

$$J(\theta) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$$

# Cost function in logistic regression

In linear regression, we had

$$J(\theta) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x) - y)^2$$

# Cost function in logistic regression

In linear regression, we had

$$J(\theta) = \frac{1}{2m} \sum_{i=1}^m \text{Cost}(h_{\theta}(x), y)$$



# Cost function in logistic regression

Here's a convex cost function:

$$\text{Cost}(h_{\theta}(x), y) = \begin{cases} -\log(h_{\theta}(x)) & \text{if } y = 1 \\ -\log(1 - h_{\theta}(x)) & \text{if } y = 0 \end{cases}$$

# Cost function in logistic regression

Here's a convex cost function:

$$\text{Cost}(h_{\theta}(x), y) = \begin{cases} -\log(h_{\theta}(x)) & \text{if } y = 1 \\ -\log(1 - h_{\theta}(x)) & \text{if } y = 0 \end{cases}$$

Exercise: Plot this (cost vs  $y$ ).

# Cost function in logistic regression

Here's a convex cost function:

$$\text{Cost}(h_{\theta}(x), y) = \begin{cases} -\log(h_{\theta}(x)) & \text{if } y = 1 \\ -\log(1 - h_{\theta}(x)) & \text{if } y = 0 \end{cases}$$

$$J(\theta) = \frac{1}{2m} \sum_{i=1}^m \text{Cost}(h_{\theta}(x), y)$$

# Cost function in logistic regression

Here's a convex cost function:

$$\text{Cost}(h_{\theta}(x), y) = \begin{cases} -\log(h_{\theta}(x)) & \text{if } y = 1 \\ -\log(1 - h_{\theta}(x)) & \text{if } y = 0 \end{cases}$$

$$J(\theta) = y \cdot \log(h_{\theta}(x)) + (1 - y) \cdot \log(1 - h_{\theta}(x))$$

# Gradient descent

$$\theta_j \leftarrow \theta_j - \frac{\alpha}{m} \sum_{i=1}^m \left( h_{\theta}(x^{(i)}) - y^{(i)} \right) \cdot x_j^{(i)}$$

for  $j = 1, \dots, n$

# Questions?

`purple.com/talk-feedback`