

# THE RISING SEA: CATEGORIES AND SHEAVES

JEFFREY JIANG

These are some notes + exercises I've compiled working through the first 2 chapters of Ravi Vakil's *The Rising Sea*, with the main purpose being to gain some familiarity and comfort with categories and sheaves.

## 1. CATEGORY THEORY

**Definition 1.1.** A *category*  $\mathcal{C}$  is a collection<sup>1</sup> of *objects*, denoted  $\text{Ob}(\mathcal{C})$  and a collection<sup>2</sup> *morphisms*  $\text{Hom}(A, B)$ <sup>3</sup> for every pair of objects  $A, B \in \text{Ob}(\mathcal{C})$  satisfying the following axioms:

- (1) Given morphisms  $f : A \rightarrow B$  and  $g : B \rightarrow C$ , there is a unique map  $g \circ f : A \rightarrow C$  that makes the following diagram commute

$$\begin{array}{ccccc} A & \xrightarrow{f} & B & \xrightarrow{g} & C \\ & \searrow & & \nearrow & \\ & & g \circ f & & \end{array}$$

- (2) For every object  $A \in \text{Ob}(\mathcal{C})$ , there exists an *identity morphism*  $\text{id}_A \in \text{Hom}(A, A)$  such that for any morphisms  $f : A \rightarrow B$  and  $g : C \rightarrow A$ , we have that  $\text{id}_A \circ f = f$  and  $g \circ \text{id}_A = g$

A morphism  $f : A \rightarrow B$  is an *isomorphism* if there exists a morphism  $g : B \rightarrow A$  such that  $f \circ g = \text{id}_B$  and  $g \circ f = \text{id}_A$ . We then call  $g$  the *inverse* to  $f$ . Isomorphisms  $A \rightarrow A$  are called *automorphisms* of  $A$ .

**Example 1.2.** The category of sets, often denoted  $\text{Set}$  has sets as its objects, and maps of sets as its morphisms.

**Example 1.3.** Vector spaces over a field  $\mathbb{F}$  also form a category, denoted  $\text{Vec}_{\mathbb{F}}$ , where the objects are  $\mathbb{F}$ -vector spaces, and the morphisms are  $\mathbb{F}$ -linear maps.

**Exercise 1.4.** Let  $A$  be an object of a category  $\mathcal{C}$ . Show that the automorphisms of  $\text{Hom}(A, A)$  form a group, called the *Automorphism group* of  $A$ . Show that two isomorphic objects in  $\mathcal{C}$  have isomorphic automorphism groups.

*Proof.* Verifying that  $\text{Aut}(A)$  is a group is mainly an exercise in definition chasing. Associativity comes from the category axioms, the identity element is the identity morphism, and inverses exist by the definition of the automorphism group.

For the second part, let  $A, B \in \text{Ob}(\mathcal{C})$  be isomorphic, with  $f : A \rightarrow B$  an isomorphism. We then define a map  $f^* : \text{Aut}(B) \rightarrow \text{Aut}(A)$  where  $f^*\sigma = f^{-1} \circ \sigma \circ f$ . This is clearly a group homomorphism, and is an isomorphism with inverse given by  $(f^{-1})^*$ . ■

Now that we have categories, the natural thing to study are maps of categories.

**Definition 1.5.** A (covariant) *functor*  $\mathcal{F}$  from a category  $\mathcal{C}$  to another category  $\mathcal{D}$  assigns each object  $A \in \text{Ob}(\mathcal{C})$  an object  $\mathcal{F}(A) \in \text{Ob}(\mathcal{D})$  and to each map  $f : A \rightarrow B$  in  $\mathcal{C}$  a map  $\mathcal{F}(f) : \mathcal{F}(A) \rightarrow \mathcal{F}(B)$  such that  $\mathcal{F}$  respects composition, i.e.

$$\mathcal{F}(f \circ g) = \mathcal{F}(f) \circ \mathcal{F}(g)$$

a *contravariant functor* can be defined similarly, except it reverses the arrows, i.e.  $\mathcal{F}(f)$  is now a map  $B \rightarrow A$ , rather than  $A \rightarrow B$ .

**Example 1.6.** For mathematical objects that are just sets with extra structure (e.g. vector spaces, groups, rings, etc), the *forgetful functor* is the functor that takes each object to its underlying set, and each map to itself (thought of as a map of sets). Categories that admit a forgetful functor into  $\text{Set}$  are called *concrete categories*.

<sup>1</sup>Loosely speaking; there's some set-theoretic issues here, but it's not that important for us

<sup>2</sup>Again, ignoring set-theoretic problems

<sup>3</sup>Vikhil uses  $\text{Mor}$ , but we'll use the more standard notation of  $\text{Hom}$

**Example 1.7 (The functor of points).** For a category  $\mathcal{C}$ , fix an object  $A \in \text{Ob}(\mathcal{C})$ . We use this to define the functor  $\mathcal{F}_A : \mathcal{C} \rightarrow \text{Set}$ , where for  $B \in \text{Ob}(\mathcal{C})$ , we let  $\mathcal{F}(B) = \text{Hom}(B, A)$  and for  $f : B \rightarrow C$ , we let  $\mathcal{F}_A(f) : \text{Hom}(C, A) \rightarrow \text{Hom}(B, A)$  be the map where  $\mathcal{F}_A(f)(g) = g \circ f$ .

Like functions, we can compose functors  $\mathcal{F} : \mathcal{A} \rightarrow \mathcal{B}$  and  $\mathcal{G} : \mathcal{B} \rightarrow \mathcal{C}$  to obtain  $\mathcal{G} \circ \mathcal{F} : \mathcal{A} \rightarrow \mathcal{C}$ , where for  $A \in \text{Ob}(\mathcal{A})$ , we have that  $\mathcal{G} \circ \mathcal{F}(A) = \mathcal{G}(\mathcal{F}(A))$ , and the same thing for morphisms in  $\mathcal{A}$ . Also like functions, we have notions of injectivity and surjectivity.

**Definition 1.8.** A covariant functor  $\mathcal{F} : \mathcal{C} \rightarrow \mathcal{D}$  is *faithful* if the induced map  $\text{Hom}_{\mathcal{C}}(A, B) \rightarrow \text{Hom}_{\mathcal{D}}(\mathcal{F}(A), \mathcal{F}(B))$  is injective and *full* if it is surjective.

**Example 1.9.** The forgetful functor  $\text{Vec}_k \rightarrow \text{Set}$  is not full, since there are more maps of sets than there are maps of vector spaces, since an element of  $\text{Hom}_{\text{Set}}(V, W)$  need not be linear. However, the “inclusion” functor  $i : \text{Ab} \rightarrow \text{Grp}$  from abelian groups into groups is full. Given two abelian groups  $G$  and  $H$ , we clearly have that  $\text{Hom}_{\text{Ab}}(G, H) = \text{Hom}_{\text{Grp}}(G, H)$ , so we have that  $i$  is a faithful functor as well.

Functors in a lot of ways act just like functions, but there’s some more things we can do with them. For example, in some sense, we can have maps between functors.

**Definition 1.10.** Given covariant functors  $\mathcal{F}, \mathcal{G} : \mathcal{A} \rightarrow \mathcal{B}$ , a *natural transformation*  $\mathcal{F} \rightarrow \mathcal{G}$  assigns each object  $A \in \text{Ob}(\mathcal{A})$  a morphism  $m_A : \mathcal{F}(A) \rightarrow \mathcal{G}(A)$ , such that for every morphism  $f : A \rightarrow B$  in  $\mathcal{A}$ , the following diagram commutes

$$\begin{array}{ccc} \mathcal{F}(A) & \xrightarrow{\mathcal{F}(f)} & \mathcal{F}(B) \\ m_A \downarrow & & \downarrow m_B \\ \mathcal{G}(A) & \xrightarrow{\mathcal{G}(f)} & \mathcal{G}(B) \end{array}$$

a *natural isomorphism* is when all the given morphisms  $m_A$  are isomorphisms.

**Example 1.11.** A trivial example of a natural transformation between  $\mathcal{F} \rightarrow \mathcal{G}$  assigns the zero map  $z : \mathcal{F}(A) \rightarrow \mathcal{G}(A)$  for every object  $A$ . Such a zero map only exists in some categories, such as  $\text{Vec}_k$  and  $\text{Grp}$ .

This gives a rigorous definition for the words “canonical” and “natural” that get thrown around a bit too often. The prototypical example for this idea is the natural isomorphism from a finite dimensional vector space to its double dual.

**Exercise 1.12.** Let  $\mathcal{D} : \text{Vec}_k \rightarrow \text{Vec}_k$  be the functor that maps each  $k$ -vector space  $V$  to its dual space  $V^*$  of linear functions  $V \rightarrow k$  and maps a linear map  $A : V \rightarrow W$  to its pullback (sometimes called the transpose)  $A^* : W^* \rightarrow V^*$ . Show that the double dual functor  $\mathcal{D} \circ \mathcal{D}$  is naturally isomorphic to the identity functor  $\text{id}$ .

*Proof.* For a vector  $v$  in a vector space  $V$ , define the map  $\xi_v : V^* \rightarrow k$  by  $\xi_v(\omega) = \omega(v)$ . Let  $\Xi_V : V \rightarrow V^{**}$  be the map that sends  $v \mapsto \xi_v$ . We claim that the morphisms  $\Xi_V$  define a natural isomorphism from  $\mathcal{D} \circ \mathcal{D}$  to  $\text{id}$ . We first show that  $\Xi_V$  defines an isomorphism. Since  $V$  and  $V^{**}$  are the same dimension, it suffices to check that  $\Xi_V$  has trivial kernel. Suppose  $v \mapsto \xi_v = 0$ . Then  $\omega(v) = 0$  for all  $\omega \in V^*$ , which is only true when  $v = 0$ . Therefore,  $\Xi_V$  is an isomorphism for every  $V$ . Showing that the  $\Xi_V$  define a natural isomorphism now amounts to showing that the following diagram commutes

$$\begin{array}{ccc} V & \xrightarrow{A} & W \\ \Xi_V \downarrow & & \downarrow \Xi_W \\ V^{**} & \xrightarrow{A^{**}} & W^{**} \end{array}$$

which amounts to showing that  $\Xi_W \circ A = A^{**} \circ \Xi_V$ . Let  $v \in V$  and  $\omega \in W^*$ . Then we compute

$$\begin{aligned} (\Xi_W \circ A)(v)(\omega) &= \Xi_W(Av)(\omega) \\ &= \xi_{Av}(\omega) \\ &= \omega(Av) \end{aligned}$$

We also compute

$$\begin{aligned}
 (A^{**} \circ \Xi_V)(v)(\omega) &= A^{**}\xi_v(\omega) \\
 &= \xi_v(A^*\omega) \\
 &= A^*\omega(v) \\
 &= \omega(Av)
 \end{aligned}$$

So we have given a natural isomorphism  $\mathcal{D} \circ \mathcal{D} \rightarrow \text{id}$  ■

It's also somewhat easy in this language why there is no natural isomorphism from a vector space to its dual. When you saw it, there was most likely some hand-waving about how the choice of a basis made the isomorphism unnatural, so we can make this rigorous now.

**Proposition 1.13.** *There exists no natural isomorphism from the dual functor  $\mathcal{D}$  to the identity functor  $\text{id}$*

*Proof.* Suppose such a natural isomorphism existed, and assigned each vector space an isomorphism  $\varphi_V : V \rightarrow \vec{v}^*$  such that the following diagram commutes for any linear map  $A : V \rightarrow W$ .

$$\begin{array}{ccc}
 V & \xrightarrow{A} & W \\
 \varphi_V \downarrow & & \downarrow \varphi_W \\
 V & \xleftarrow{A^*} & W
 \end{array}$$

If we let  $A$  be the zero map, we have that  $A^* \circ \varphi_W \circ A = 0$ , so it cannot be  $\varphi_V$ . ■

One product from category theory that ends up being extremely useful is the notion of a **universal property**. One example you may have seen before is the universal property of a product—Given two objects  $A, B$  in some category  $\mathcal{C}$ , their product, denoted  $A \times B \in \text{Ob}(\mathcal{C})$  is (up to unique isomorphism) the unique object with maps  $\pi_A : A \times B \rightarrow A$  and  $\pi_B : A \times B \rightarrow B$  such that given any pair of maps  $f : P \rightarrow A$  and  $g : P \rightarrow B$  factor uniquely through  $A \times B$ , i.e. there exists a unique map  $f \times g : P \rightarrow A \times B$  such that the following diagram commutes

$$\begin{array}{ccc}
 P & \xrightarrow{f} & A \\
 g \downarrow & \searrow f \times g & \uparrow \pi_A \\
 B & \xleftarrow{\pi_B} & A \times B
 \end{array}$$

Some other useful objects defined by universal properties

**Definition 1.14.** An **initial object**  $I$  in a category  $\mathcal{C}$  if there exists exactly one map  $I \rightarrow A$  for every  $A \in \text{Ob}(\mathcal{C})$ . A **final object** is an object  $F$  in which there exactly one map  $A \rightarrow F$  for every object  $A$ . An object  $Z$  is a **zero object** if it is both an initial and final object.

**Exercise 1.15.** Show any two initial objects are uniquely isomorphic. Show any two final objects are uniquely isomorphic

*Proof.* Let  $A, B$  be initial objects. We note that the definition of an initial object implies that the only map  $A \rightarrow A$  and  $B \rightarrow B$  must be  $\text{id}_A$  and  $\text{id}_B$  respectively. Therefore, the unique maps  $A \rightarrow B$  and  $B \rightarrow A$  must compose to identity, so they are isomorphisms.

The same proof applies to final objects. For final objects the unique maps  $A \rightarrow A$  and  $B \rightarrow B$  are necessarily the identity maps, and the unique maps  $A \rightarrow B$  and  $B \rightarrow A$  must compose to identity. ■

For concreteness, we should determine what these some of objects are

**Exercise 1.16.** What are the initial/final objects in  $\text{Set}$ ,  $\text{Ring}$ <sup>4</sup> and  $\text{Top}$ ?

*Solution.*

<sup>4</sup>Since this book is for algebraic geometry, all rings are commutative and unital. Ring maps are required to map  $1 \mapsto 1$

- (1) The initial object for  $\mathbf{Set}$  is the empty set. Given any set  $S$ , we have a unique map  $\emptyset \rightarrow S$  (the empty map). The final object is the single point set  $*$ . Given any set  $S$ , there exists only a single map to  $*$ , namely the map that takes all of  $S$  to the point. We note that the empty set cannot be a final object, since we cannot map a nonempty set to an empty set.
- (2) The initial object in  $\mathbf{Ring}$  is the integers  $\mathbb{Z}$ . Given any commutative unital ring  $R$  (which from now on we'll just say ring), the unique map  $\mathbb{Z} \rightarrow R$  is entirely determined by the map  $1 \mapsto 1$  since 1 generates  $\mathbb{Z}$  as an abelian group. The final object is the zero ring  $0 = 1$ , where the unique map  $R \rightarrow 0$  is the map that sends everything to 0.
- (3) The initial and final objects for  $\mathbf{Top}$  are the same as for  $\mathbf{Set}$ . We note that the empty set and a single point set have unique topologies, so we do not need to specify a topology.

■

One reason that universal properties are nice is that they give an extremely helpful way in proving that some object is the one we desire – by showing it satisfies the universal property. Given an arbitrary set  $P$ , how can I conclude that it is the product  $A \times B$ ? We can look for an explicit isomorphism, which can be difficult, or we can simply show that it satisfies the same universal property as the product. Another good example of a universal property comes from an important concept in algebraic geometry – the localization of a ring.

**Example 1.17 (Localization of a ring).** Given a ring  $R$  and a multiplicative subset  $S \subset R$  (a set closed under multiplication that contains 1). We can then define the ring  $S^{-1}R$  to be the set of formal fractions  $\{r/s : r \in R, s \in S\}$  modulo the equivalence relation  $\sim$  where  $r/s \sim p/q$  if and only if there exists  $t \in S$  such that  $t(qr - sp) = 0$ <sup>5</sup>. Addition and multiplication of fractions is exactly the same as in  $\mathbb{Q}$ , namely  $r/s + p/q = (rq + ps)/pq$  and  $r/s \cdot p/q = rp/sq$ . We then have a canonical map  $R \rightarrow S^{-1}R$  where  $r \mapsto r/1$ .

**Exercise 1.18.** Show the canonical map  $R \rightarrow S^{-1}R$  is injective if and only if  $S$  contains no zero divisors.

*Proof.* Suppose there exists two distinct elements  $p, q$  that map to the same element, i.e.  $p/1 \sim q/1$ . This means that there exists some  $s \in S$  such that  $s(p - q) = 0$ , which is true if and only if  $s$  is a zero divisor for the nonzero element  $p - q$ . ■

**Exercise 1.19.** Show that  $S^{-1}R$  satisfies the following universal property:  $S^{-1}R$  is initial among  $R$ -algebras  $A$  where every element of  $S$  is mapped to an invertible element. In other words any map  $R \rightarrow A$  in which every element is mapped to an invertible element factors uniquely through the map  $R \rightarrow S^{-1}R$ .

*Proof.* Let  $\varphi : R \rightarrow A$  be such a map where every element of  $S$  maps to an invertible element. Then we want to find a map  $\tilde{\varphi} : S^{-1}R \rightarrow A$  such that the following diagram commutes

$$\begin{array}{ccc} R & \xrightarrow{\varphi} & A \\ \downarrow \iota & \nearrow \tilde{\varphi} & \\ S^{-1}R & & \end{array}$$

where  $\iota : R \rightarrow S^{-1}R$  is the canonical map. We note that in order for the diagram to commute, we must necessarily have that  $\tilde{\varphi}(r/1) = \varphi(r)$  for every  $r \in R$ . Therefore, the only possible mapping for  $\tilde{\varphi}$  must be  $\tilde{\varphi}(r/s) = \varphi(r)/\varphi(s)$ , which is possible since  $\varphi(s)$  is invertible, so we have found our unique map. ■

You can localization for  $R$ -modules  $M$  as well. Let's define it in terms of a universal property this time. Let  $S^{-1}M$  be the  $R$ -module equipped with the map  $\varphi : M \rightarrow S^{-1}M$  where any map  $f : M \rightarrow N$  such that scalar multiplication  $M_s : N \rightarrow N$  is an isomorphism factors through  $S^{-1}M$  as a unique map  $\tilde{f} : S^{-1}M \rightarrow N$ , i.e. the following diagram commutes

$$\begin{array}{ccc} M & \xrightarrow{f} & N \\ \downarrow \varphi & \nearrow \tilde{f} & \\ S^{-1}M & & \end{array}$$

<sup>5</sup>This might look strange if you compare it to the fractions you know in  $\mathbb{Q}$ , but special care needs to be taken in the case that  $R$  contains zero divisors

**Exercise 1.20.** Prove that the localization  $(S^{-1}M, \varphi)$  exists.

*Proof.* Define the localization  $S^{-1}M$  similarly to  $S^{-1}R$  as the set

$$\left\{ \frac{m}{s} : m \in M, s \in S \right\} / \sim$$

modulo the relation  $\sim$  where  $m/s \sim n/p$  if there exists  $t \in S$  such that  $t(mp - ns) = 0$ , and let  $\varphi$  be the map  $m \mapsto m/1$ . We claim that this satisfies the universal property given above. Let  $f : M \rightarrow N$  be a map of modules where  $S$  acts on  $N$  by isomorphisms. Then define  $\tilde{f} : S^{-1}M \rightarrow N$  by  $\tilde{f}(m/s) = 1/s \cdot f(m)$ , where  $1/s$  is the inverse to the multiplication map  $m_s : R \rightarrow R$ , which is defined since scalar multiplication by  $s$  defines an isomorphism  $N \rightarrow N$ . It is clear that the diagram commutes, so the set and map we gave satisfies the universal property. ■

Another example is the tensor product  $\otimes$ , where given  $R$ -modules  $M$  and  $N$ , we can form  $M \otimes_R N$  as the set of sums formal symbols  $m \otimes n$  where  $m \in M$  and  $n \in N$  where  $\otimes$  is bilinear, i.e.  $(am_1 + bm_2) \otimes n = am_1 \otimes n + bm_2 \otimes n$  and  $m \otimes (an_1 + bn_2) = am \otimes n_1 + bm \otimes n_2$ .

**Exercise 1.21.** Show that  $\mathbb{Z}/10\mathbb{Z} \otimes_{\mathbb{Z}} \mathbb{Z}/12\mathbb{Z} \cong \mathbb{Z}/2\mathbb{Z}$ .

*Proof.* Let  $[a]_{10}$  denote the equivalence class of  $a \in \mathbb{Z}$  in  $\mathbb{Z}/10\mathbb{Z}$  and define  $[a]_{12}$  similarly. Then define the map  $\varphi : \mathbb{Z}/10\mathbb{Z} \otimes_{\mathbb{Z}} \mathbb{Z}/12\mathbb{Z} \rightarrow \mathbb{Z}/2\mathbb{Z}$  by  $[a]_{10} \otimes [b]_{12} \mapsto [ab]_2$  and extending linearly to sums. This is clearly surjective and  $\mathbb{Z}$ -linear, so to show that  $\varphi$  is an isomorphism, we show that the kernel is trivial. To show this, it suffices to check elements of the form  $[a]_{10} \otimes [b]_{12}$ . Suppose we have  $[a]_{10} \otimes [b]_{12} \mapsto 0$ . Noting that  $[a]_{10} = a[1]_{10}$ , and the same for  $[b]_{12}$ . Therefore, we have that  $[a]_{10} \otimes [b]_{12} = ab([1]_{10} \otimes [1]_{12})$ . Therefore in order for  $\varphi([a]_{10} \otimes [b]_{12}) = 0$ , we must have  $ab = 0$ , since  $[1]_{10} \otimes [1]_{12} \mapsto [1]_2$ . Therefore,  $[a]_{10} \otimes [b]_{12} = 0$ , so the kernel is trivial. ■

**Exercise 1.22.** Fix an  $R$ -module  $N$ . Define a functor  $\mathcal{N} : \text{Mod}_R \rightarrow \text{Mod}_R$  by  $\mathcal{N}(M) = M \otimes_R N$ , and for any map  $\varphi : A \rightarrow B$ , let  $\mathcal{N}(\varphi) : A \otimes_R N \rightarrow B \otimes_R N$  be the map defined by  $\mathcal{N}(\varphi)(a \otimes n) = \varphi(a) \otimes n$ . Show that  $\mathcal{N}$  is indeed a covariant functor, and that it is **right-exact**, i.e. if we are given a short exact sequence of  $R$ -modules

$$0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$$

the corresponding sequence

$$0 \rightarrow A \otimes_R N \rightarrow B \otimes_R N \rightarrow C \otimes_R N$$

is also exact.

*Proof.* To show that  $\mathcal{N}$  defines a functor, all we need to show is that it respects composition. Let  $\varphi : A \rightarrow B$  and  $\psi : B \rightarrow C$ . Then we need to show that  $\mathcal{N}(\psi \circ \varphi) = \mathcal{N}(\psi) \circ \mathcal{N}(\varphi)$ . To show this, it suffices to show that they agree on elements of the form  $a \otimes n$ . We compute

$$\mathcal{N}(\psi \circ \varphi)(a \otimes n) = (\psi \circ \varphi)(a) \otimes n = \mathcal{N}(\psi \circ \varphi)(a \otimes n)$$

so  $\mathcal{N}$  is a functor.

We now show that  $\mathcal{N}$  is right exact. Let

$$0 \longrightarrow A \xrightarrow{\varphi} B \xrightarrow{\psi} C \longrightarrow 0$$

be exact, i.e.  $\ker \psi = \text{Im } \varphi$ ,  $\varphi$  is injective, and  $\psi$  is surjective. We want to show that

$$0 \longrightarrow A \otimes_R N \xrightarrow{\mathcal{N}(\varphi)} B \otimes_R N \xrightarrow{\mathcal{N}(\psi)} C \otimes_R N \longrightarrow 0$$

Is also exact, i.e.  $\mathcal{N}(\varphi)$  is injective,  $\ker \mathcal{N}(\psi) = \text{Im } \mathcal{N}(\varphi)$ , and  $\mathcal{N}(\psi)$  is surjective. We note that if  $\mathcal{N}(\varphi)(a \otimes n) = 0$ , then this means that  $\varphi(a) \otimes n = 0$ , which means either  $\varphi(a) = 0$  or  $n = 0$ . Since  $\varphi$  is injective, this implies that either  $a$  or  $n$  is 0, so  $a \otimes n = 0$ . Therefore,  $\mathcal{N}(\varphi)$  is also injective. Next, we want to show that  $\mathcal{N}(\varphi)$  surjects onto  $\ker \mathcal{N}(\psi)$ . To do this, we first characterize the elements in  $\ker \mathcal{N}(\psi)$ . Suppose  $\mathcal{N}(\psi)(b \otimes n) = \psi(b) \otimes n = 0$ . Then we must have that either  $\psi(b) = 0$  or  $n = 0$ . Therefore, the kernel of  $\mathcal{N}(\psi)$  is exactly  $\ker \psi \otimes_R N$ . We note that the image of  $\mathcal{N}(\varphi)$  is exactly  $\text{Im } \varphi \otimes_R N$ , and since the original sequence is exact, this gives us that  $\text{Im } \mathcal{N}(\varphi) = \ker \mathcal{N}(\psi)$ . Finally, we must show that  $\mathcal{N}(\psi)$  is surjective. We note that the image of  $\mathcal{N}(\psi)$  is exactly  $\text{Im } \psi \otimes_R N = C \otimes_R N$  since  $\psi$  was surjective. Therefore, the sequence is exact. ■

The characterization we gave for the tensor product  $A \otimes B$ , while not wrong, is pretty inelegant, and obfuscates the real reason we like tensor products. It is much better to define the tensor product in terms of the universal property it satisfies. In this case, given  $R$ -modules  $A, B$ , the tensor product  $A \otimes B$  is an  $R$ -module equipped with a bilinear map  $\varphi : A \times B \rightarrow A \otimes B$  such that for any bilinear map  $\psi : A \times B \rightarrow C$ , it factors uniquely as a linear map  $A \otimes B$ , i.e. the following diagram commutes

$$\begin{array}{ccc} A \times B & \xrightarrow{\psi} & C \\ \varphi \downarrow & \nearrow & \\ A \otimes B & & \end{array}$$

So the tensor product is another  $R$ -module in which linear maps are equivalent to bilinear maps out of  $A \times B$ . The construction we gave above equivalent, with the map  $\varphi$  being given by the mapping  $(a, b) \mapsto a \otimes b$ .

**Exercise 1.23.**

- (1) Let  $M$  be an  $A$  module, and let  $\varphi : A \rightarrow B$  be a ring homomorphism. Give  $B \otimes_A M$  the structure of a  $B$ -module. Show that this defines a functor  $\text{Mod}_A \rightarrow \text{Mod}_B$ .
- (2) Furthermore, if we are given another ring homomorphism  $\rho : A \rightarrow C$ , show that we can endow  $B \otimes_A C$  the structure of a ring.

*Proof.*

- (1) We note that the homomorphism  $\varphi : A \rightarrow B$  gives  $B$  the structure of an  $A$ -module where the scalar multiplication is given by  $a \cdot b = \varphi(a)b$ , so it makes sense to construct  $B \otimes_A M$ . To make  $B \otimes_A M$  a  $B$ -module, we just need to specify a scalar multiplication by  $B$ . Let  $b \otimes m \in B \otimes_A M$ , and let  $\lambda \in B$ . Let

$$\lambda \cdot b \otimes m = \lambda b \otimes m$$

and extending linearly to sums. By definition this distributes over the addition of  $B \otimes_A M$  as an  $A$ -module, so this makes it a  $B$ -module as well. We note by the previous exercise, tensoring with a fixed  $A$ -module (in this case  $B$ ), defines an functor.

*Remark.* This process is commonly referred to as *extension of scalars*, especially when the map  $\varphi$  is like an inclusion, e.h.  $\mathbb{R} \hookrightarrow \mathbb{C}$ . In that case, this process takes a real vector space and complexifies it into a complex vector space.

- (2) Let addition on  $B \otimes_A C$  be the standard addition as  $A$ -modules. Then we define multiplication as

$$(b_1 \otimes c_1)(b_2 \otimes c_2) = b_1 b_2 \otimes c_1 c_2$$

and extending linearly to sums, which covers distributivity. In addition, multiplication is commutative since  $B$  and  $C$  are commutative, so this forms a ring. ■

**Exercise 1.24.** Let  $S$  be a multiplicative subset of  $A$ , and let  $M$  be an  $A$ -modules. Describe a natural isomorphism  $(S^{-1}A) \otimes_A M \cong S^{-1}M$ .

*Proof.* We already have that the assignment  $M \mapsto (S^{-1}A) \otimes_A M$  defines a functor, and it is simple to check that the assignment  $M \mapsto S^{-1}M$  also defines a functor, so we are looking for isomorphisms  $\varphi_M : (S^{-1}A) \otimes_A M \rightarrow S^{-1}M$  such that the following diagram commutes for any  $A$ -module homomorphism  $\psi : M \rightarrow N$

$$\begin{array}{ccc} (S^{-1}A) \otimes_A M & \xrightarrow{\tilde{\psi}} & (S^{-1}A) \otimes_A N \\ \varphi_M \downarrow & & \downarrow \varphi_N \\ S^{-1}M & \xrightarrow{\psi} & S^{-1}N \end{array}$$

Where  $\tilde{\psi}$  denotes the map where  $r/s \otimes m \mapsto r/s \otimes \psi(m)$  and  $\bar{\psi}$  denotes the map where  $m/s \mapsto \psi(m)/s$ . Define  $\varphi_M : (S^{-1}A) \otimes_A M \rightarrow S^{-1}M$  by

$$\frac{r}{s} \otimes m \mapsto \frac{rm}{s}$$

we note that the kernel is trivial, since  $rm/s = 0$  if and only if  $r = 0$  or  $m = 0$ . In addition, the map is surjective, since we have that  $1/s \otimes m \mapsto m/s$ , so this defines an isomorphism. Finally, to show that the diagram commutes, we want to show that  $\varphi_N \circ \tilde{\psi} = \tilde{\psi} \circ \varphi_M$ . We compute

$$\begin{aligned} (\varphi_N \circ \tilde{\psi})(r/s \otimes m) &= \varphi_N(r/s \otimes \psi(m)) \\ &= r\psi(m)/s \\ (\tilde{\psi} \circ \varphi_M)(r/s \otimes m) &= \tilde{\psi}(rm/s) \\ &= \psi(rm)/s \\ &= r\psi(m)/s \end{aligned}$$

So the diagram commutes, and we have specified a natural isomorphism. ■

**Example 1.25 (Fibered products).** In any category, suppose we have morphisms  $\alpha : X \rightarrow Z$  and  $\beta : Y \rightarrow Z$ . The *fibered product* is another object, denoted  $X \times_Z Y$  equipped with maps  $\pi_X : X \times_Z Y \rightarrow X$  and  $\pi_Y : X \times_Z Y \rightarrow Y$  such that  $\alpha \circ \pi_X = \beta \circ \pi_Y$ , and for any object  $W$  with maps  $W \rightarrow X$  and  $W \rightarrow Y$  whose compositions with  $\alpha$  and  $\beta$  agree, there exists a unique map  $W \rightarrow X \times_Z Y$ . This is summarized by the following commutative diagram

$$\begin{array}{ccc} W & \xrightarrow{\quad} & Y \\ & \searrow \quad \nearrow & \\ & X \times_Z Y & \\ & \swarrow \quad \searrow & \\ X & \xrightarrow{\quad} & Z \end{array}$$

$\alpha$

In many cases where the maps  $\alpha$  and  $\beta$  are clear, they are omitted, and we refer to the fibered product as just  $X \times_Z Y$ . The commutative square

$$\begin{array}{ccc} X \times_Z Y & \xrightarrow{\pi_X} & X \\ \pi_Y \downarrow & & \downarrow \alpha \\ Y & \xrightarrow{\beta} & Z \end{array}$$

is often referred to a Cartesian diagram/square.

**Exercise 1.26.** Show that in the category **Set**, the fibered product is given by

$$X \times_Z Y = \{(x, y) \in X \times Y : \alpha(x) = \beta(y)\}$$

*Proof.* It suffices to show that this satisfies the universal property described above. Suppose we are given a set  $W$  with maps  $f : W \rightarrow X$  and  $g : W \rightarrow Y$  such that  $\alpha \circ f = \beta \circ g$ . Then we have a map  $\varphi : W \rightarrow X \times_Z Y$  where  $\varphi(w) = (f(w), g(w))$ , which lands in  $X \times_Z Y \subset X \times Y$  by the definition, and visibly makes the diagram above commute. ■

**Exercise 1.27.** Let  $X$  be a topological space, and show that the fibered product always exists in the category of open sets of  $X$  by describing what it is

*Proof.* We recall that the category of open sets is the category where the objects are open sets of  $X$ , with a unique arrow existing between open sets  $U$  and  $V$  if  $U \subset V$ . Then let  $A, B, C$  be open sets of  $X$ , where we have arrows  $A \rightarrow C$  and  $B \rightarrow C$ , which in more standard language means that  $A \subset C$  and  $B \subset C$ . Then for any  $W$  such that  $W \subset A$  and  $W \subset B$ , there must exist an arrow  $W \rightarrow A \times_C B$  so  $W \subset A \times_C B$ , and we have that  $A \times_C B \subset A$  and  $A \times_C B \subset B$ . Therefore,  $A \times_C B = A \cap B$ . ■

**Exercise 1.28.** Let  $Z$  denote the final object in a category. Assuming a relevant fibered products exist, show that for any objects  $X, Y$ , the fibered product  $X \times_Z Y$  is uniquely isomorphic to the product  $X \times Y$ .

*Proof.* We note that since there only exists a single map  $X \rightarrow Z$  and  $Y \rightarrow Z$ , we don't need to specify the maps. Let  $X \times Y$  be the product (in the categorical sense). Then by the universal property, we have maps  $\pi_X$  and  $\pi_Y$  such that for any object  $P$  with maps  $f : P \rightarrow X$  and  $g : P \rightarrow Y$ , we get a map  $f \times g : P \rightarrow X \times Y$ . We note that these maps also satisfy the condition for a map to exist from  $P \rightarrow X \times_Z Y$ . We then know that there exists a unique map  $X \times Y \rightarrow Z$ , since  $Z$  is a final object. Then since all maps to  $Z$  must compose to the unique map to  $Z$ , we have that  $X \times Y$  satisfies the universal property, so it must be the fibered product as well. ■

**Exercise 1.29.** Show that if the two squares in the following diagrams are commutative, then the outer rectangle is commutative as well

$$\begin{array}{ccc} U & \longrightarrow & V \\ \downarrow & & \downarrow \\ W & \longrightarrow & X \\ \downarrow & & \downarrow \\ Y & \longrightarrow & Z \end{array}$$

*Proof.* Showing that the outer rectangle commutes equates to proving that the composition  $U \rightarrow W \rightarrow Y \rightarrow Z$  is the same as  $U \rightarrow V \rightarrow X \rightarrow Z$ . From the commutativity of the upper square, we have that  $U \rightarrow W \rightarrow X \rightarrow Z$  is the same as  $U \rightarrow V \rightarrow X \rightarrow Z$ . Then from commutativity of the lower square, we have that  $W \rightarrow X \rightarrow Z$  is the same as  $W \rightarrow Y \rightarrow Z$ , so  $U \rightarrow V \rightarrow X \rightarrow Z$  is the same as  $U \rightarrow W \rightarrow Y \rightarrow Z$ , proving commutativity of the outer rectangle. ■

**Exercise 1.30.** Given morphisms  $X_1 \rightarrow Y$ ,  $X_2 \rightarrow Y$ , and  $Y \rightarrow Z$ , show there is a natural map  $X_1 \times_Y X_2 \rightarrow X_1 \times_Z X_2$ , assuming these fibered products exist.

*Proof.* By composing the maps  $X_i \rightarrow Y \rightarrow Z$ , we get two maps with which we can construct the fibered product  $X_1 \times_Z X_2$ . We then note that the projection maps  $X_1 \times_Y X_2 \rightarrow X_i$  satisfy the condition that their composition with the maps  $X_i \rightarrow Y \rightarrow Z$  are the same, so we get a map  $X_1 \times_Y X_2 \rightarrow X_1 \times_Z X_2$ . ■

**Exercise 1.31 (The magic diagram).** Given maps  $X_i \rightarrow Y$  and  $Y \rightarrow Z$ , show that the following diagram commutes, assuming all relevant fibered products exist

$$\begin{array}{ccc} X_1 \times_Y X_2 & \longrightarrow & X_1 \times_Z X_2 \\ \downarrow & & \downarrow \\ Y & \longrightarrow & Y \times_Z Y \end{array}$$

*Proof.* We first do some bookkeeping to figure out what each map is. The map  $X_1 \times_Y X_2 \rightarrow X_1 \times_Z X_2$  is the map discussed in the exercise above. The map  $X_1 \times_Y X_2 \rightarrow Y$  is either  $\alpha \circ \pi_{X_1}$  or  $\beta \circ \pi_{X_2}$ , which are the same by the definition of the fibered product, where  $\alpha$  and  $\beta$  are the maps  $X_i \rightarrow Y$ , the map  $X_1 \times_Y X_2 \rightarrow X_2 \rightarrow Y \times_Z Y$  is the one induced by the composition of the projection maps with  $\alpha$  and  $\beta$ , and the map  $Y \rightarrow Y \times_Z Y$  is the one obtained by factoring the identity map  $\text{id}_Y$  through the fibered product.


Then to prove that the diagram commutes, we appeal to the previous exercise, by providing two squares that commute. Consider the diagram

$$\begin{array}{ccc} X_1 \times_Y X_2 & \longrightarrow & X_1 \times_Z X_2 \\ \downarrow & & \downarrow \\ X_1 & \longrightarrow & X_1 \\ \downarrow & & \downarrow \\ Y & \longrightarrow & Y \times_Z Y \end{array}$$

Where the middle map  $X_1 \rightarrow X_1$  is the identity map. Then we want to show both of the squares commute. For the upper square, we note that the map  $X \times_Y X_2 \rightarrow X_1 \times_Z X_2$  is the one obtained by factoring the



projection maps  $X_1 \times_Z X_2 \rightarrow X_i$ , which gives the following commutative diagram



which gives us commutativity of the upper square. For the lower square, we note that the identity map on  $Y$  gives us a map  $Y \rightarrow Y \times_Z Y$ , and that the map  $X_1 \rightarrow Y$  trivially commutes with the identity map, so it gives us a map  $X_1 \rightarrow Y \times_Z Y$ , so mapping  $X_1 \rightarrow Y \rightarrow Y \times_Z Y$  is the same as going straight  $X_1 \rightarrow Y \times_Z Y$ . ■

A lot of constructions of objects in category theory come from taking the universal property of another object, and flipping all of the arrows to get a “dual” object. To reflect this, we usually take the old object and slap “co-” to the front<sup>6</sup>.

**Definition 1.32.** Define the *coproduct* by taking the universal property for the product and flipping all of the arrows. That is, given two objects  $X, Y$ , the coproduct, denoted  $X \amalg Y$ , is another object equipped with maps  $X \rightarrow X \amalg Y$  and  $Y \rightarrow X \amalg Y$  such that for any object  $C$  with maps  $X \rightarrow C$  and  $Y \rightarrow C$ , there exists a unique map  $X \amalg Y \rightarrow C$  such that the following diagram commutes



**Definition 1.33.** Let  $X, Y, Z$  be objects, and we are given maps  $Z \rightarrow X$  and  $Z \rightarrow Y$ , then the fibered coproduct  $X \amalg_Z Y$  is an object equipped with maps  $X \rightarrow X \amalg_Z Y$  and  $Y \rightarrow X \amalg_Z Y$  such that given another object  $W$  with maps  $X \rightarrow W$  and  $Y \rightarrow W$  such that their compositions  $Z \rightarrow W$  agree, the map factors through to a map  $X \amalg_Z Y \rightarrow W$ , such that the following diagram commutes



**Exercise 1.34.** Show that the disjoint union is the coproduct in  $\mathbf{Set}$ .

*Proof.* Let  $X$  and  $Y$  be sets, and let  $X \sqcup Y$  be the disjoint union, which is equipped with the inclusion maps  $X \hookrightarrow X \sqcup Y$  and  $Y \hookrightarrow X \sqcup Y$ . Then given any maps  $f : X \rightarrow Z$  and  $g : Y \rightarrow Z$ , we get a map  $X \sqcup Y \rightarrow Z$  given by  $f$  and  $g$  on the disjoint components  $X$  and  $Y$ , showing that the disjoint union satisfies the universal property. ■

**Exercise 1.35.** Suppose we have ring homomorphisms  $A \rightarrow B$  and  $A \rightarrow C$  (giving  $B$  and  $C$  the structure of  $A$ -modules). From a previous exercise.  $B \otimes_A C$  has a ring structure. Show that there is a ring homomorphism

<sup>6</sup>What is coffee? It's ffee with all the arrows reversed!

$B \rightarrow B \otimes_A C$  defined by  $b \mapsto b \otimes 1$  (which also exists for  $C$ ). Show this gives a fibered coproduct in  $\text{Ring}$ , i.e. the diagram

$$\begin{array}{ccc} B \otimes_A C & \longleftarrow & C \\ \uparrow & & \uparrow \\ B & \longleftarrow & A \end{array}$$

satisfies the universal property of a fibered coproduct.

*Proof.* We first verify that the mapping  $b \mapsto b \otimes 1$  is a ring homomorphism. By the bilinearity of  $\otimes$ , we get that  $(b_1 + b_2) \otimes 1 = b_1 \otimes 1 + b_2 \otimes 1$ , and that  $rb \otimes 1 = r(b \otimes 1)$ , so this is a ring homomorphism. The same proof goes for the map  $C \rightarrow B \otimes_A C$ .

Then let  $R$  be another ring with ring homomorphisms  $\varphi : B \rightarrow R$  and  $\psi : C \rightarrow R$  such that the compositions with the maps  $A \rightarrow B$  and  $A \rightarrow C$  respectively give the same map  $A \rightarrow W$ . Then we get a map  $B \otimes_A C \rightarrow R$ , which is uniquely defined by  $b \otimes c \mapsto \varphi(b)\psi(c)$  and extending linearly to sums, so  $B \otimes_A C$  is the fibered coproduct in the category of rings. ■

There is also a generalization we have for injective and surjective maps, which are monomorphisms and epimorphisms.

**Definition 1.36.** A morphism  $\pi : X \rightarrow Y$  is a monomorphism, if given two maps  $f, g : Z \rightarrow X$  satisfying  $\pi \circ f = \pi \circ g$ , then  $f = g$ .

You should think of this as an injective map, but note that monomorphisms don't necessarily have to be injective.

**Exercise 1.37.** Show the composition of two monomorphisms is a monomorphism.

*Proof.* Let  $\varphi : X \rightarrow Y$  and  $\psi : Y \rightarrow Z$  be monomorphisms, and let  $f, g : A \rightarrow X$  satisfying

$$\psi \circ \varphi \circ f = \psi \circ \varphi \circ g$$

Since  $\psi$  is a monomorphism, we conclude that  $\varphi \circ f = \varphi \circ g$ , and since  $\varphi$  is a monomorphism, we conclude that  $f = g$ . Therefore, we have that  $\psi \circ \varphi$  is a monomorphism as well. ■

**Exercise 1.38.** Prove a morphism  $\pi : X \rightarrow Y$  is a monomorphism if and only if the fibered product  $X \times_Y X$  exists, and the induced morphism  $X \rightarrow X \times_Y X$  is an isomorphism

*Proof.* We note that if we show that the map  $X \rightarrow X \times_Y X$  is an isomorphism, we have shown existence, since  $X$  certainly exists as an object. Therefore, we can assume that the fibered product exists, and comes equipped with a map  $p : X \times_Y X \rightarrow X$ . In addition, we have that the identity map on  $X$  induces a morphism  $\iota : X \rightarrow X \times_Y X$ . We claim that these two maps are inverses. We have that  $\pi \circ \text{id}_X = \pi \circ p \circ \iota$ , from the universal property of the fibered product, so we conclude that  $\text{id}_X = p \circ \iota$ . We now want to show that  $\iota \circ p = \text{id}_{X \times_Y X}$ . We note that  $p \circ \iota \circ p = \iota \circ p$  using the fact that  $\pi$  is a monomorphism like we did earlier, so  $\iota \circ p$  must be identity.

Conversely, suppose we are given that the map  $\iota : X \rightarrow X \times_Y X$  is an isomorphism. Then any map  $Z \rightarrow X$  induces a map  $Z \rightarrow X \times_Y X$ . Then suppose we are given two maps  $f, g : Z \rightarrow X$  such that  $\pi \circ f = \pi \circ g$ . Then we have that they induce the same map  $Z \rightarrow X \times_Y X$ , which when composing with the isomorphism  $p = \iota^{-1}$ , gives us that  $f = g$ , so  $\pi$  is a monomorphism. ■

**Exercise 1.39.** Show that if  $Y \rightarrow Z$  is a monomorphism, then the induced map  $X_1 \times_Y X_2 \rightarrow X_1 \times_Z X_2$  is an isomorphism.

*Yikes.* As per the hint, there are two ways to prove this. For the first way, let  $V$  be any object. We want to give a bijection  $\text{Hom}(V, X_1 \times_Y X_2) \rightarrow \text{Hom}(V, X_1 \times_Z X_2)$ . By the universal property, we know that giving a map  $V \rightarrow X_1 \times_Y X_2$  is the same as giving maps  $V \rightarrow X_i$  such that their compositions with the maps  $X_i \rightarrow Y$  agree, and likewise for maps  $V \rightarrow X_1 \times_Z X_2$ . We have that the maps that define the fibered product  $X_1 \times_Y X_2$  are the compositions of the maps to  $Y$  with the map  $Y \rightarrow Z$ , so we have that any map  $V \rightarrow X_i$  that satisfies the conditions for  $Y$  will also satisfy the conditions for  $Z$ , so we have that the map  $\text{Hom}(V, X_1 \times_Y X_2) \rightarrow \text{Hom}(V, X_1 \times_Z X_2)$  given by composing with the map  $Y \rightarrow Z$  is surjective. In addition, since  $Y \rightarrow Z$  is a monomorphism, if two maps  $V \rightarrow X_i$  composed with  $\alpha$  or  $\beta$  agree on the compositions

with  $Y \rightarrow Z$ , they induce the same map to  $X_1 \times_Y X_2$ , so this map is injective, so we get a natural bijection  $\text{Hom}(V, X_1 \times_Y X_2) \rightarrow \text{Hom}(V, X_1 \times_Z X_2)$ . We then note that the map  $X_1 \times_Y X_2 \rightarrow X_1 \times_Z X_2$  is induced by the projection maps  $X_1 \times_Y X_2 \rightarrow X_i$ , and by how we constructed the bijection, we have that the bijection  $\text{Hom}(V, X_1 \times_Y X_2) \rightarrow \text{Hom}(V, X_1 \times_Z X_2)$  is given by post-composing with our map  $X_1 \times_Y X_2 \rightarrow X_1 \times_Z X_2$ , reading a little further to Yoneda's lemma tells us that it must be an isomorphism. ■