

# Sum rule for the spin-dependent structure function $b_1(x)$ for spin-one hadrons

F. E. Close

*Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, England,\*  
Department of Physics and Astronomy, University of Tennessee, Knoxville, Tennessee 37996,  
and Physics Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831*

S. Kumano

*Department of Physics and Astronomy, University of Tennessee, Knoxville, Tennessee 37996,  
Physics Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831,  
and Nuclear Theory Center, Indiana University, Bloomington, Indiana 47408-0768\**

(Received 29 May 1990)

We show that the spin-dependent structure function of spin-one hadrons,  $b_1(x)$ , is related to the electric quadrupole moment of the target, and obtain  $\int dx b_1(x) = \lim_{(t \rightarrow 0)} -\frac{5}{3}(t/4M^2)F_Q(t) = 0$  for isoscalar targets if the sea of quarks and antiquarks is unpolarized. We show how this sum rule is modified in the presence of a polarized sea.

The recent measurement by the European Muon Collaboration<sup>1</sup> of the spin-dependent structure function of the proton [ $g_1(x)$ ] has stimulated intense interest in the details of spin structures in the proton and neutron.<sup>2</sup> Measurements of  $g_1(x)$  for the neutron are planned<sup>3</sup> and will require the existence of polarized nuclear targets, such as the deuteron. Nuclear targets with  $J \geq 1$ , such as the deuteron, are also interesting in their own right.<sup>4</sup> In particular there is a new effect which does not exist for spin- $\frac{1}{2}$  hadrons, namely, the existence of a further spin-dependent structure function  $b_1(x)$  which could be measured by polarizing the spin-one target. For real photons, this structure function is essentially that discussed by Pais<sup>5</sup> in 1967.

The only available fixed targets with  $J \geq 1$  are nuclei, and so early discussions of  $b_1(x)$  have tended to be based upon models where the nucleus consists of nonrelativistic nucleons. For nucleons in an  $S$  state,  $b_1(x) \equiv 0$ . For nucleons in a  $D$  state,<sup>4,6</sup>  $b_1(x) \neq 0$  in general. However, we note that these models have the property  $\int dx b_1(x) = 0$ . We find that in a quark-parton model this sum rule is generally true if the sea of quarks and antiquarks is unpolarized. After completing this work, we learned that Mankiewicz<sup>7</sup> has studied  $b_1(x)$  for the  $\rho$  meson and noticed empirically that  $\int dx b_1(x) = 0$  in his model. He noticed also that this need not be the case on the light plane where the probability for the  $q\bar{q}$  to have  $S_z = +1$  is independent of that for  $S_z = 0$ . Our sum rule confirms this and quantifies the effect of sea polarization.

The lepton scattering cross section from a hadron target involves the hadron tensor shown in Fig. 1(a):

$$W_{\mu\nu}(p, q, H_1, H_2) = \frac{1}{4\pi} \int d^4\xi e^{iq \cdot \xi} \langle p, H_2 | [J_\mu(\xi), J_\nu(0)] | p, H_1 \rangle, \quad (1)$$

where the  $H_1$  and  $H_2$  are  $z$  components of the target spin.

Parity and time-reversal invariances lead to eight independent structure functions. We may write a general expression for  $W_{\mu\nu}$  of a spin-one hadron by considering current conservation<sup>4</sup>

$$\begin{aligned} W_{\mu\nu} = & -F_1 g_{\mu\nu} + F_2 \frac{p_\mu p_\nu}{v} - b_1 r_{\mu\nu} \\ & + \frac{1}{6} b_2 (s_{\mu\nu} + t_{\mu\nu} + u_{\mu\nu}) + \frac{1}{2} b_3 (s_{\mu\nu} - u_{\mu\nu}) \\ & + \frac{1}{2} b_4 (s_{\mu\nu} - t_{\mu\nu}) + i \frac{g_1}{v} \epsilon_{\mu\nu\lambda\sigma} q^\lambda s^\sigma \\ & + i \frac{g_2}{v^2} \epsilon_{\mu\nu\lambda\sigma} q^\lambda (p \cdot q s^\sigma - s \cdot q p^\sigma), \end{aligned} \quad (2)$$




FIG. 1. (a) The imaginary part of the forward virtual Compton amplitude and (b) elastic form factor in the quark-parton model.

where

$$r_{\mu\nu} = \frac{1}{\nu^2} (q \cdot E^* q \cdot E - \frac{1}{3} \nu^2 \kappa) g_{\mu\nu}, \quad (3)$$

$$s_{\mu\nu} = \frac{2}{\nu^2} (q \cdot E^* q \cdot E - \frac{1}{3} \nu^2 \kappa) \frac{p_\mu p_\nu}{\nu}, \quad (4)$$

$$t_{\mu\nu} = \frac{1}{2\nu^2} (q \cdot E^* p_\mu E_\nu + q \cdot E^* p_\nu E_\mu + q \cdot E p_\mu E_\nu^* + q \cdot E p_\nu E_\mu^* - \frac{4}{3} \nu p_\mu p_\nu), \quad (5)$$

$$u_{\mu\nu} = \frac{1}{\nu} (E_\mu^* E_\nu + E_\nu^* E_\mu + \frac{2}{3} M^2 g_{\mu\nu} - \frac{2}{3} p_\mu p_\nu), \quad (6)$$

$$s^\sigma = -\frac{i}{M^2} \epsilon^{\sigma\alpha\beta\tau} E_\alpha^* E_\beta p_\tau, \quad (7)$$

and  $\nu$  and  $\kappa$  are defined by  $\nu = p \cdot q$  and  $\kappa = 1 + M^2 Q^2 / \nu^2$ .  $E^\mu$  is the polarization of the target and it is normalized as  $E^2 = -M^2$ .

The helicity amplitudes are defined by

$$A_{h_1 H_1, h_2 H_2} = \epsilon_{h_2}^{\mu} \epsilon_{h_1}^{\nu} W_{\mu\nu}(p, q, H_1, H_2), \quad (8)$$

where  $\epsilon_h^\mu$  is the photon polarization vector with helicity  $h$ . The relations between the structure functions  $b_1(x)$  and  $F_1(x)$  in the scaling limit and these helicity amplitudes are

$$F_1(x) = \frac{1}{3} (A_{+,0,+0} + A_{+,+,++} + A_{+,-,+}), \quad (9a)$$

$$b_1(x) = A_{+,0,+0} - \frac{A_{+,+,++} + A_{+,-,+}}{2}. \quad (9b)$$

Measurement of  $b_1(x)$  requires that the target be polarized.

In the parton model, we define  $q_\uparrow^m(x)$  [ $q_\downarrow^m(x)$ ] as the probability to find a quark with momentum fraction  $x$  and spin up [down] in the spin-one hadron with the  $z$  component of spin  $m$  moving with infinite momentum along the  $z$  axis. This gives

$$F_1(x) = \frac{1}{2} \sum_i e_i^2 [q_i(x) + \bar{q}_i(x)], \quad (10a)$$

$$b_1(x) = \sum_i e_i^2 [\delta q_i(x) + \delta \bar{q}_i(x)], \quad (10b)$$

where  $q_i(x)$  and  $\delta q_i(x)$  are defined by

$$q_i(x) = \frac{2}{3} [q_{\uparrow i}^0(x) + q_{\uparrow i}^1(x) + q_{\downarrow i}^1(x)], \quad (11a)$$

$$\delta q_i(x) = q_{\uparrow i}^0(x) - \frac{q_{\uparrow i}^1(x) + q_{\downarrow i}^1(x)}{2}, \quad (11b)$$

and analogously for  $\bar{q}_i(x)$  and  $\delta \bar{q}_i(x)$ .

To form sum rules in the parton model, one may calculate the dependence of amplitudes on Bjorken  $x$  for a spin-one target moving fast in the  $z$  direction, Fig. 1(a), and compare with static properties integrated over  $x$  at zero-momentum transfer, Fig. 1(b). For example, in the spin-averaged case one has the Gottfried sum rule

$$\int dx [F_1^p(x) - F_1^n(x)] = \frac{1}{6} [F_C^p(0) - F_C^n(0)] = \frac{1}{6}, \quad (12)$$

where  $F_C(0)$  is the charge of the target, which follows if

the number of  $\bar{u}$  and  $\bar{d}$  in the sea are equal (see Ref. 8). Analogously one can form a sum rule for the deuteron

$$\int dx F_1^d(x) = \frac{5}{6} F_C^d(0) + \frac{1}{18} (Q + \bar{Q})_s, \quad (13)$$

where  $F_C^d(0)$  is the deuteron's charge and  $(Q + \bar{Q})_s$  is the number of charge weighted partons in the sea

$$(Q + \bar{Q})_s \equiv 5(U + \bar{U} + D + \bar{D})_s + 2(S + \bar{S})_s. \quad (14)$$

This quantity is infinite and so the sum rule (13) has no utility, but we exhibit it in order to facilitate comparison with our sum rule for  $b_1(x)$ .

We may rederive these familiar spin-averaged sum rules, keeping target polarizations explicit throughout, and thereby immediately see the relation to our new sum rule. Defining [Fig. 1(b)]

$$\Gamma_{H,H} = \langle p, H | J_0(0) | p, H \rangle, \quad (15)$$

then in the parton model we have

$$\begin{aligned} \frac{1}{3} (\Gamma_{00} + \Gamma_{11} + \Gamma_{-1-1}) &= \sum_i e_i \int dx q_{i,v}^A(x) \\ &= \frac{A}{6} \int dx [u_v(x) + d_v(x)], \end{aligned} \quad (16a)$$

$$\begin{aligned} \frac{1}{2} (\Gamma_{00} - \Gamma_{11}) &= \sum_i e_i \int dx \delta q_{i,v}^A(x) \\ &= \frac{A}{6} \int dx [\delta u_v(x) + \delta d_v(x)], \end{aligned} \quad (16b)$$

where the subscript  $v$  refers to the valence quarks, and where we restrict ourselves to  $I=0$  targets. The  $\Gamma_{HH}$  amplitudes in Eqs. (15) and (16) are related to the electric charge and electric quadrupole form factors as<sup>9-11</sup>

$$\Gamma_{00} = \lim_{t \rightarrow 0} \left[ F_C(t) - \frac{t}{3M^2} F_Q(t) \right], \quad (17a)$$

$$\Gamma_{11} = \Gamma_{-1-1} = \lim_{t \rightarrow 0} \left[ F_C(t) + \frac{t}{6M^2} F_Q(t) \right], \quad (17b)$$

where the form factors  $F_C$  and  $F_Q$  are measured in the units of  $e$  and  $e/M^2$ . Thus,

$$\frac{1}{3} (\Gamma_{00} + \Gamma_{11} + \Gamma_{-1-1}) = F_C(0), \quad (18a)$$

$$\frac{1}{2} (\Gamma_{00} - \Gamma_{11}) = \lim_{t \rightarrow 0} \left[ -\frac{t}{4M^2} F_Q(t) \right] = 0. \quad (18b)$$

Substituting Eq. (18a) into Eq. (16a) and Eqs. (10a) and (11a) for isoscalar targets leads to Eq. (13). In a similar way, we use Eqs. (18b), (16b), (10b), and (11b) to obtain a polarized analog of Eq. (13):

$$\begin{aligned} \int dx b_1^d(x) &= \lim_{t \rightarrow 0} \left[ -\frac{5}{3} \frac{t}{4M^2} F_Q(t) \right] + \frac{1}{9} (\delta Q + \delta \bar{Q})_s \\ &= \frac{1}{9} (\delta Q + \delta \bar{Q})_s. \end{aligned} \quad (19)$$

Thus we see that the sum rule for the structure function  $b_1(x)$  is closely related to the electric quadrupole struc-

ture of the target and that the integral vanishes in any model with an unpolarized sea.

This quark-model sum rule provides insight into the  $b_1(x)$  calculated in various models. Models involving nu-

cleons alone, even in a  $D$  state, must give vanishing  $\int dx b_1(x)$  since there need be no nonzero  $\delta Q_s, \delta \bar{Q}_s$ . This can be verified explicitly by integrating the equations given in Refs. 4 and 6:

$$\int dx b_1(x) = \sum_{k=p,n} \left[ \sin^2 \alpha \int dy \Delta f_{dd}(y) - 4\sqrt{2/5} \sin \alpha \cos \alpha \int dy \Delta f_{sd}(y) \right] \int dz F_1^k(z) = 0, \quad (20)$$

because  $\int dy \Delta f_{dd}(y) = 0$  and  $\int dy \Delta f_{sd}(y) = 0$ .

Models with  $\pi$  exchange generate a tensor force and thereby a quadrupole moment for the target. However, with vanishing  $\delta Q$  they still preserve  $\int dx b_1(x) = 0$  as can be seen by inspection of the explicit  $b_1(x)$  in Refs. 4 and 6. Models involving  $\rho$  exchange could give a nonvanishing integral as  $\rho$  can effectively transport a nonzero  $\delta Q$ . This is essentially noted in Ref. 7 where in the light cone, or in  $SU(6)_W$ , the longitudinal and transverse  $\rho$  components are, in principle, independent and hence  $\delta Q + \delta \bar{Q}$  need not vanish.

We are indebted to Dr. L. Mankiewicz and participants at the U.S. Nuclear Theory Institute for discussions. We thank Dr. J. B. McGrory and A. S. Tate for reading the manuscript. This research was supported in part by the U.S. Department of Energy under Contract No. DE-AC05-84OR21400 with Martin Marietta Energy Systems, Inc., and by the State of Tennessee Science Alliance Center under Contract No. R01-1061-68. S. K. was also supported by the U.S. Department of Energy under Contract No. DE-FG02-87ER40365.

\*Present address.

<sup>1</sup>EMC Collaboration, J. Ashman *et al.*, Phys. Lett. B **206**, 364 (1988).

<sup>2</sup>F. E. Close, in *Few Body Problems in Physics*, proceedings of the Twelfth International Conference, Vancouver, Canada, 1989, edited by H. W. Fearing [Nucl. Phys. A**508**, 413c (1990)], for a summary of recent work.

<sup>3</sup>HERMES Collaboration (unpublished).

<sup>4</sup>P. Hoodbhoy, R. L. Jaffe, and A. Manohar, Nucl. Phys. B**312**, 571 (1989).

<sup>5</sup>A. Pais, Phys. Rev. Lett. **19**, 544 (1967).

<sup>6</sup>G. A. Miller, in *Electronuclear Physics with Internal Targets*, proceedings of the Topical Conference, Stanford, California, 1989, edited by R. G. Arnold (World Scientific, Singapore, 1989).

<sup>7</sup>L. Mankiewicz, Phys. Rev. D **40**, 255 (1989).

<sup>8</sup>F. E. Close, *Introduction to Quarks and Partons* (Academic, New York, 1979).

<sup>9</sup>M. Gourdin, Nuovo Cimento **28**, 533 (1963).

<sup>10</sup>F. E. Close, Phys. Lett. **65B**, 55 (1976).

<sup>11</sup>S. Kumano, Phys. Lett. B **214**, 132 (1988); Nucl. Phys. A**495**, 611 (1989), with modifications in Clebsch-Gordan coefficients.