

The following response is done with the material that was used to investigate the beam and track efficiency at the time it was presented to the g12 working group. There is no other means of supplying more information in an efficient time frame.

1 Beam Energy Corrections

1. “Clarify if the photon-energy corrections derived and reported in the note were obtained after the application of the phi-dependent momentum corrections reported in 3.3.1. This correction should be derived after e-loss and phi-dependent momentum corrections are applied.”

Stated on page 75 of the note procedures:

”To validate the corrections of the entire g12 data set, the missing neutron mass was recalculated for each run, shown in Fig. 72, using several correction schemes, i.e. a scheme of just ”energy-loss“ corrections, a scheme of ”energy-loss“ and momentum corrections (JTG PCor), a scheme of ”energy-loss“, momentum corrections (JTG PCor) and beam corrections (MK BeamCor) and a scheme of ?energy-loss? and beam corrections (MK BeamCor). It can be seen in Fig. 72 that the only scheme that sufficed was the combination of ”energy-loss“ and beam corrections.“

However, the plot was modified, making the above statement not understandable. In Figure 1 below shows the plot that this text is referring to. It should be noted that one method of deriving the momentum corrections before the beam corrections did not improve either the neutron mass measurement or the proton mass measurement. This can be seen in Figure 1 by the red data points. Therefore the beam corrections were derived after e-loss but before the phi-dependent momentum corrections. It also should be noted that after the beam corrections were derived and after the phi-dependent momentum corrections were derived, the beam correction algorithm was re-ran to investigate any possible systematic, however the ratio of the largest derivation, for a run, was on the order of 10^{-4} which had only a 0.3 MeV increase at 1 GeV photon beam energy.

2. “Some of us are concerned that the photon energy correction may absorb any remaining systematic biases in the pion momenta. In such a case, the method by which the photon energy correction is derived ensures that the missing mass peaks are placed right at the nominal mass of the missing particle, but the calculated kinematics at which you report your observables (W, Cm angles) will be incorrectly estimated. We would like to see control distributions that provide evidence that after e-loss and phi-dependent momentum corrections, there are no remaining systematic biases in the particles’ momenta.” The observables in which we report (W, C.M. angles) agree well with previous measurements of CLAS and other reputable institutions.

- “Plot the $\pi^+\pi^-$ invariant mass (Ks) as a function of the π^+ and π^- momenta. On each plot show with a solid line the mean of the integrated invariant-mass distribution.”






Figure 1: Plot showing the series of analyses that contributed to the final chain of procedures for the beam correction.



(a)




(b)


Figure 2: (Top): K_s mass vs. π^+ momentum. (Bottom): K_s mass vs. π^- momentum.

- “Plot the $\pi^+\pi^-$ invariant mass (Ks) as a function of the π^+ and π^- theta lab angles. On each plot show with a solid line the mean of the integrated invariant-mass distribution.”


Mass of K0




of Entries




of Entries




Difference from mass of K0




of Entries



Difference from mass of K0



of Entries



For some bins of θ and ϕ the K_s mass difference (from PDG value) are more than 1 MeV, this is due to the number of entries and the fit performed for that bin. See Appendix for fits.

- “Plot the $\pi^+\pi^-$ invariant mass (Ks) as a function of the π^+ and π^- common Z vertex. On the plot show with a solid line the mean of the integrated invariant-mass distribution.”

For all plots involving the K_0 mass distribution, a distance of closest approach cut was placed on the $\pi^+\pi^-$.

- “Plot the $\pi^+\pi^-$ invariant mass (Ks) as a function of run number. On the plot show with a solid line the mean of the integrated invariant-mass distribution.”

This can be seen in figure 62, where the top panel is for run 56515 and the bottom panel is for run 57130, which depicts a plot of the Ks mass being 1 MeV in difference. Since, the proton mass from 56515 and 57130 was 0.930 and 0.939 respectively it was determined that the beam was the culprit.

2 Single Particle Efficiency

1. “The only independent (of analysis) g12 argument at this time that can be acceptable in my view about the validity of the correction would be that after the corrections, the omega cross sections from three-track and two-track events are consistent with each other, but I did not see a comparison of these in MKs thesis”

In section 4.13.4 of MK thesis, there depicts a comparison of several π^0 differential cross-sections from g12 using the dynamic g12 correction compared to the g11 method derived for the g11 experiment. Shown is a good agreement between the 2 methods, however there is slight deviations in the forward $\cos\theta$ direction. We attribute this deviation to the difference in the method of calculating the efficiency, where g12 is dynamic, while the g11 correction was a static “global” correction. Furthermore, in the analysis note procedure, the π^0 differential cross sections are shown to be in good agreement with previous measurements obtained from CLAS and other world measurements.

2. “According to MK thesis, the kinematic fit was done after e-loss and beam-energy corrections. Phi-dependent momentum correction was not done. This is not consistent with the analysis chain in the g12 latest note.”

It is not explicitly stated, however when kinematic fit is performed, the phi-dependent momentum corrections are performed (Prior to the fit, but in the fit procedure).

3. “The parameters of the pull distributions of the simulated and the real data are not consistent with each other within their reported uncertainties. (The error matrices for the simu- and real-data fits seem to have been tuned independently. I wonder how the error matrices compare with each other). I suspect that one of the reasons for that discrepancy are residual differences between the resolutions of the simulation and the data. These

would not be necessarily caught by comparing integrated distributions, such as invariant and missing masses, but would become prominent as one "zooms" in a particular bin of (p,theta,phi, z)."

I require clarification on how the data and simulated pull distributions are not consistent with each other. From MK's thesis, page 127, the pull distributions are shown and are consistent with each other. It is true that there exists two tunes, one for MC and one for data. Each tune has a flag to tell the fitter if it is MC or data. The independent tune was performed because it was seen that using the residual parametrization from the data for the TBER matrices was not sufficient. However, the simulated widths, momentum, θ and ϕ angles were consistent between MC and data, this can be seen in MK thesis pages 134-139.


4. "For a better estimate of the magnitude of the issue above, one would need to see differences, such as p_calculated-p_measured for the narrow kinematic bin (theta,phi,z) as a function of momentum (etc), plotted together for simulated and real data (mean values, widths)."
see above
5. "The counting of detected particles was done by checking if a real particle was detected with kinematics falling within the kinematic bin of interest. On average, that is not a problem, if one derives an overall correction, since bin migration tends to average out. However, that can be very dangerous if the local resolutions in the simulation and the real detector do not match. Different bin migrations will cause a local bias in the efficiency."
The local resolutions of MC and ata are consistent in the g12 experiment. This is documented in section 5.2 of the analysis note procedure.
6. "The uncertainty of the correction has not been estimated (this may take care of my concern in d) as long as the biases are randomly distributed over the (p,theta,phi,z) bins) as far as I could see. The quoted value of 3% for 3-prong events seems to account for photon multiplicity effects in the correction, but not for inherent effects, such as resolution mismatches simulation-data."
7. "In MK's thesis, the detection efficiencies for p, pi+, and pi- are shown both for the simulation and for the data. While, the simulated results nicely show the location of the coils, the real data do not. A relatively high efficiency is for the regions of the coils for the data, where we expect zero. One explanation would be that in the determination of the thetacosphi and thetasinphi bins, the bins covering the coil areas contain areas both with zero efficiency and with high efficiency. I would like to see if the efficiency for that bin was reported for the actual average of the events' kinematics in that bin, or the for the middle of the bin (which is unacceptable). I also wonder if the procedure of correction evaluation did check that the actual mean values of the kinematic bins, for which the ratio of efficiencies simu/real was formed, did indeed match, or one trusted the middle of the bins. Again, this may not be an issue for observables reported for wide bins, but brings into question potential errors of the correction (i.e. systematic biases)." It should be realized that all analyses should do geometric fiducial cuts, these cuts will eliminate any events seen in the

coils of dead space of the DC's. The evaluation done on the track efficiency was to only show the over simulation of events.


3 Appendix


Plots involving K_s mass in bin of π^+ angles

PhiThetaMassPip




PhiThetaMassPip







PhiThetaMassPip




PhiThetaMassPip




PhiThetaMassPip




PhiThetaMassPip




PhiThetaMassPip




PhiThetaMassPip




PhiThetaMassPip




PhiThetaMassPip




PhiThetaMassPip




PhiThetaMassPip




PhiThetaMassPip




PhiThetaMassPip




PhiThetaMassPip




PhiThetaMassPip




PhiThetaMassPip




PhiThetaMassPip




PhiThetaMassPip




PhiThetaMassPip




PhiThetaMassPip




PhiThetaMassPip




PhiThetaMassPip




PhiThetaMassPip




Plots involving K_s mass in bin of π^+ angles




PhiThetaMassPim








PhiThetaMassPim




PhiThetaMassPim









PhiThetaMassPim




PhiThetaMassPim




PhiThetaMassPim







PhiThetaMassPim




PhiThetaMassPim




PhiThetaMassPim




PhiThetaMassPim




PhiThetaMassPim




PhiThetaMassPim




PhiThetaMassPim




PhiThetaMassPim




PhiThetaMassPim




PhiThetaMassPim




PhiThetaMassPim



PhiThetaMassPim





PhiThetaMassPim

