

1. Correction question and g12 answer

- **Tagger and RF calibration Figure 3 shows a jump in resolution. Why is that?**

This is due to the finalization of the production trigger, which requires higher energy photons. As also shown in Fig. 2, the high energy T-counters have better resolutions. With a trigger that requires the presence of a higher energy photon, the overall resolutions changes. However, the individual T-counter timing resolution remains the same.

- **Fig 12 shows good alignment of start counter for entire run period. However the resolution has steps. Any explanations why, for earlier runs, the resolution is a bit worse? Is there an understanding of this effect? How does this affect the data analysis? How does the start counter timing resolution compares to other run periods?**

The ST resolution was obtained by assuming a gaussian distribution, while in reality the ST-RF timign differences are not exactly gaussian. Therefore, it IS affected by background, beam current, trigger. The steps are about 30ps and 10ps respectively, and we use the best start counter hit in our event selection. Therefore, we do not expect any significant effect.

- **ST efficiency. How did you determine this number? Is it for any particle of any momentum at any angle? Have you looked at the efficiency for protons and pions separately? Have you looked at the efficiency for different paddles? For instance infamous paddle 9 has known problems with efficiency and resolution. How is this efficiency used in the analysis?**

The efficiency of the start counter was calculated by examining the number of tracks with a ST hit after track reconstruction. The ST fired 91.6% of the time, this was calculated from run 57000 since it is a good run and included in production data.

We do not account for the bad resolution of paddle 9 specifically. The effect should be absorbed by the track-dependent inefficiency corrections.

- **DC calibrations. Has the DC alignment with B=0 data been done? Show the results.**

Yes, the alignment was done using $B = 0$ data. The note (section 2.5) has been updated accordingly.

- **DC efficiencies. Show a comparison of the real data and MC data with DC efficiency map applied.**

The DC wire efficiencies were computed for each wire using the whole data set, using the CLAS utility PDU. An example of this is shown in the beginning of DC

calibration section. Simulation data all use the GPP to apply this efficiency map. This is standard practice in CLAS. The data compares very well with simulation.

- **There is nothing in the note about CC calibration and efficiency. It is also not clear if CC timing is used in the analysis or not.**

This has been addressed now in the note in Section 2.6.1. Also, the CC timing is not used in standard g12 analysis.

- **EC calibration and efficiency. You don't say anything about energy calibration and energy response. You have quite a few dead strips. They not only affect efficiency but also affect energy response. How did you handle this? What procedures (corrections) were used to reconstruct photon energy and π^0 2gamma decay?**

The EC energy calibration quality can be demonstrated by the two photon invariant mass (π^0) as a function of run number (Fig. 35 in section 2.8). The mean is very close to the expected value. The knock-out of bad and inefficient EC strip are summarized in Table 18.

- **Target thickness. You quote target length uncertainty of 0.2 cm. Please explain where is this number coming from. Have you accounted for the thermal contraction of the cell? At the end, for normalization purposes, you need the number of target protons. Just put this number with all corrections applied. This is the single number everybody should use for this dataset.**

Point taken. We do not quote a separate systematic uncertainty associated only with target length.

- **For yield and PID stability check, and evaluation of the related systematic uncertainty, show the average number of protons and pion per event on a run-by-run basis. The pull distributions of these quantities can be used to estimate the associated systematic uncertainty.**

The average number of protons and pions on a run-by-run basis was monitored during the storage, and we did not observe any significant instability. Furthermore, we believe this is already shown in the omega yield plot in section 4.3. The uncertainty associated with this effect is estimated to be around 0.5 %.

- **For the photon flux systematic uncertainty, evaluate two sources. Comparison of different normalization runs. Pull distribution of normalized yield.**

sections 4.2 and 4.3. plots address this, just need to add a number to the text (sys unc) ????

- **Describe the procedure for photon flux binning to arbitrary bin size.**

Done! (Details in Sec 4).

- **When you compare yields with different beam current you apply a multiple photon correction. There may be different ways of doing it. Explain the procedure.**

For this particular study, only one photon was chosen (randomly) if there is multiple photons in the chosen time bucket. If the probability of having two photons is (ignoring more than two photons) P , the normalized yields is corrected by a factor of $\frac{1}{1-0.5P}$. indeed dependent on the beam current, and is corrected accordingly.

- **When you discuss the photon energy correction, you say that the tagger magnetic field is not measured. This is not correct. We had a Hall probe in the tagger magnet and it was read in EPICS together with magnet current.**

It has been corrected accordingly in the note.

- **Fig. 57 caption says beam current, but shows beam energy.**

It was a typo and has been corrected.

- **Fig. 51 is confusing. What is JTG PCor? Yellow points show the best match to neutron mass, but if the legend is correct the momentum correction is not applied. Please, clearly explain what exact procedure must be used for momentum/energy corrections.**

The g12 momentum correction was derived by Johann, hence the name of JTG. Fig. 51 was indeed confusing, and it has now been replaced (at a different location), to demonstrate the effect of all g12 corrections being applied together (End of Section 3.3, Fig. 68).

- **In addition to missing mass distributions, show some invariant mass distributions. They are independent of photon energy and will show if the momentum corrections are good regardless of photon energy correction.**

The invariant mass of $K^+\pi^0$, showing the K^{*+} , and the K^+K^- invariant mass showing the $\phi(1020)$, have been added in the note (chapter 6). The mass of K^0 reconstructed from $\pi^+\pi^-$ pairs can also be seen to be where it is expected (Fig. 58). All known narrow states in g12 have been shown to be within 1 MeV of the PDG values.

- **Reference for [eloss] is CLAS Note 2007-016**

It has been corrected.

- **You refer to the g9 momentum correction. In general the procedure may be applied to g12 data. However direct comparison does not make sense since the configuration of the experiments were very different.**

We do not use g9 momentum correction. And the note has been modified accordingly.

- **The lepton id is based on the analysis of the EC and CC information, however no discussion of the CC calibration is included in chapter 2.**

This is related to the previously addressed issue.

- **The CC cuts used in the present analysis includes a cut on Npe and a cut on the CC hit angle that is related to the track direction. No cut on the time correlation between CC hit and, for example, TOF hit associated to the same track is used: this is a standard cut used in all recent electron beam analysis used to reject misidentified pion. Why is this not applied?**

Using the current cuts of NPE and hit angle, the suppression of di-leptons was sufficient without including additional cuts on the CC. This method of lepton PID, involving di-leptons, was established during the g7 run period. Further g12 analyses that involve single lepton PID could include this as a cut.

- **The EC cuts used in this analysis are based on the total energy deposited in the calorimeter, E_calor, and exploit the fact that E_calor/p is almost a constant for leptons. This is a typical selection criteria used in most electron run analyses: usually the way this selection is illustrated is by plotting E_calor vs p and displaying the cuts on this graph. We suggest replacing figures 69-75 with these simple plots: this would make easier for a reader, familiar with lepton analysis, to understand the selection cuts**

???

- **The note refers to the g7 lepton id scheme in explaining the chosen cuts: in g7 however, not only cuts on the total calorimeter energy but also cuts on the inner and outer energy were used: why were these not considered necessary in the present analysis?**

IsLepton() in g12 was made to be a balanced between good lepton selection without being aggressively restrictive. Like any other PID, it is up to the individual analyses to show the validity of their particle selection, and make additional cuts as needed.

- **In chapter 2.7.1, the knockout of bad EC strips is explained in details. We assume the loss of efficiency due to the strip knockout was accounted for by applying the same knockout function to both real data and mc data (used for the efficiency evaluation). Can you please confirm this is what was done**

(the explanation in the note is not very clear about this point)? The strip knockout will induce a worsening of the calorimeter energy resolution that becomes dependent on the sector and on the track impact point. How was this accounted for?

The bad strips are knocked out both in data and simulation. The effect of energy resolution worsening is not accounted. If this is an important issue for a particular analysis, it has to be investigated separately.

- **Was the stability of the EC calibration checked as a function of run number (for example monitoring the E_{calo}/p ratio)?**

Not exactly. However, the π^0 mass was monitored as a function of run number and stable.

- **To be able to judge the effectiveness of the lepton id cuts, we suggest showing the CC photoelectron distribution for the initial and final lepton sample: a typical hint of how clean the final sample is the disappearance of the peak at about 2 photoelectrons which is visible in the initial plots.**

In our procedure, which is derived from the g7 group, a cut on the npe is part of the lepton ID. The effectiveness can be demonstrated in the figures in section 3.4.

- **You use the DC residual match to determine smearing parameters. Once this is done, show how the reconstructed momentum resolution compares with the one from simulation.**

While deriving the smearing parameters, we also monitored the TOF timing resolution as well as invariant mass resolutions such as $\phi(1020)$. The data compares very well with the data. For example, the overall TOF resolution is about 215 ps in the simulation, very close to the data which is just above 200 ps.

- **To demonstrate that efficiencies are accounted for correctly, show comparisons of various distributions (angular, occupancies?) between data and simulation.** This has been done for g12 in many topologies. For illustration purpose, the momentum and angular distributions for e^+ are shown here, comparing data and simulation.

- **Did you knock out bad tagger counters in the simulation?**

Two known bad energy regions in the tagger: (2.975,3.175) (dead/bad paddle) and (3.475,3.575). G12 group do not use events from these two regions.

- **If input events for GSIM are in the form MCVX/MCTK banks, GSIM can properly generate the TAGR bank. Among other things it modifies photon energy and assigns it to the center of the Ebin. If you use the PART bank as**




Figure 1: TOF resolution for data (left) and simulation(right)

an input to GSIM, your event generator is supposed to generate the TAGR bank. How it is done in your event generator? Does it account for tagger sag?

The TAGR bank was generated in the same manner in both cases. The tagger sag was accounted in the data during the cooking stage. For simulation, there is no tagger sag.

- **Fiducial Region Selection: Were these determined from simulated events? Why not from real data?**

It is derived from real data.

- **p.87 bottom: Why 50%? How was this criterion chosen? Above 50% the acceptance still varies as fast as below 50% - is the acceptance in regions above 50% better described by GSIM?**

g12 has three sets of fiducial cuts (nominal, tight and loose), not just the 50% (nominal) cuts. One should always try them all and used the difference to derive systematic uncertainties associated with the fiducial cuts. This of course, is applicable only when statistics of the specific topology allows.

- **p.88: We would like to see all the phi distributions - can be given with a link**




Figure 2: Momentum and angular comparison of data to simulation for the positron.

to a web site.

Unfortunately we did not keep all the plots. Once the numbers were extracted after careful examination of those plots, we only kept the numbers and eventual software tools for applying the fiducial cuts.

- **Fig. 83: We would like to see these plots for all sectors. The point at 0.4 GeV/c has a very small uncertainty, while the two neighboring points have much larger uncertainties - is it understood why?**

This is really an artifact of statistics. Even if this point was taken out, the fit would still work.

- **Provide the size of the different skims, how many events?**

total size:

211 TiB (exclusive)
 31 TiB (additional)
 26.2 Ge (Billion Events)

exclusive skims:

size	events	skim
50 TiB	6.2 Ge	1-1ckaon1ctrk
31 TiB	3.8 Ge	2-2pos1neg_not_1ckaon1ctrk
68 TiB	8.4 Ge	3-2ctrk_not_2pos1neg_1ckaon1ctrk
62 TiB	7.7 Ge	4-not_2ctrk_2pos1neg_1ckaon1ctrk
263 GiB	32.6 Me	5-other

additional skims:

size	events	skim
17 TiB	2.1 Ge	6-1lepton
13 TiB	1.6 Ge	7-4ctrk
1 TiB	124 Me	8-ppbar

- **What banks are included in the cooked files? Are the SEB and GPID banks included?**

here they are in alphabetical order:

CC CC01 CCPB CCRC CL01 DCPB DSTC EC EC1 EC1R ECHB ECPB ECPC
ECPI ECS EPIC EVNT FBPM HBER HBID HBTR HEAD HEVT HLS LCPB
MVRT PART RGLK S1ST SC SC1 SCPB SCR SCRC SCS ST1 STN0 STN1
STPB STR STSN SYNC TAGE TAGI TAGR TBER TBID TBTR TDPL TGBI
TGPB TGS TRGS TRKS TRL1

SEB related banks (EVNT etc) are included. GPID is not.

- **Non-physics events are included in in a separate skim. How can one synchronize them with physics events? They have different numbering sequences.**
Events can be coordinated by timestamp; Alternatively, one can always go back to the raw data if someone in the future find it useful to include these non-physics data.
- **Yesterday we found out from Zulkaida's talk that some additional normalization correction of 15% has been applied to cross section. I don't think there is anything said about this correction in g12 note. Is this common correction to be applied to all cross sections? If so it has to be included in the note with explanation how it was obtained.**

That global correction is not technically correct, but it happens to be very close to what should be applied. G12 compares the reconstruction efficiency between real and mc data (ratio of the acceptances for each track). This leads to an overall correction about 5% for each track, but it is slightly kinematical variable dependent. If someone is looking at 2-track events, then the correction would have been smaller.

The g12 procedure for applying this track-dependent inefficiency correction has now been documented (Section 6.1). It is important to point out that this correction naturally absorbs any detector/trigger inefficiency that was not fully accounted for in our procedures such as the GPP smearing.

The following questions are still being worked on:

- **TOF efficiency and bad paddles.** The selection of bad paddles is based on raw ADC and TDC information. This is only good for the initial selection. After that you should look at the PID quality of individual paddles. We know from other data sets that sometimes the occupancy looks good but particle ID is not good. The bad particle ID could be due to bad resolution or sometimes we see double peaking in which the timing is jumping between two different states. If it is stable within the run it is fixable by adjusting offsets run-by-run. However these instabilities appear even within a single file. It would be good to see timelines for vertex time difference and resolutions for protons and pions vs. run number for individual paddles.
- **TOF uses different discriminators with different thresholds for TDCs and for trigger.** It was discovered by the g13 group that some of the counters are inefficient in the trigger but look fine in TDC. What has been done to address this trigger inefficiency?

Being worked on by Rafael. We should be able to show that the resolutions are stable. However, we do not plan to knock out additional paddles, because the track dependent inefficiency correction absorbed this effect. If we were to do a new knockout scheme, everything would have to be changed again and there is nothing concrete we could gain from it.

- **When you compare cross sections to previous measurements please compare in the same kinematical bins.** A strong forward peak pushes other point to the axis and it is not possible to judge if they agree or not. In addition to overlaying two data sets, show their ratio. Even without it we see that in some bins the agreement is not very good. Make something similar to pull distributions as well, to see if there is a bias.

Being done by Zulkaida/MK/Johann.

- **Beam polarization.** While you well describe the study performed to verify helicity assignment, you don't provide an exact recipe for the user on how to obtain the sign of the helicity. Was it direct or delayed reporting of helicity? Should the user include the half-wave plate state or it was already accounted for? Ideally would be good to have a function which takes care of

all this and returns the sign of helicity for a given event. The degree of the photon polarization then will have to be calculated for the 'good' photon using electron beam polarization and polarization transfer. This can be made a standard function which will pull out electron beam polarization for a given run. This will make life easier for those who will analyze this data in the future. Eq. 11 is not correct - need to explain how the photon polarization was implemented so that the yield asymmetry is converted into a polarization observable (event by event, integrated over all events, else?); level of agreement between g12 and g1c control asymmetries must be quantified; Was I_{circ} monitored per run? Why do you not reproduce exactly the same I_{circ} as in [17] and visualize/quantify the level of agreement for exactly the same energy bin. We agree that this method is good to determine the overall sign of the helicity assignment, however there still may be a polarization bias, which is not determinable from this comparison, due to the acceptance effect on the extracted I_{circ} .

Being worked on by Rafael.